



Effect of Mulberry Irrigated with Treated Sewage Water on Silkworms and Cocoon Production

M. Arun Kumar^{a++*}, S. Chandrashekhar^{a#},
C. Seenappa^{b#} and V. S. Ramya^a

^a Department of Sericulture, University of Agricultural Sciences, GKVK, Bengaluru-65, India.

^b Department of Agronomy, University of Agricultural Sciences, GKVK, Bengaluru-65, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i121517

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93760>

Original Research Article

Received: 23/09/2022

Accepted: 19/11/2022

Published: 21/12/2022

ABSTRACT

Aim: The experiment was aimed at evaluating the effect of mulberry irrigated with treated sewage water on silkworms and cocoon production.

Study Design: The experiment was laid out in a Randomized Complete Block Design (RCBD) with six treatments and four replications.

Place and Duration of Study: The study was conducted during Rabi 2019 in pre-established irrigated V-1 mulberry garden at Department of Sericulture, University of Agricultural Sciences, Gandhi Krishi Vigyana Kendra, Bengaluru, Karnataka, India.

Methodology: A field experiment was conducted in a V1 mulberry garden established under paired row system to study the influence of raw and treated sewage water irrigation on growth of mulberry

++Ph. D. student;

Professor;

*Corresponding author: E-mail: m.a.k.arunkumar01@gmail.com;

and subsequent effect on silkworm ($PM \times CSR_2$) growth and cocoon parameters. The raw and treated sewage water was collected from GKVK sewage treatment plant and irrigated to mulberry garden. Silkworms fed with mulberry leaves raised on recommended dose of NPK kg/ha/year + FYM along with raw and treated sewage water irrigation showed positive influence on late age silkworms.

Results: Observations recorded on silkworm growth and cocoon parameters revealed that significant higher fifth instar larval weight (32.02 g/10 larvae), ERR (99.0 %), single cocoon weight (1.706 g), pupal weight (1.385 g), shell weight (0.321 g), shell ratio (18.65%) and shorter fifth instar larval duration (181.91 hrs). Similarly, reeling parameters of cocoons showed significantly longer filament length (853.00 m) and higher filament weight (0.28 g) in silkworms fed with mulberry leaves from the plots irrigated with raw sewage water.

Keywords: Silkworm growth; cocoon parameters; raw sewage water; treated sewage water.

1. INTRODUCTION

"Mulberry foliage is the sole food for the silkworm (*Bombyx mori*, named after the mulberry genus *Morus*. Mulberry is a perennial crop which can be maintained for many years. Selection of land, recommended package of practices and water management are the imperial factors for producing quality leaf. Among these, irrigation water plays a significant role as one of the key inputs in mulberry cultivation. Water in the soil-plant system is a necessary medium for the distribution of nutrients through the plant. It acts as a solvent for biochemical reactions, a medium of distribution for solutes, temperature regulator and as a source of hydrogen in photosynthesis" [1].

Bongale and Krishna [2] evaluated "the effect of raw sewage and borewell water irrigation on leaf quality of mulberry (*Morus indica*) and cocoon crops of silkworm (*Bombyx mori*). The results showed that significantly higher cocoon yields were recorded in raw sewage water irrigated gardens, compared to borewell irrigated mulberry gardens".

Chandrakala et al. [3] conducted "a field survey to study the cocoon yield pattern of the silkworm crops raised on mulberry irrigated with wastewater as against irrigation by bore wells in the vicinity of Bangalore city who irrigate their mulberry gardens with Vrishabhavathy stream water, which is polluted with domestic and industrial wastewater from the city. The average yield for 100 layings was 46 to 57 kg in the control group (borewell water) whereas in test groups (stream water), it ranged between 34 to 51 kg".

Chandrajou et al. [4] reared "Kolar gold, CSR19, and $CSR_2 \times CSR_4$ breeds of silkworm using M5

variety mulberry leaves cultivated by various proportions of spent wash (SW) irrigation. Influence of spent wash irrigated M5 Mulberry leaves on the yield of different breeds of silkworms was evaluated. Results showed that the cocoon yields of CSR19, Kolar gold and $CSR_2 \times CSR_4$ were high in case of mulberry leaves irrigated with 33 per cent spent wash than raw water and 50 per cent distillery spent wash irrigation. The percentage yield was maximum in the case of Kolar gold (26%) and minimum in $CSR_2 \times CSR_4$ (21.87%) and it was moderate in CSR19 (25.6%). This concludes that the diluted 33 per cent distillery spent wash irrigated mulberry leaves is most beneficial for the rearing of the above breeds of the silkworm, which increases the economy of the farmers".

Saad [5] carried out "a study to compare the silkworm performance by feeding mulberry leaves from mulberry gardens irrigated with borewell water and sewage water in spring season. The results revealed that the larvae reared on mulberry leaves from gardens irrigated with borewell water showed the same non-significant performance in larval weights, while larvae fed on leaves from mulberry garden irrigated with sewage water exhibited a significant increase in the mean weight of fresh cocoon, pupae and cocoon shell by about 18.9, 23.9 and 18 per cent respectively which is higher than silkworms reared on leaves from mulberry plants irrigated with borewell water".

In 2014, the Karnataka government envisaged two projects viz., Koramangala-Challaghatta (KC) Valley project and Hebbal-Nagavara (HN) Valley project to supply treated waste water from Bengaluru to irrigation tanks in neighbouring districts of Kolar and Chikkaballapur in the view of recharging groundwater level and making water available for irrigation. In sericulture,

irrigation with various waste waters like silk reeling waste water, distillery spent wash, industrial waste discharge is emphasized which are rich in plants essential nutrients for mulberry growth and development.

As available literature regarding treated sewage water and the effect of treated sewage water on mulberry production and its impact on cocoon crop are very few, the current investigation was taken up with the following objective to study the effect of mulberry irrigated with treated sewage water on silkworms and cocoon production.

2. MATERIALS AND METHODS

The study was conducted during Rabi 2019 in pre-established irrigated V1 mulberry garden at Department of Sericulture, University of Agricultural Sciences, GKVK, Bengaluru, Karnataka, India. The type of soil is clay loam and annual rainfall ranges from 528 mm to 1374.4 mm with the mean of 915.8 mm. The Experiment was laid out in a Randomized Complete Block Design (RCBD) with four replications and six treatments, comprising of different proportions of raw sewage water, treated sewage water and borewell water. The II instar silkworms procured from Basaveshwara chawki rearing center, Devanahalli taluk, Bengaluru and reared in plastic trays by feeding three times a day with chopped tender mulberry leaves of Victory-1 variety. Bed cleaning was done twice and thrice during II and III instars, respectively, by lifting unfed leaves and excreta of the silkworm. Optimum spacing was provided according to the age of the silkworms. After each bed cleaning and when all the silkworms settled for moulting, the lime powder was dusted on silkworms. Shelf rearing was conducted during third, fourth and fifth instar of silkworm by feeding raw sewage water and treated sewage water irrigated V1 mulberry leaves by following standard rearing practices [6].

The treatment details in the silkworm rearing room are presented below.

Silkworm Hybrid	PM × CSR ₂ Kolar gold)
Number of treatments	6
Number of replications	4
Number of worms/treatment/replication	50

The treatment details are presented below.

Treatment details:

- T₁ - 100% borewell water irrigation (control)
- T₂ - 25% treated sewage water + 75% borewell water irrigation
- T₃ - 50% treated sewage water + 50% borewell water irrigation
- T₄ - 75% treated sewage water + 25% borewell water irrigation
- T₅ - 100% treated sewage water irrigation
- T₆ - 100% raw sewage water irrigation

Silkworm growth and cocoon parameters of PM × CSR₂ hybrid: Ten silkworms and ten cocoons were randomly selected in each treatment for recording observations on growth and cocoon parameters such as Vth instar larval weight (g /10 larvae), Vth instar larval duration (hrs), Effective rate of rearing (%), mortality (%), single cocoon weight (g), pupal weight (g), single shell weight (g), shell ratio (%), cocoon filament length (m), non-breakable filament length (NBFL) (m) and denier.

Effective rate of rearing (%): The number of cocoons harvested at the end of the rearing was recorded and the ERR was calculated by using the formula.

$$\text{ERR (\%)} = \frac{\text{Number of cocoons harvested}}{\text{Total number of silkworms reared}} \times 100$$

Mortality (%): The number of silkworms dead at the end of rearing was recorded and the mortality rate (%) was calculated by using the formula.

$$\text{Mortality (\%)} = \frac{\text{Number of silkworms dead}}{\text{Total number of silkworms reared}} \times 100$$

Shell ratio (%): Shell ratio indicates the total quantity of silk available from the single cocoon and is expressed as a percentage. It was calculated by using the following formula.

$$\text{Shell ratio (\%)} = \frac{\text{Weight of the cocoon shell}}{\text{Weight of the whole cocoon}} \times 100$$

Non-breakable filament length (NBFL) (m): It is the average length of the filament that can be unwound from the cocoon without break.

$$\text{Nonbreakable filament length (m)} = \frac{\text{Total filament length}}{1 + \text{Number of breaks in filament}}$$

Denier: The raw silk filament was removed from the Epruvette and weighed after conditioning at room temperature for 24 hrs to determine the denier using the standard formula.

$$\text{Denier} = \frac{\text{Weight of single cocoon filament (g)}}{\text{Length of single cocoon filament (m)}} \times 9000$$

The data recorded on various parameters were subjected to Fisher's method of Analysis of Variance (ANOVA) and interpreted according to Gomez and Gomez [7]. The level of significance used in F and t-tests was $P=0.05$ for RCBD. The critical difference (CD) values were computed where the F test was found significant.

3. RESULTS AND DISCUSSION

Silkworms fed with mulberry leaves from the V1 mulberry variety raised on recommended dose of NPK kg/ha/year + FYM along with raw and treated sewage water irrigation showed positive influence on late age silkworms. However, the different parameters viz., fifth instar larval weight, fifth instar larval duration, effective rate of rearing, mortality percentage, cocoon parameters and reeling parameters showed significant difference among the treatments.

Effect of raw and treated sewage water irrigation on rearing performance of late age silkworm (PM × CSR₂): Raw and treated sewage water irrigation to mulberry significantly increased rearing performance of late age silkworm (PM × CSR₂) (Table 1). Significant higher fifth instar larval weight (32.02 g/10 larvae), ERR (99.0 %) and significantly shorter fifth instar larval duration (181.91 hrs) mortality (1.0%) were recorded when silkworms were fed with mulberry leaves from the plots irrigated with 100% raw sewage water and that were statistically on par with 100% treated sewage water irrigation. The lowest larval weight (26.86 g/10 larvae), longer fifth instar larval duration (191.64 hrs), lowest ERR (93.25%), higher mortality (6.75%) were observed when silkworms were fed with mulberry leaves from 100% borewell water irrigated plots.

Increase in larval weight is apparently due to superior leaf quality. The present investigation recorded higher larval weight, which was in accordance with the results of Chikkaswamy et al. [8] who had recorded higher Vth instar larval weight (53.01 g/10 larvae) in NB₄D₂ larvae fed with leaves from raw sewage treated plot. Paramanik [9] reported that "sewage water had pronounced influence on biosynthesis of carbohydrates and proteins in leaves which help to improve the growth, metabolism and physiological activity of larvae". Similarly, "shorter larval duration may be due to the balanced

nutritional status of the leaves which enable the silkworms to mature early due to the faster metabolic activity" [10]. Rao et al. [11] reported that sewage water had pronounced influence on biosynthesis of three amino acids namely threonine, methionine and histidine which help to improve the growth of the silkworms.

Effect of raw and treated sewage water irrigation on Cocoon parameters of silkworm (PM × CSR₂): The cocoons harvested from silkworms fed with mulberry leaves from the plots irrigated with 100% raw sewage water recorded maximum single cocoon weight (1.706 g), pupal weight (1.385 g), shell weight (0.321 g), cocoon shell ratio (18.65%) and cocoon yield per 100 dfls (74.25 kg) were recorded when silkworms were fed with mulberry leaves from the plots irrigated with 100% raw sewage water (Table 2). The lowest single cocoon weight (1.337 g), pupal weight (1.114 g), cocoon shell ratio (16.56%), shell weight (0.223 g) and the cocoon yield (56.47 kg) were observed when silkworms were fed with mulberry leaves from 100% borewell water irrigated plots.

Same trend was observed by Chandraju et al. [4] who reported increase in cocoon yield of CSR₂ × CSR₄ by 28 per cent with the application of distillery spentwash water. In line with the present observations, Shankar et al. [10] also reported that "feeding of mulberry leaves with higher concentration of nutrients recorded significantly higher pupal weight compared to control. Therefore, the quality of leaves largely determines the performance of silkworms during their development and spinning of cocoons".

Effect of raw and treated sewage water irrigation on cocoon reeling parameters of silkworm hybrid (PM × CSR₂): The filament length (853.00 m), non-breakable filament length (373.25 m), Filament weight (0.28 g) were significantly longer when silkworms fed with V1 mulberry leaves from plots irrigated with 100% raw sewage water. While comparatively shorter filament length (797.75 m), non-breakable filament length (297 m) and lower filament weight (0.22 g) was recorded when silkworms were fed with mulberry leaves from 100% borewell water irrigated plots. The filament denier was non-significant among the different treatments. However, lower denier (2.44) was recorded when silkworms were fed with mulberry leaves from 100% borewell water irrigated plots (Table 3).

The results showed same trend as of Chandraju et al. [4] who recorded significantly higher

filament lengths (800.00 m, 868.35m and 878.34m) in three different breeds of silkworm CSR 18, CSR 19 and Kolar gold, respectively with the application of distillery spentwash as one of the treatments to mulberry. Further he reported that spentwash irrigation results enhanced nutrition in mulberry leaves which in

turn influenced better growth of silkworms, yielding higher proportions of silk proteins and spinning of long silk thread. These results only signify the close relationship between higher nutrient status of the leaf and silkworm nutrition which resulted in higher cocoon parameters.

Table 1. Effect of V1 mulberry leaves on silkworm (PM × CSR2) as influenced by raw and treated sewage irrigation

Treatments	V th instar larval weight (g/10 larvae)	Effective rate of rearing (%)	Mortality (%)	V th instar larval duration (h)
T ₁ (100 % BW)	26.86	93.25	6.75	191.64
T ₂ (25% TSW + 75% BW)	28.62	94.00	6.00	190.80
T ₃ (50% TSW + 50% BW)	29.35	95.25	4.75	188.33
T ₄ (75% TSW + 25% BW)	29.74	96.00	4.00	184.55
T ₅ (100% TSW)	31.26	99.00	1.00	183.50
T ₆ (100% RSW)	32.02	97.50	2.50	181.91
F-test	**	*	*	**
S.Em.±	0.33	1.07	1.07	1.59
CD	1.40	3.21	3.21	6.31

➤ *Significant at 5% and **Significant at 1%.

➤ BW= borewell water, TSW= treated sewage water and RSW= raw sewage water.

Table 2. Effect of V1 mulberry leaves on cocoon parameters of PM × CSR2 as influenced by raw and treated sewage water irrigation

Treatments	Single cocoon weight (g)	Pupal weight (g)	Single shell weight (g)	Cocoon shell ratio (%)	Cocoon yield kg/100 dfls
T ₁ (100 % BW)	1.337	1.114	0.223	16.56	56.47
T ₂ (25% TSW + 75% BW)	1.464	1.216	0.248	16.73	62.15
T ₃ (50% TSW + 50% BW)	1.495	1.234	0.261	17.28	64.12
T ₄ (75% TSW + 25% BW)	1.530	1.253	0.276	17.81	66.15
T ₅ (100% TSW)	1.618	1.321	0.294	17.96	72.19
T ₆ (100% RSW)	1.706	1.385	0.321	18.65	74.25
F-test	**	*	**	**	**
S.Em.±	0.030	0.048	0.009	0.407	0.956
CD	0.131	0.146	0.040	1.228	3.951

➤ *Significant at 5%, and **Significant at 1%.

➤ BW= borewell water, TSW= treated sewage water and RSW= raw sewage water.

Table 3. Silk filament characteristics of PM × CSR2 as influenced by raw and treated sewage water irrigation to V1 mulberry

Treatments	Filament length (m)	Non-breakable filament length (m)	Filament weight (g)	Denier
T ₁ (100 % BW)	797.75	297.00	0.22	2.44
T ₂ (25% TSW + 75% BW)	810.75	310.75	0.23	2.56
T ₃ (50% TSW + 50% BW)	814.25	321.25	0.24	2.64
T ₄ (75% TSW + 25% BW)	821.00	329.75	0.26	2.79
T ₅ (100% TSW)	841.75	366.75	0.27	2.84
T ₆ (100% RSW)	853.00	373.25	0.28	2.93
F-test	**	NS	**	NS
S.Em.±	5.76	-	0.01	-
CD	23.98	-	0.02	-

➤ *Significant at 5%, and **Significant at 1%.

➤ BW= borewell water, TSW= treated sewage water and RSW= raw sewage water.



Plate 1. Vth instar larvae fed with V1 mulberry leaves from plots irrigated with raw sewage water and borewell water



Plate 2. General view of silkworm rearing (PM x CSR₂)

4. CONCLUSION

The Raw sewage water irrigation along with recommended dose of NPK and FYM had positive influence on cross breed silkworm PM x CSR₂ when fed with mulberry leaves. The mulberry irrigated with raw sewage increased the nutrient content of leaves with carbohydrates and proteins which in turn influenced the cocoon parameters positively. Further analyses should

be carried out on the biochemical parameters of silkworm to find out the bio-transportation and bio-accumulation of heavy metals from mulberry to silkworms which affects the silkworm growth.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Subbaswamy MR, Basavanna HM and Suryanarayana N. Quality of irrigation water for mulberry gardens. Indian silk. 1987;47-49.
2. Bongale UD and Krishna M. Leaf quality of mulberry (*Morus indica* L.) and cocoon crops of the silkworm (*Bombyx mori* L.) as influenced by sewage and bore well water irrigation. Indian Journal of Sericulture, 2000;39(2): 165-168.
3. Chandrakala MV, Maribashetty VG, Aftab Ahamed CA and Jyothi HK. Cocoon yield pattern and analysis of water, soil and leaves from mulberry gardens irrigated with polluted water around Bangalore, India. International Journal of Industrial Entomology. 2009;18(2): 97-103.
4. Chandraju S, Nagendraswamy G and Kumar CSC. Yields of *Bombyx mori* L. races cocoons CSR19, Kolar gold and CSR₂xCSR₄ reared fed with V1 variety of mulberry leaves cultivated by spentwash irrigation. Journal of Chemical and Pharmaceutical Research. 2012;4(3): 1812-1814.
5. Saad IA. Comparative study between silk production from mulberry gardens irrigated with clean and sewage water in Egypt. Journal of Agricultural Research, Kafr El-Shaikh University. 2014;40(2): 384-389.
6. Dandin SB, Jayaswal J and Giridhar K. Handbook of Sericulture Technologies. Central Silk Board, Bengaluru. 2001;287.
7. Gomez KA and Gomez AA. Statistical procedures agricultural research, an international rice research institute book. A Willey Inter Science Publication, John Willey and sons. New York, 1984.
8. Chikkaswamy BK and Paramanik RC. Effects of tannery effluents on mulberry genotypes for biomass production. International Journal of Biosciences and Nanosciences. 2014;1(2): 54-62.
9. Paramanik RC. Impact of sewage irrigation on mulberry varieties. International Journal of Advanced Research in Engineering and Applied Sciences. 2015;4(2): 59-71
10. Shankar MA. Nutritional management of mulberry and its effect on silkworm growth, development and silk quality. Ph.D. (Agronomy) Thesis, UAS, Bangalore, 1990; 410.
11. Rao DMR, Munirathnam Reddy M, Gopinath OK and Vindhya GS. Physio-biochemical characterization of two mulberry genotypes under sewage water irrigation. Sericologia. 2011;51(2): 249-258.

© 2022 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/93760>