



Effect of Feldspar, Compost and Biochar on Cultivating Cowpea (*Vigna unguiculata* ssp. *unguiculata*) Plant and Soil Sandy Clay Loam Properties

M. Doaa ^{a*} and A. E. Ashmawi ^b

^a Soils, Water and Environmental Res. Inst., (SWERI), Agriculture Research Center, (ARC) Giza, Egypt.

^b Department of Horticulture, Faculty of Agriculture Alazhar University, Egypt.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ASRJ/2022/v6i130123

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/85385>

Received 02 February 2022

Accepted 05 April 2022

Published 19 April 2022

Original Research Article

ABSTRACT

The crop production in Egypt relies completely on imports to meet its annual requirement of potash fertilizers besides; the high cost of conventional, water soluble K fertilizers constrains their use by most of the farmers in the country. This work was conducted cowpea (*Vigna unguiculata* ssp. *unguiculata*) plants cv. Dokki 126 at Ismailia Research Station, Agricultural Research Center (A.R.C), Egypt, during the two summer seasons of 2020 and 2021 to study the effect of some treatments as potassium feldspar, biochar and compost on soil sandy clay loam properties and the physical and chemical contents of cowpea. A randomized complete block design was used. The results show that the potassium feldspar + biochar + Compost significantly increased the vegetative growth expressed as plant length, number of leaves, number of branches, pod length, Pod diameter, Pod fresh weight, Pod dry weight, Plant fresh weight, Plant dry weight. Also the results indicated that soil pH and EC was not significantly affected in the soil treated with potassium feldspar + biochar + compost. There are positive significant effects on nutrient availability (NPK) in both seasons by using potassium feldspar + biochar + compost in soil. The application of OMW can improve soil quality indices (nutrients (N, P, and K), organic matter, pH, total porosity, bulk density).

*Corresponding author: E-mail: doaa.mousa@gmail.com;

Keywords: Cowpea; *Vigna unguiculata*; plant length; vegetative growth; potassium feldspar; biochar; compost; sandy soil; organic matter; bulk density; total porosity.

1. INTRODUCTION

Cowpea (*Vigna unguiculata* ssp. *unguiculata*) has been introduced to the Egyptian agriculture as a promising field crop. It is a short duration legume crop with low water requirements, with high nutritive value and known in both southern parts of Asia and Africa for human consumption [1]. Cowpea as a summer crop will compete with other summer dominant crops in Egypt. Another usage of cowpea it could be employed as vegetable crop for its green pods which could be cooked, however cowpea cultivation as a vegetable crop haven't a role in the Egyptian crop structure and untraditional practices like intercropping could be used for such purpose [2]. Due to intensified agricultural mechanization, soils are exposed to increased levels of degradation. Soil compaction is the beginning of this degradation which further proceeds to decrease organic matter content of soils. Among the various methods applied to solve these problems, the addition of organic and non-organic matter to the soil has an important place. In Egypt, it should be used large amount of K chemical fertilizers to maximize crop yield per unit area and to compensate K-decreases in soils due to crop uptake, runoff, leaching and soil erosion [3]. Also, the high price of these fertilizers is responsible for increasing production cost and environmental pollution. The use of natural potassium fertilizer and/or bio-fertilizer is low cost resources for providing plants with K which could alternate the expensive applied K-chemical fertilizers [4,5]. The main natural sources of K come from the weathering of minerals (K-feldspar, leucite, K-mica and illite [6]. Many authors reported that K-feldspar may be valuable as a low releasing K and cheaper source of potassium [7, 8]. Compost, which originates from the decomposition of plant material or organic wastes, is considered as amid fertilizer [9].

The application of composted organic bio-wastes as organic fertilizers is still popular, and agricultural use of compost is found to have a definite advantage to the soil quality improvement. Infusion of compost into the soil increases soil nutrients and organic matter, thus positively affecting the physical, chemical and biological properties of the soil. [10]. In recent years, biochar, which is a solid product of biomass pyrolysis, is considered an evidenced

soil amendment because it is attributed to soil carbon pool and can remain intact in the soil. Biochar, which is recognized as a valuable soil ameliorant, can offset the effects of climate change because it has a definite capability of retaining too much carbon [11]. An sample number of evidence shows that it affects soil physical, chemical and biological properties by acting as soil amendment [12,13,14,15,16]. Furthermore, the effects of biochar application on other soil physical properties, such as PR, hydraulic conductivity, bulk density, and soil structure have not been fully evaluated in the field conditions [17,18,19, 20,21,22,23]. In this study, we hypothesized that the concurrent application of feldspar, compost and biochar influences the soil compaction, yet this influence can be modified by adjusting compost components and application rates. The objective of this study was to determine the effect of different compost and biochar rates on physical and chemical characteristics loam soil of sandy clay, as well as the effect of this productivity of cowpeas plants.

2. MATERIALS AND METHODS

A field experiment was conducted at Ismailia Research Station, Agricultural Research Center (A.R.C), Egypt during two successive seasons of summer 2020 and 2021 to study the effect some treatments on the physical and chemical contents of cowpea (*Vigna unguiculata* ssp. *unguiculata*) cv. Dokki 126 variety. The first treatment was control (without treatment). The soil samples were taken at depth 0-30 cm before cultivation and after harvesting to determine physical and chemical characteristics of the investigated soil according to Page et al. [24] as shown in Table (1). Some properties of the used feldspar, compost and biochar were carried out as described by Brunner and Wasmer [25] as illustrated in Table 2.

The investigation was conducted to evaluate the effect of feldspar, compost and biochar on soil properties, and productivity of plant cowpea grown in sandy loam clay soil. The initial soil was analyzed and showed in Table 1 soil analysis showed that soil was sandy texture and low fertility for available macronutrients. feldspar, compost and biochar analysis were showed in Table 2.

The applied biochar to this experiment made of different types of citrus trees it was produced using pyrolysis at a final temperature of 500 °C with a retention time of 2 h. Biochar samples were ground and sieved at <0.5 mm diameter. Biochar was applied to the soil at 4 ton/fed (feddan = 0.42 Hectare) rate while, was applied to the soil. Before planting, cowpea seeds were inoculated with the specific rhizobium (*Rhizobium radiobactersp* strain) using gum media. The inoculants was prepared by the Soil Microbiology Dept., Soil, Water and Environment Res. Inst., Agricultural Research Center (ARC), Egypt. Seeds of cowpea were sown in the first week of May in the two growing seasons. Also, before sowing the soil was prepared using agricultural gypsum ($\text{CaSO}_4 \cdot 2\text{H}_2\text{O}$) at a rate of 1190 kg/fed. Potassium sulphate (41.5 % K) at a rate of 119 kg/fed was applied in two equal portions. The first half was added at sowing and the second one was added after 30 days.

Superphosphate was applied during soil preparation. Ammonium nitrate was added at four split equal doses after 2, 4, 6 and 8 weeks from sowing. Potassium fertilizers were divided into two equal doses, the first was added at sowing after 31 days and the second was added after 55 days from sowing. At harvesting, ten plants from each plot were taken randomly, and threshed. Grain and straw were dried using an electrical oven on 75 °C until constant weight obtained. Then weighted to obtain their dry weight and transferred to seed yield. The grain and straw were ground 0.5 g powder and digested by concentrated digestion mixture of $\text{H}_2\text{SO}_4 + \text{HClO}_4$ acids according to Sommers and Nelson [26]. Nitrogen was determined by micro Keldahl, according to Phosphorus was determined by spectrophotometric ally using ammonium molybdate stannous chloride method according to Chapman and Pratt [27]. Potassium was determined by a flame photometer, according to Page *et al.* [24]. Available P and K were extracted by Ammonium-Bicarbonate- according to Soltanpour [28]. Field capacity was determined according to USDA [29]. Soil bulk density, total porosity and saturation percent were determined according to Black *et al.* [30].

At harvest, Sample of vegetative growth were taken after 85 days from sowing.

Determine the yield components as follows

1- Plant length cm. 2- Leaf number/plant. 3- No. of branches/plant. 4- Pod length cm. 5- Pod diameter cm. 6- Pod fresh weight g. 7- Pod dry weight (g) was determined by drying 100 g in an oven at 70 °C till constant weight was reached 8- Plant fresh weight g. 9- Plant dry weight (g) was determined by drying 100g in an oven at 70 °C till constant weight was reached 10- Chlorophyll a, b and carotenoids contents in the leaves were determined according Robblen, (1975)11- total Carbohydrates contents were determined in dry seeds according Herbert *et. al.*, [31] 12- Nitrogen content was determined as g/100 g dry weight according to the micro-kjeldahl method [32]. 13- Phosphorus content was determined calorimetrically as g/100 g dry weight using the hydroquinone and sodium sulphite method [32]. 14- Potassium contents were determined as g/100g dry weight using flam photometer according to Dewis and Freitas [33].

The used compost was analyzed for some physical and chemical properties and its content of some essential plant nutrients according to the methods described by Page *et al.* [24] and the obtained data are recorded in Table 2. Some physical and chemical properties of biochar in Table (3) . This experiment included 8 treatments, which were the combinations between three soil amendment with rate biochar (7.350 m³ /fed.), compost (15.750 m³ /fed.) and feldspar applications of potassium feldspar with soil inoculation of both phosphate and potassium dissolving bacteria (3.150 m³ /fed.).

- 1- Control treatment without any addition with supplying the full recommended dose of phosphorus (200 kg/ fed. of super phosphate, 15.5 % P₂O₅), The full dose of potassium (50 kg/fed. of potassium sulphate, 48 % K₂O) and nitrogen (150 kg/fed. of ammonium nitrate, 33 % N) for all experimental plots
- 2- potassium feldspar
- 3- Biochar
- 4- Compost
- 5- potassium feldspar + Biochar+ Compost
- 6- potassium feldspar + Biochar
- 7- potassium feldspar + Compost
- 8- Biochar+ Compost

Table 1. Some physical and chemical properties of the investigated surface layer of soil (0-30 cm)

Chemical properties						
pH	EC** dS/m	Soluble cations**(meq.L)				
7.59	1.31	Ca ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺	
		5.51	2.75	10.69	1.03	
SAR	ESP	Soluble anions**(meq.l)				
1.64	3.34	Co ₃ ⁻	Hco ₃ ⁻	Cl ⁻	SO ₄ ⁼	
		7.70	3.20	14.80	2.91	
SP	CaCO ₃ %	O.M%	Available (mg kg ⁻¹)			
21.23	5.95	0.41	N	P	K	
			44.2	4.99	171.0	
Physical properties						
Particle size distribution (%)						Texture class
Coarse sand	Fine	Total fined	Silt	Clay	sand	Sandy clay loam
18.69	34.06	44.89	9.00	27.31	63.69	
Hydraulic conductivity (cmhr ⁻¹)		8.40	Available water %			7.20
Field capacity % v/v		9.13	Total porosity %			47.5
Wilting point % v/v		1.90	Bulk density (Mg m ⁻³)			1.62
<i>pH⁻ : in suspension 1:2.5</i>						
<i>EC**(ds/m) , soluble cations ** and anions (meq.l) :in saturated past extract</i>						

Table 2. Some physical and chemical properties of tested compost

Chemical properties		Total nutrient (%)			
pHSus.1:10dSm ⁻¹	EC dSm ⁻¹ 1:10 Extract	Organic matter	N	P	K
2.27	7.47	38.17	1.06	0.94	1.27
Available nutrients concentration mg kg ⁻¹					
Macronutrients			Micronutrients		
N	P	K	Fe	Mn	Zn
442	237	673	283	0.98	0.82
physical properties					
Bulk density kg m ³		Moisture content %	Water holding capacity %		
500		25.72	310		

Table 3. Some physical and chemical properties of biochar

Organic carbon%	26.91	Total K %	8.11
Organic matter%	46.39	CEC (cmol / kg)	32.15
Ash (g kg ⁻¹)	3.2	Moisture (g kg ⁻¹)	60
Fixed (g kg ⁻¹)	60	Volatiles (g kg ⁻¹)	233
EC (dSm ⁻¹) (1:10 bio char water suspension) dS/m	1.4	pH	6.87
Total N %	1.55	Bulk density (g/cm ³)	0.16
Total P %	0.65		

3. RESULTS AND DISCUSSION

Effect of potassium feldspar, biochar and compost on the physical properties of soil.

Bulk density, total porosity and hydraulic conductivity

Data presented in Table 4 showed that application of feldspar + biochar + compost improved bulk density (BD) values, total porosity (TP) and hydraulic conductivity (K_{sat}) in two seasons compared with control. The best value of bulk density, total porosity and hydraulic conductivity were 1.28 (Mg m⁻³)

³), 39.3 (%) and 8.17 (cm/h) respectively, in the second season with treatment by potassium feldspar + biochar + compost was more than the other treatments. Application of biochar + compost reduced bulk density and hydraulic conductivity values by percentage 22.42 and 9.3% respectively, compared with control, while total porosity values was increased by percentage 10.3 % compared with control. These results were agreement with biochar could also reduce bulk density in the long term by interacting with soil particles and improving aggregation and porosity. The latter requires the monitoring of changes in bulk density, total porosity and hydraulic conductivity through extended periods of time [34, 35]. Compost application to agricultural lands has been recognized as a reliable way to improve the physical properties of most soils, especially soils changes in physical properties in compost-amended urban soils have included bulk density, infiltration rate, hydraulic conductivity, water content, aggregate stability and porosity. [36 ,37]. These results were in harmony with El-Shony et al. [38], who mentioned that the combined application of biochar and compost improved the porosity and bulk density of the soil compared with the sole application of biochar.

Field capacity, wilting point and available water

Data presented in Table (5) showed the application effect of the feldspar + biochar + compost on soil field capacity (FC), wilting point (WP) and the content of available water (AW) in the studied sandy clay loam soil. The values of FC, WP and AW in testing soils positively affected responded to the application of different treatments as comparatively observed with the control treatment. Whereas, the values of FC, WP and AW values increased from 11.16, 2.14 and 9.02% in the control treatment to (13.88, 6.33 and 7.55) under potassium feldspar + biochar + compost addition, 13.98, 6.44 and 7.45 under potassium feldspar + Biochar + Compost addition, first and second season respectively. The highest increase percentages of retained at field capacity, wilting point and available water were noticed in treatment with feldspar + biochar + compost and biochar + compost. Application of biochar + compost. Caused raised of water retained at either field capacity or wilting point. These results are in similar with these

obtained by [39]. The increase in water availability could be ascribed to the beneficial effect of such materials on soil aggregation. On the other hand, the lowest values of field capacity, wilting point and available water were recorded in feldspar, biochar, compost mix together when compared by control and the other treatment [40]. Also, these results may be attributed to the addition of soil organic conditioners either addition increased the specific surface area and the water holding pores area which increased water field capacity, wilting point and available water [41]. Similar finding was observed by [42] and [43]. Also this can be attributed to the inferiority in the composition and stability of soil aggregates, [44]. The increases in available water, hence, lead to a feasible route conditions and plant growth. In general, the aforementioned sequence illustrates the superiority of compost and biochar at its higher which is a logic result because of its relatively which indicates that decomposition of the compost and biochar would take longer time than the other used composites and hence its conditioning effect would still remain for longer periods.

Effect of the feldspar + biochar + compost on soil chemical properties

4. SOIL PH

Data presented in Table (6) indicated that, the values of soil pH were slightly decreased with addition the different soil amendments feldspar + biochar + compost compared to the control. But the other hand, the effects of the different soil amendments showed non-significant effect on soil pH. These results may be due to contain low plant residual which base forming cautions so cause decreasing pH. In the same trend with feldspar agreement by [45] and might be attributed to organic acids produced during the decomposition of the organic fraction of the soil organic conditioner's production of organic acids, which formed as a result of organic matter decomposition from compost [46]. Also who found that the biochar combined with feldspar was slightly reduced the pH by 0.22 than biochar alone. Albert and Kwame [47] suggested that the added Biochar as soil amended level reduce pH. Reduction in soil pH may be related to the residual organic matter after different biochemical and chemical changes.

Table 4. Effect of olive mill waste water and boichar combined with different NPK rate on soil bulk density (BD), total porosity (TP) and hydraulic conductivity (K_{sat}) in two seasons

Treatments	First season			Second season		
	BD (g/cm ³)	TP %	Ksat (cm/h)	BD (g/cm ³)	TP%	K_{sat} (cm/h)
Control	1.62g	47.5g	8.40f	1.65g	47.8f	8.38f
potassium feldspar	1.43d	44.1f	8.31e	1.41e	42.1e	8.27c
Biochar	1.46e	41.3c	8.23bc	1.42e	41.3c	8.211b
Compost	1.49f	40.7b	8.22ab	1.46f	40.7b	8.20b
potassium feldspar+ Biochar+ Compost	1.33a	40.3a	8.21a	1.28a	39.3a	8.17a
potassium feldspar + Biochar	1.37c	42.5d	8.35c	1.37c	41.6c	8.33e
potassium feldspar+ Compost	1.34b	42.6d	8.34cd	1.32b	42.1d	8.31d
Biochar+ Compost	1.34b	42.70e	8.33c	1.33b	41.40b	8.31d
LSD. at 0.05 Treatment	0.39	2.19	1.57	0.32	2.13	0.31

Table 5. Effect of potassium feldspar + biochar + compost on soil physical properties on field capacity (FC), wilting point (WP) and available water (AW) in two seasons

Treatments	First season			Second season		
	FC %	WPI %	AW %	FC %	WP I%	AW%
Control	11.16g	2.14g	9.02f	11.66g	2.24e	9.42g
potassium feldspar	13.01e	5.56c	7.45a	13.33d	5.60b	7.73d
Biochar	13.22b	5.61b	7.61c	13.62b	5.68b	7.94f
Compost	13.11c	5.58c	7.53b	13.47c	5.64b	7.83e
potassium feldspar+ Biochar+ Compost	13.88a	6.33a	7.55b	13.98a	6.44a	7.54b
potassium feldspar+ Biochar	13.09d	5.15d	7.93d	13.33d	5.32c	7.89c
potassium feldspar+ Compost	12.15c	4.5f	7.65c	12.20e	5.01d	7.19a
Biochar+ Compost	12.98f	4.79e	8.19e	13.03f	5.32c	7.71d
LSD. at 0.05 Treatment	2.72	3.93	1.57	2.32	4.23	2.34

5. EC

Data in Table (6) showed that the electric conductivity (EC) values were affected with addition of feldspar, biochar, and compost with non-significant differences between all the experimental treatments in both season. The low EC value was (1.091) (dS/m⁻¹) at second season compared with other treatments. The decrease in EC may be related to the salt leaching of salts outside the root zone [48]. These results agree with Jones et al. [49] who reported that the applying of biochar together with N fertilizer led to decrease soil salinity

(EC). These results may be due to that biochar application has reduced soil bulk density and improved soil aggregate structure, which led to increase total porosity in soil and increase in macro pores and in turn to increased water content at low suction pressures led to movement of leaching water that enhance progressive removal for Na-salts [50]. Also [51] confirmed the average of electrical conductivity of soil during both season were decreased by increasing the amount of organic fertilizers applied. This may be due to increasing water holding capacity. This led to a decrease in the rate of evaporation from the soil surface, and

thus a decrease in the concentration and accumulation of salts in the soil. "EC" were decreased by increasing the volume of applied water where as a result, the process of leaching of salts repeated with full irrigation outside the root spread area by deep percolation. The same trend in decrease the electric conductivity (EC) because that the dominant conduction mechanism in dry K-feldspar at high temperature is ionic conduction, and potassium as the major charge carriers migrates in the crystal lattice by the interstitial hopping mechanism in sandy soil [52].

6. OM

Table (6) indicated that, the values of the percentage of soil organic matter content during both season were significantly increased by addition of feldspar, biochar, and compost applied mix together. The percentage value of increase organic matter of feldspar, biochar, and compost mix in first season was 32.5 % and 37.2 %. The increase in organic matter following biochar application could be due to high carbon (C) associated with biochar [53]. While several studies showed an increase in organic matter content, total N and C/N ratio following irrigation with biochar and may have a beneficial effect on soil fertility [54, 55,56]. Improved sandy loam soils by compost application to sustain crop production. All compost types significantly increased soil total carbon [57]. They added that compost has a positive effect on overcoming the depressive effect of water stress [58]. This increase of organic matter could attributed to the activity of micro organisms for bio-fertilizer which enhanced the hydrolysis of applied compost and consequently the soil organic matter

concentration increased, [59,60]. Application of different K sources to tomato plants led to varying amounts of organic C and available NPK in the soil after harvest [61]. Organic C and available N content increased appreciably by the application of all organic sources, while mineral sources (feldspar and potassium sulphate) had no or little impact on fertility build up [61].

7. NPK

Generally, data showed Table 7 that the available N, P, and K in soil were significantly enhanced by using treatments feldspar alone or mix with other treatment biochar and compost, compared with control. The highest values of N, P and K contents in soil were 48.1, 562 and 185.0 mg/kg soil respectively, at first season and 49.1, 6. 62 and 187 mg/kg soil respectively, at second season for soil treated by feldspar + biochar + compost than other treatments. These results are in agreement with those reported by [62] the use of natural potassium fertilizer as feldspar rock is low cost resources for providing plants with K which could alternates the expensive applied K-chemical fertilizers. Also, it can able to solubilize rock – K powder, such as feldspar through plant production and excretion of soil organic acids or chelate silicon ions to bring K into solution [5,63]. [64] strongly suggested that using biofertilizers plus half dose of feldspar and chemical fertilizers have resulted in the greatest yield. They revealed that 50 % of required potassium fertilizer could be replaced by bio and feldspar fertilizers, because bio and feldspar fertilizers improved the efficiency use of recommended potassium fertilizer and reduced the cost of chemical fertilizer also reduced the environment pollution from extensive application of chemical fertilizer.

Table 6. Effect of potassium feldspar + biochar + compost on soil pH, EC and OM of two seasons in tested soil

Treatments	First season			Second season		
	pH	EC	OM	pH	EC	OM
Control	7.80e	1.29d	0.50e	7.81f	1.31e	0.53f
Potassium feldspar	7.71c	1.15b	0.56d	7.67c	1.13b	0.59e
Biochar	7.73d	1.18c	0.60c	7.70e	1.15c	0.66d
Compost	7.72c	1.16c	0.63b	7.71e	1.14b	0.683c
potassium feldspar + Biochar + Compost	7.61a	1.11a	0.66a	7.59a	1.09a	0.71a
potassium feldspar + Biochar	7.71c	1.17c	0.62b	7.69d	1.16d	0.70b
potassium feldspar+ Compost	7.63b	1.16b	0.61c	7.61b	1.15c	0.67d
Biochar+ Compost	7.64b	1.15b	0.64b	7.60b	1.14b	0.68c
LSD. at 0.05						
Treatment	0.04	0.14	0.08	0.05	0.18	0.09

Table 7. Effect of potassium feldspar + biochar + compost on NPK of two seasons in tested soil

Treatments	First season (mg Kg ⁻¹ soil)			Second season (mg Kg ⁻¹ soil)		
	N	P	K	N	P	K
Control	39.20e	4.99e	171.00d	45.10	5.09g	166.00e
Potassium feldspar	44.50d	5.19d	175.00c	47.20b	5.819e	183.00b
Biochar	45.90c	4.61b	179.00b	46.90c	6.48c	171.00d
Compost	45.10c	4.60b	183.00a	46.10c	5.57f	179.00c
potassium feldspar+ Biochar+ Compost	48.10a	5.62a	185.00a	49.10a	6.62a	187.0a
potassium feldspar+ Biochar	46.90b	5.56c	181.00b	47.19b	6.56b	183.00b
potassium feldspar+ Compost	46.10c	5.45c	184.10a	46.80c	6.55b	185.10b
Biochar+ Compost	47.10b	5.61b	182.00a	47.10b	6.12d	183.00b
LSD. at 0.05 Treatment	2.6	0.77	3.33	2.34	0.75	2.3

Table 8. Effect of potassium feldspar + biochar + compost on NPK contents in seeds of cowpea

Treatments	First season (g plant ⁻¹)			Second season (g plant ⁻¹)		
	N	P	K	N	P	K
Control	3.77f	0.30f	0.45e	3.97g	0.37g	0.55e
potassium feldspar	4.86c	0.51c	0.94d	4.98c	0.59b	1.14b
Biochar	4.89b	0.54b	0.96c	4.93d	0.54d	1.06c
Compost	4.86c	0.52c	0.95d	4.91e	0.57c	1.07c
potassium feldspar+ Biochar+ Compost	4.90a	0.58a	1.01a	5.09a	0.678a	1.21a
potassium feldspar+ Biochar	4.22d	0.36e	0.940d	4.29e	0.41e	1.04c
potassium feldspar+ Compost	4.85c	0.52c	0.991b	5.05b	0.57c	1.011d
Biochar+ Compost	4.15e	0.34e	0.951c	4.25f	0.39f	1.03c
LSD. at 0.05 Treatment	1.58	0.25	0.476	1.87	0.21	0.671

7.1 NPK Contents in Seeds of Cowpea

Data presented in Table 8 that the concentration of macronutrients i.e., N, P and K in the studied seed under the effect of different treatments. Data show that the increasing of available N, P, and K concentrations in seed as significantly by using treatments mix feldspar, biochar and compost, with two seasons compared with control. The highest values of N, P and K contents in seeds were 4.90, 0.58 and 1.01 % seed at first season and 5.09, 0.678 and 1.21 (mg plant⁻¹) seed at second season for plant treated than other treatments. These results are in agreement with those reported by [65] reveal that feldspar application had a positive effect on the studied seed chemical contents from NPK on

growth of cotton and its development as well as K %, chlorophyll A and/or B only by increasing the solubility of rock feldspar, consequently released available K to plants. These results are agreement with those obtained by [7,66,67,68]. Data revealed that application of compost to sandy soil significantly increased seed contents (%) of N, P and K that result revealed by Abou Hussien et al., [69] reported that compost application to soil increased soil content of available macro and micro nutrients as well as nutrients uptake by growing plants. Increased uptake of nutrients upon application of compost is mainly due to effect of compost on soil physical, chemical and biological properties which increase nutrients availability and finally their uptake by plants [70,71]. Also, biochar for good soil fertility and nutrient use efficiency of crops it becomes imperative that addition of a

fast releasing nutrient source to biochar be sought, These results are in agreement with those reported by Aruna et al., [72] who showed biochar (4 ton/fed) was responsible improve nutrient status content in plant. Therefore, N and P availability could be expected to increase with biochar application rather than responsible improve crop growth and nutrient status.

8. PLANT GROWTH

Results in Tables 9 & 10, revealed that there are significant effect by the using treatments mix feldspar, biochar and compost on all studied traits of plant growth in both seasons compared with control. The maximum values of plant length (cm), number of leaves, number of branches/plant was 305.00 and 278.80 cm, 84.67 and 82.00, 5.33 and 5.67 respectively, also pod length cm, pod diameter, cm Pod fresh weight g was 23.83 and 24.23, 0.82 and 0.85, 13.23 and 12.33 in the first and second seasons. In the Tables (11 & 12) the same trend, pod dry weight, plant fresh weight (g/fed), Plant dry weight (g /fed), Carbohydrates g/100 g D.W. was 422.14 (g/fed), 114.90 (g/fed) in the first and second season were achieved with mix feldspar, biochar and compost, application than other treatments at first and second seasons. These results are agreement with the

application of biochar helps in improving of soil physiochemical properties, which leads in the increase of grain yield [73] and sufficient amount of soil nitrogen availability lead to increase in plant growth and yield [74] and also the increase of 43–68 % in grain yield is due to nitrogen application [75,76]. Also, the addition of compost stimulates plant root growth and increases root system that makes a plant more drought resistant due to its ability to absorb more water from the 17 soil. The increased root system also helps the plant to increase its nutrient uptake. Leaching is reduced drastically because of increased water and nutrient retention capacities of the soil as a result of increased organic matter provided by the compost [77]. Compost improve soil fertility and plant water availability to plant. [78, 79]. [5] reported that the positive influence of the fine grains of the K mineral bearing rocks is improving the poor structure of loose sandy soil, consequently the water and nutrient capacities of this soil will be enhanced and increase their ability to cowpea plant uptake. Sugumaran [80] pointed to the feldspar bearing rocks are quite resistant to weathering and supply relatively small quantities of K during given growing season. In the meantime their cumulative release of K over several years is of some importance. This release is enhanced by the solvent action of carbonate and organic acids.

Table 9. Effect of potassium feldspar + biochar + compost on plant length, No. of leaves and No. of branches/plant in cowpea plant

Treatments	First season			Second season		
	Plant length (cm)	Number of leaves	No. of branches / plant	Plant length (cm)	Number of leaves	No. of branches / plant
Control	144.30 f	42.67 e	3.67 bc	127.00 f	49.33 g	3.33 b
potassium feldspar	199.90 e	52.33 de	3.33 c	185.78 e	61.67 e	3.67 b
Biochar	224.70 d	56.33 cd	4.33 abc	214.43 d	56.00 f	4.00 b
Compost	191.87 e	71.33 b	3.33 c	180.18 e	52.00 g	3.33 b
potassium feldspar+ Biochar+ Compost	305.00 a	84.67 a	5.00 ab	278.80 a	82.00 a	5.67 a
potassium feldspar+ Biochar	265.93 b	72.00 b	5.33 a	258.43 b	74.33 b	4.33 b
potassium feldspar+ Compost	259.77 b	66.00 bc	4.33 abc	249.43 b	70.00 c	4.00 b
Biochar+ Compost	250.23 c	59.00bcd	4.00 abc	237.17 c	65.33 d	3.67 b
LSD. at 0.05	9.52	12.28	1.223	9.63	3.43	1.059

Table 10. Effect of potassium feldspar + biochar + compost on pod length, Pod diameter and Pod fresh weight in cowpea plant

Treatments	First season			Second season		
	Pod length cm	Pod diameter cm	Pod fresh weight g	Pod length cm	Pod diameter cm	Pod fresh weight g
Control	18.17 f	0.58 d	8.35 f	17.70 f	0.60 g	7.85 g
potassium feldspar	20.67 e	0.70 c	9.84 e	20.10 d	0.69 e	8.85 e
Biochar	21.40 d	0.72 c	10.69 d	20.53 d	0.72 d	9.19 d
Compost	20.23 e	0.57 d	8.89 f	19.03 e	0.65 f	8.45 f
potassium feldspar+ Biochar+ Compost	23.83 a	0.82 a	13.23 a	24.23 a	0.85 a	12.33 a
potassium feldspar+ Biochar	22.93 b	0.80 a	12.53 b	23.30 b	0.82 b	12.08 a
potassium feldspar+ Compost	22.63 bc	0.77 ab	11.85 c	22.87 b	0.78 c	11.53 b
Biochar+ Compost	22.13 c	0.74 bc	11.35 c	21.77 c	0.73 d	10.32 c
LSD. at 0.05	0.59	0.046	0.58	0.92	0.028	0.32

Table 11. Effect of potassium feldspar + biochar + compost on pod dry weight, plant fresh weight and plant dry weight in cowpea plant

Treatments	First season			Second season		
	Plant fresh weight g	Pod dry weight g	Plant dry weight g	Plant fresh weight g	Pod dry weight g	Plant dry weight g
Control	358.74 h	2.60 f	92.50 f	353.28 h	2.64 e	98.53 g
potassium feldspar	372.75 f	3.36 d	98.07 e	369.03 f	3.20 d	106.48 e
Biochar	378.17 e	3.59 d	102.59 d	377.91 e	3.14 d	109.22 d
Compost	366.05 g	3.07 e	95.29 ef	360.32 g	3.04 d	102.49 f
potassium feldspar+ Biochar+ Compost	421.90 a	5.07 a	118.78 a	441.25 a	4.59 a	121.33 a
potassium feldspar + Biochar	397.46 b	4.52 b	111.76 b	412.54 b	4.22 b	116.83 b
potassium feldspar + Compost	390.28 c	4.14 c	106.32 c	401.91 c	4.14 b	112.22 c
Biochar+ Compost	Plant fresh weight g	3.96 c	104.08 cd	392.34 d	3.78 c	108.68 d
LSD. at 0.05	358.74 h	0.27	2.93	6.53	0.15	1.74

Table 12. Effect of potassium feldspar + biochar + compost on carbohydrates and chlorophyll contents of cowpea plant in two seasons

Treatments	First season		Second season	
	Carbohydrates g/100 g D.W.	Chlorophyll A mg/100g F.W.	Carbohydrates g/100 g D.W.	Chlorophyll A mg/100g F.W.
Control	59.81 e	16.27 h	59.99 g	13.10 e
potassium feldspar	60.39 d	18.60 f	61.38 e	14.75 d
Biochar	60.89 c	21.62 e	61.92 de	16.15 c
Compost	58.06 f	17.73 g	60.74 f	13.86 de
potassium feldspar+ Biochar+ Compost	63.23 a	26.07 a	64.80 a	20.86 a
potassium feldspar+ Biochar	62.89 a	24.68 b	63.59 b	19.97 a
potassium feldspar+ Compost	61.80 b	23.73 c	62.69 c	18.55 b
Biochar+ Compost	61.27 c	22.68 d	62.01 d	16.88 c
LSD. at 0.05	0.447	0.77	0.579	1.00

Table 13. Effect of potassium feldspar + biochar + compost on chlorophyll b and carotenoids contents in leaves of cowpea plant in two seasons

Treatments	First season		Second season	
	Chlorophyll B mg/100g F.W.	Carotenoids mg/100g F.W.	Chlorophyll B mg/100g F.W.	Carotenoids mg/100g F.W.
Control	9.16 h	3.59 g	5.57 f	3.59 g
potassium feldspar	11.29 f	5.04 ef	6.52 de	5.04 ef
Biochar	12.09 e	5.45 de	6.95 cd	5.45 de
Compost	10.43 g	4.50 f	7.42 c	4.50 f
potassium feldspar+ Biochar+ Compost	16.13 a	8.45 a	9.89 a	8.45 a
potassium feldspar+ Biochar	14.83 b	7.28 b	9.38 a	7.28 b
potassium feldspar+ Compost	14.05 c	6.55 c	8.44 b	6.55 c
Biochar+ Compost	12.903 d	5.96 cd	7.42 c	5.96 cd
LSD. at 0.05	0.67	0.70	0.52	0.35

The correlation analysis revealed that plant length, number of leaves, effective nodule, and chlorophyll content had positive correlation to growth and yield production in cowpea. This is an indication that the taller the plant the more likely it is to have more leaves and greater the opportunity for more chlorophyll content in leaves and the greater the chance for photosynthesis to occur leading to vegetative growth. This agrees with report by [81] that tall plants are likely to have more green areas for more photosynthetic activities that facilitate increase in growth and grain yield of crops.

9. CONCLUSION

Based on the above discussion, it can be concluded that we can achieve increasing soil water and nutrient retention of low fertility soil by applying organic amendments like biochar, compost and potassium feldspar [82]. Moreover, biochar, compost and potassium feldspar as slow-release fertilizers improved soil's physical and chemical properties and enhanced plant growth and yield parameters of cowpea crop. Concerning the application method, the mixed application of two organic amendments and one inorganic, especially at mix treatment together superiority in soil properties and yield productivity as compared with sole application of biochar alone or compost alone or potassium feldspar. The beneficial effect of biochar compost and potassium feldspar mix could be due to the organic acids released from the mixture, which positively reflected the availability of nutrients and their concentration in different plant parts of cowpea plants.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Elsayed E.A. Elsobky and Hend H.M. Hassan2021 Optimizing Cowpea Productivity by Sowing Date and Plant Density to Mitigate Climatic Changes .Egyptian Journal of Agronomy Volume 43, Issue 3,Page 317-331.
2. Abd El Lateef EM, Eata AEM, Asal M Wali, MS Abd El-Salam. Evaluation of Mungbean (*Vigna radiata* L. Wilczek) as Green Pod and Seed Crop under Different Cropping Systems in Egypt. Asian Journal of Crop Science. 2020;12:115-123.
3. Shams AS, WA Fekry Efficiency of applied K-feldspar with potassium sulphate and silicate dissolving bacteria on sweet potato plants. Zag. J. Agric. Res. 2014;41(3):467-477.
4. Manning, D. A. C. (2010): Mineral sources of potassium for plant nutrition. A Review Agron. Sustain Dev. pp: 281-294.
5. Labib BF K Ghabour, IS Rahim, MM. Wahba Effect of potassium bearing rock on the growth and quality of potato crop (*Solanum tuberosum*). J. Agric. Biotech. Sustainable Dev. 2012;4(1):7 -15.
6. Hellal FA, Abd El-Hady M, AAM Ragab Influence of organic amendments on nutrient availability and uptake by faba bean plants fertilized by rock phosphate and feldspar. Amer-Eurasian J. Agric. & Environ. Sci. 2009;6 (3):271-279.
7. Shafeek MR, OA H EL- Zeiny and MA. Ahmed Effect of phosphate and potassium fertilizer on growth, yield seed composition of pea plant in new reclaimed soil. Ass. J. plant. Sci. 2005;4(6):608-612.
8. Abou-el-Seoud B, A Abdel- Mageed. Impact of rock materials and bio fertilizations on P and K availability for maize (*Zea maize*) under calcareous soil conditions. Saudi J. Biol. Sci. 2012;19 (1):55–63.
9. Aggelides SM, PA Londra Effects of compost produced from town wastes and sewage sludge on the physical properties of a loamy and a clay soil. Bio resource Technology. 2000;71:253–59. DOI: 10.1016/S0960-8524 (99) 00074 -7
10. Hargreaves JC, MS Adl and PR. Warman. A review of the use of composted municipal solid waste in agriculture. Agric Ecosyst Environ. 2008;123:1–14.
11. Peake L, Freddo A, BJ Reid Sustaining soils and mitigating climate change using biochar, Chapter 7, in A. Des Las Heras TR. Macagno (Eds.). Sustainability science and technology, an introduction. Taylor&Francis Group, CRC Press; 2014. Available:https://doi.org/10.1201/b16701-8
12. Głąb T, A Żabiński, U Sadowska, K Gondek, M Kopeć, M Mierzwa-Hersztek and S. Tabor Effects of co-composted maize, sewage sludge, and biochar mixtures on hydrological and physical qualities of sandy soil. Geoderma. 2018;315(3):27–35.

13. Lehmann J.; M. C. Rillig; J. Thies; C. A. Masiello; W. C. Hockaday and D. Crowley (2011) Biochar affects on soil biota—a review. *Soil Biol Biochem* 43:1812–1836
14. Li Y, SJ, Hu K, Chen Y, Müller W, Li Z, Fu Lin and H. Wang. Effects of biochar application in forest ecosystems on soil properties and greenhouse gas emissions: A review. *Journal of Soils and Sediments* 2018;18:546–63.
15. Mukherjee A, R Lal Biochar impacts on soil physical properties and greenhouse gas emissions. *Agronomy*. 2013;3:313. DOI: 10.3390/agronomy3020313.
16. Wang T, CE, C Stewart, YS. Wang and J. Zheng Effects of biochar addition on evaporation in the five typical Loess Plateau soils. *Catena*. 2018;162:29–39. DOI:10.1016/j.catena.2017.11.013.
17. Asai H, BK, H Samson, M Stephan, K Songyikhangsuthor, K Homma, Y Kiyono, Y Inoue, T. Shiraiwa, T Horie Biochar amendment techniques for upland rice production in Northern Laos: 1. Soil physical properties, leaf SPAD and grain yield. *Field Crops Research*. 2009;111:81–84. DOI:10.1016/j.fcr.2008.10.008.
18. Busscher, WJ, JM Novak, DE Evans, D. W. Watts, MAS. Niandou; and M. Ahmedna. Influence of pecan biochar on physical properties of a norfolk loamy sand. *Soil Science*. 2010;175:10–14. DOI:10.1097/SS.0b013e3181cb7f46.
19. Chan KY, L Van, IZ. Meszaros A. Downie and S Joseph. Agronomic values of greenwaste biochar as a soil amendment. *Soil Research* 2007;45:629–34. DOI:10.1071/SR07109.
20. Glaser B, J Lehmann and W. Zech Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal, a review. *Biology and Fertility of Soils*. 2002;35:219–30. DOI:10.1007/s00374-002-0466-4.
21. Laird, D. A.; P. Fleming; D. D. Davis; R. Horton; B. Wang and D. L. Karlen. Impact of biochar amendments on the quality of a typical Midwestern agricultural soil. *Geoderma*. 2010;158:443–49. DOI:10.1016/j.geoderma.2010.05.013.
22. Masulili A, WH. Utomo and M. Syechfani (2010): Rice husk biochar for rice based cropping system in acid soil 1. The characteristics of rice husk biochar and its influence on the properties of acid sulfate soils and rice growth in West Kalimantan, Indonesia. *The Journal of Agricultural Science* 2:39.
23. Peng X, LL Ye, CH Wang, H Zhou, B Sun. Temperature and duration dependent rice straw derived biochar: Characteristics and its effects on soil properties of an Ultisol in southern China. *Soil & Tillage Research*. 2011;112:159–66. DOI:10.1016/j.still.2011.01.002.
24. Page AL, RH Miller, DR Keeney. *Methods of Chemical Analysis. Part2: Chemical and properties (Second Edition)*. American Society of Agronomy, Inc. and Sci. Soc. Of America, Inc. Publishers. , Madison, Wisconsin USA;1982.
25. Brunner PH, HR Wasmer. *Methods of analysis of sewage sludge solid wastes and compost*. W.H.O. International Reference Center for Wastes Disposal (H-8600), Dülendorf Switzerland;1978.
26. Sommers LE, DW Nelson. Determination of total phosphorus in soil: A rapid perchloric acid procedure. *Soil Science Society of America Journal*. 1972; 36:902-904.
27. Chapman HD, PF Pratt. *Methods of Analysis for Soils, Plants and Waters*. Univ. California Div. Agric. Sci. Priced Publication, Oakland;1978.
28. Soltanpour PN. Use of Ammonium Bicarbonate DTPA Soil Test to Evaluate Elemental Availability and Toxicity. *Communications in Soil Science and Plant Analysis*. 1985;16:323-338.
29. USDA. *Soil survey manual*. U.S. Dept. of Agric. (USDA), Hand Book No.18. Washington Dc;1962.
30. Black GR, DD Evans, LE, Ensminge JL, White, and FE Chark. *Methods of soil analysis, Agron. Series , AM. Agron., Madison, Wisconsin, USA;1965*.
31. Herbert D, P Phlipps, R Strange. Determination of total carbohydrates. *Methods in Microbial*. 1971;58: 209-344.
32. AOAC. *Association Official methods of analysis of the Association of Official Analytical Chemists*. 2 vols. 15th ed. Washington, DC;1990.
33. Dewis J, F Freitas. *Physical and chemical methods of soil and water analysis*. Food and Agric. Organ United Nations. *Soils Bulletin*, No. 1970;10.
34. Lucia Toková, Dušan Igaz, Ján Horák and Elena Aydin. Effect of Biochar Application and Re-Application on Soil Bulk Density, Porosity, Saturated Hydraulic

- Conductivity, Water Content and Soil Water Availability in a Silty Loam HaplicLuvisol. *Agronomy*. 2020;10(7): 1005.
35. Vitková J, Šurda P. Soil moisture changes after biochar application in *Acta Hydrol.Slovaca* 2019;20:74–79.
 36. Somerville PD, PB May, SJ Livesley. Effects of deep tillage and municipal green waste compost amendments on soil properties and tree growth in compacted urban soils. *J. Environ. Manag.* 2018 ;227:365–374.
 37. Usowicz B, J Lipiec. Spatial variability of saturated hydraulic conductivity and its links with other soil properties at the regional scale. *Sci. Rep.* 2021;11:8293.
 38. El-Shony, M., Farid, I.M., Alkamar, F., Abbas, M.H., Abbas, H. 2019. Ameliorating a sandy soil using biochar and compost amendments and their implications as slow-release fertilizers on plant growth. *Egyptian Journal of Soil Science*, 59(4), 305-322.
 39. Jennifer Cooper; Isabel Greenberg; Bernard Ludwig; Laura Hippich; Daniel Fischer Bruno and Glaser Michael (2020): Effect of biochar and compost on soil properties and organic matter in aggregate size fractions under field conditions. *Agriculture, Ecosystems & Environment* Volume 295, 15 June, 106882
 40. N.Li¹M.-Y.You*¹B.Zhang[†]X.-Z.Han*¹S.K.Panakoulia[‡]Y.-R.Yuan[§]K.Liu*[¶]Y.-F.Qiao*^W.-X.Zou*^N.P.Nikolaidis[¶]S.A.Banwart[‡] 2017 Modeling Soil Aggregation at the Early Pedogenesis Stage From the Parent Material of a Mollisol Under Different Agricultural Practices *Advances in Agronomy* Volume 142, , Pages 181-214
 41. Christina N. Kranz, Richard A. Mc Laughlin, Amy Johnson, Grady Miller and Joshua L. Heitman (2020): The effects of compost incorporation on soil physical properties in urban soils . A concise review *J. Environ. Manag.*, 261.
 42. Schmid CJ, Murphy JA, S Murphy Effect of tillage and compost amendment on turfgrass establishment on a compacted sandy loam. *J. Soil Water Conserv.* 2017;72:55-64.
 43. Sax MS, Bassuk N, H van Es, D Rakow Long-term remediation of compacted urban soils by physical fracturing and incorporation of compost. *Urban For. Urban Green.* 2017;24:149-156.
 44. Bashir A, M Rizwan, MZ urRehman, M Zubair, M Riaz, MF Qayyum, HF Alharby, AA. Bamagoos, S Ali Application of co-composted farm manure and biochar increased the wheat growth and decreased cadmium accumulation in plants under different water regimes. *Chemosphere.* 2020; 246:125809.
 45. Davide C. and C. Thomas (2019): Fertilizing Properties of Potassium Feldspar Altered Hydrothermally *Communications in Soil Science and Plant Analysis* 50 (4): 1-10.
 46. Jeffery S.; D. Abalos and M. Prodana (2017): "Biochar boosts tropical but not temperate crop yields," *Environmental Research Letters*, vol. 12, no. 5, p. 053001.
 47. Albert K, A Kwame Biochar and or compost applications improve soil properties, growth and yield of maize grown in Acidic rainforest and coastal savannah soils in Ghana, *Research Article. Inter. J. of Agron.* 2018;6-8.
 48. Lehrsch, G. A.; R. E. Sojka and P. M. Jolley (1993): Freezing effects on aggregate stability of soil amended with lime and gypsum. *Soil surface sealing and crusting. Catena Suppl.* 24: 115 -127.
 49. Jones, D.; J. Rousk; G. Edwards-Jones; T. DeLuca and D. Murphy (2011): Biochar-mediated changes in soil quality and plant growth in a three-year field trial. *Soil Biol. Biochem.*, 12: 1- 10.
 50. Lei, O. and R. Zhang (2013): Effects of biochars derived from different feedstocks and pyrolysis temperatures on soil physical and hydraulic properties. *Journal of Soils and Sediments*, 13(9):1561-1572.
 51. Ghanema H. G.; Y. A. B. El-Gabry; E. M. C. Okasha and Shimaa K. Ganzour (2021): Improving Some Irrigation Efficiencies, Soil Fertility, Yield and Quality of Wheat under Deficit Irrigation by Integrated N-Fertilization *Egyptian Journal of Chemistry*. Vol. 64, No. 4 pp. 2201 – 2212
 52. Hu Y.; D. Xiang; S. D. Veresoglou; F. Chen; and Y. Z. Hao (2014): Soil organic carbon and soil structure are driving microbial abundance and community composition across the arid and semi-arid grasslands in northern China. *Soil Biol. Biochem.* 77 51–57. 10.1016/j. soil biol. 2014. 06.014 [CrossRef] [Google Scholar]

53. Abdulaziz GA. Biochar as a potential soil additive for improving soil physical properties-a review. *Arabian J. of Geosciences*. 2018;11:766.
54. Mekki A. A.; Dhouib and S. Sayadi (2006): Changes in microbial and soil properties following amendment with treated and untreated olive mill wastewater. *Microbiol. Res.*, 161: 93–101.
55. Brunetti G. N.; Senesi and C. Plaza (2007): Effects of amendment with treated and untreated olive oil mill wastewaters on soil properties, soil humic substances and wheat yield. *Geoderma*, 138: 144-152.
56. Mekki A. A.; Dhouib and S. Sayadi (2013): Effects of olive mill wastewater application on soil properties and plants growth. *International Journal of Recycling of Organic, Waste in Agriculture*, 2(1): 1–7.
57. Mahmoud A. A.; M. M. Rady; E. E. Belal; R. Al-Qthanin; H. M. Al-Yasi and E. F. Ali (2021): Revitalizing Fertility of Nutrient-Deficient Virgin Sandy Soil Using Leguminous Biocompost Boosts *Phaseolus vulgaris* Performance Plants, 10, 1637. <https://doi.org/10.3390/plants10081637>
58. Kusparwanti, T. R.; Eliyatningsih and R. Wardana (2020): Application Legume Compost with Bio-Activator *Trichoderma* sp as Inorganic Fertilizer Substitution in Sweet Corn (*Zea mays L. Saccharata*) Cultivation. *IOP Conf. Series. Earth Environ. Sci.* 2020, 411, 012063. [CrossRef]
59. Alshankiti A, S Gill Integrated Plant Nutrient Management for Sandy Soil Using Chemical Fertilizers, Compost, Biochar and Biofertilizers- Case Study in UAE. *J. Arid Land Stud.* 2016;26:101–106.
60. Chapman, H. D. (1965): Cation-Exchange capacity. In *Methods of Soil Analysis: Part 2 Chemical and Microbiological Properties*; Norman, A.G., Ed.; Madison, WI, USA; Volume 9, pp. 891–901.
61. Badr MA. Efficiency of K-feldspar combined with organic materials and silicate dissolving bacteria on tomato yield. *J. Appl. Sci. Res.* 2006;2:1191- 1198
62. Shimaa K, HH, AA M Ragab. Effect of potassium fertilizer, feldspar rock and potassium releasing bacterium (*Bacillus circulans*) on sweet potato plant under sandy soil conditions. *Scientific Journal of Agricultural Sciences*. 2020;2 (2):56-63,
63. Bennett PC, Choi WJ, JR Rogers Microbial destruction of feldspars. *Mineral Mag.* 1998; 62(A):129-150.
64. Habibi, A.; G. Heidari; Y. Sohrabi; H. Badakhshan and K. Mohammadi (2011): Influence of bio, organic and chemical fertilizers on medicinal pumpkin traits. *Journal of Medicinal Plants Research*. 5(23): 5590-5597.
65. Sarhan MGR. 1, SSh Abd El-Gayed. Possibility of using Feldspar as Alternative Potassium for Cotton Fertilization Combined with Silicate Dissolving Bacteria, Humic Acids and Farmyard Manure and its Effect on Soil Properties *J. Soil Sci. and Agric. Eng., Mansoura Univ.* 2016; V8 (12):761 – 767.
66. Abdel-Hak RS, S El-Shazly A, EA El-Gazzar, MS El-Shamma Response of valencia orange trees to rock-feldspar applications on in reclaimed soils. *J. Appl. Sci. Res.* 2012;8(7):3160-3165
67. Ismail, S. A.; A. M. Abd El-Hafeez; O. A. Galal and H. A. Awadalla (2014): Impact of some alternative fertilizers such as anhydrous ammonia, humic acid, rock phosphate and feldspar on growth, yield and its components and nutrient uptake of wheat as well as nutrient availability. *Fayoum J. Agric. Res. & Dev.*, 28(1): 89-107.
68. Merwad, A. M. A. (2016): Efficiency of K-sulphate and K-feldspar combined with silicate dissolving bacteria on yield and nutrient uptake by maize plants. *Egypt. J. Soil Sci.* 56(2): 249- 259
69. Abou Hussien EA, AM Elbaalawy, MM. Hamad. Chemical properties of compost in relation to calcareous soil properties and its productivity of wheat. *Egypt. J. Soil. Sci.* 2019;59:85-97.
70. Wardah L ,S, Prijono, Z Kusuma Soil health improvement by addition the manure and compost for peanut plant (*Arachis hypogaea l*) in Leranwetan, Palang, Tuban. *IOSR J. Agri. Vet. Sci.* 2014;7:64-71.
71. Emam, A. A. E. (2018): Effect of Some Organic and Inorganic Fertilizers on Plant Irrigated by Saline Water. Ph.D. Thesis, Fac. of Agric., Menoufia University, Egypt.
72. Aruna OA, TM Agbede, WS Ejue, CM Aboyeji, O Dunsin, CO. Aremu, AO. Owolabi, BO Ajiboye, OF Okunlola, OO Adesola. Biochar, poultry manure and NPK

- fertilizer: sole and combine application effects on soil properties and ginger (*Zingiber officinale* Roscoe) performance in a tropical Alfisol Source: Open Agriculture. 2020;5(1):30-39. ISSN:2391-9531
73. Muhammad F.; L. Romdhane; Abdul Rehman; A. K. M. Al-Alawi; W. M. Al-Busaidi; S. A. Asad and D. Lee (2021): Integration of Seed Priming and Biochar Application Improves Drought Tolerance in Cowpea Regulation J. of Plant Growth volume 40, pages1972–1980
74. Farooq M.; L. Romdhane; M. K. Al Sulti; A. Rehman; W. M. Al-Busaidi and D. J. Lee (2019) Morphological, physiological and biochemical aspects of osmopriming-induced drought tolerance in lentil. J Agron Crop Sc 206:176–186
75. Farooq M.; A. Ullah; M. Usman and K. H. M. Siddique (2020 a) Application of zinc and biochar help to mitigate cadmium stress in breadwheat raised from seeds with high intrinsic zinc. Chemosphere 260:127652
76. Farooq M.; A. Rehman; A. K. M. Al-Alawi; W. M. Al-Busaidi and D. J. Lee (2020 b) Integrated use of seed priming and biochar improves salt tolerance in cowpea. Sci Horti 272:109507
77. Graves, R. E. and G. M. Hattemer (2000): Chapter 2 Composting. Part 637 Environmental Engineering National Engineering Handbook. United States Department of Agriculture, Natural Resources Conservation Service. (210-VI-NEH). Available:<http://ftp.wcc.nrcs.usda.gov/wntsc/AWM/neh637c2.pdf> (Accessed on 24/6/13)
78. Curtis, M. J. and V. P. Claassen (2005): Compost incorporation increases plant available water in a drastically disturbed serpentine soil. Soil Sci. 170:939–953.
79. Farrell, M. and D. I. Jones (2009): Critical evaluation of municipal solid waste composting and potential compost markets. Bioresource Technology 100(19):4301 – 4310. Fodder crops production. Agridigest - An in house, Zari Taraqati Bank Ltd, Pakistan Journal of Agriculture. 23: 24–30.
80. Sugumaran PB. Solubilization of potassium containing minerals by bacteria and their effect on plant growth. World J Agric Sci. 2007;3:350–355.
81. Brhgh, P.D. Compost and nitrogen fertilizer on maize (*Zea mays* L) growth and yield and residual effects on cowpea (*Vigna unguiculata* L.) Walp). In a submitted to the University of Ghana, legon in partial fulfillment of the requirement for the award of master of philosophy in crop Science Degree;2017.
82. Robblen G. Quantitative analysis of chloroplast pigments. Untersuchungen an etrahlenin duzierten Blatterbumtanten von Arbidopsis Thaliana (L) verer burg Leher.1975; 8:189.

© 2022 Doaa and Ashmawi; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/85385>