



Frequency of Hypophosphatemia in Children Presenting with Sepsis in a Tertiary Care Hospital: A Cross-sectional Study

Kashif Habib Qadri ^a, Muhammad Saad Raza ^{b#}, Yousuf Yahya ^{c†},
Suneet Kumar ^{d‡}, Rumana Sangi ^{e*}, Aliya Kemal Ahsan ^{e*}
and Muhammad Nadeem Chohan ^{†*}

^a Civil Hospital Thatta, Pakistan.

^b United Medical and Dental College, Karachi, Pakistan.

^c Dow University of Health Sciences, Karachi, Pakistan.

^d Lady Dufferin Hospital, Karachi, Pakistan.

^e Post Fellow Pediatric Cardiology, NICVD, Karachi, Pakistan.

[†] Bilawal Medical College, Jamshoro, Pakistan.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2022/v34i39A36238

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/87473>

Original Research Article

Received 11 March 2022

Accepted 21 May 2022

Published 27 May 2022

ABSTRACT

Aim: To determine the frequency of hypophosphatemia in the children presenting with sepsis in a tertiary care hospital.

Study Design: A cross-sectional study.

Place and Duration: Department of paediatric, civil hospital, Karachi, Pakistan from october 2020 to april 2021.

^o Medical officer

[#] Senior Registrar Pediatrics.

[†] Assistant Professor Pediatrics.

[‡] Senior Medical Officer NICU,

^{*} FCPS Pediatric Cardiology

[†] Associate Professor Pediatrics.

*Corresponding author: E-mail: Nadeem.chohan@lumhs.edu.pk, nadeemchohanpink76@yahoo.com;

Methodology: A total of 190 children with clinical sepsis were included in the study. Serum phosphate level, CBC, CRP, and blood culture of all children were sent within 24 hours of admission. Serum phosphate level less than 2.5mg/dL was labeled hypophosphatemia. It was categorized as mild, moderate and severe at level of 2-2.5 mg/dL, 1-2 mg/dL, and < 1 mg/dL respectively.

Results: Mean \pm SD age of study participants was 2.69 \pm 3.19 years. Out of 190 patients 106 (55.8%) were male while 84 (44.2%) were female. Hypophosphatemia was found in 36 (18.9%) patients among them. Mild and moderate hypophosphatemia was noted in 18 (50%) of the children from each category of patients.

Conclusion: This study concluded that hypophosphatemia was documented in considerable number of children who presented with clinical sepsis. Serum phosphate levels should be checked in children who present with sepsis. To avoid the issues of a double burden of disease in these patients, strategies to prevent these diseases should be supported.

Keywords: Hypophosphatemia; illness; mortality; outcome; prognosis; sepsis.

1. INTRODUCTION

A dysregulated host response to the infection causes the sepsis, which is a life-threatening organ failure. Sepsis represents a significant socioeconomic burden worldwide [1]. Severe sepsis is one of the primary cause of the admission to the intensive care units (ICUS) [2-3]. The incidence and rate of hospitalization for the severe sepsis is increasing every year. In studies published in the previous decade, the incidence of sepsis in all patients admitted to ICUs ranges from 9 to 37 percent, and severe sepsis remains a prominent cause of the death. Severe sepsis mortality rates in patients admitted to intensive care units range from 30 to 50% [4-5].

Blood Culture remains the gold standard but has limitations due to final report after seven days. Hematological and inflammatory markers such as ESR, CRP, raised total leukocyte count, ANC, band cells, toxic granulations are better indicator for the sepsis but have certain limitations [6-8]. Phosphate is a component of adenosine triphosphate, 2, 3-diphosphoglycerate and intracellular chemical messengers among other intermediate molecules involved in critical physiological processes [9]. Hypophosphatemia (HP) has been linked to sepsis and has been suggested as a useful diagnostic tool [10]. During the first 24 hours of admission, patients with sepsis had a high rate of HP [10]. Phosphate (P) values of less than 2 mg/dl were particularly indicative of gram-negative sepsis. The exact process by which HP develops in sepsis is uncertain. The elimination of the illness was linked to an increase in P levels [11-13].

In the majority of the cases, hypophosphatemia symptoms are nonspecific and include fatigue

and irritability [14]. During childhood, the normal level of phosphorus in the blood changes. Normal blood phosphorus levels in children aged one to three years are (3.8 mg/dl to 6.5 mg/dl), four to eleven years (3.7 mg/dl to 5.6 mg/dl) and twelve to fifteen years (3.7 mg/dl to 5.4 mg/dl) [15].

In the postoperative period of cardiac surgery, severe hypophosphatemia can lead to leukocyte dysfunction, and neuromuscular disturbances [16-18]. Childhood sepsis played a role, with 37.37 percent having sepsis with a negative blood culture (NBC) and 30.30 percent having sepsis with a positive blood culture (PBC) (PBC). In both the PBC and NBC groups, hypophosphatemia was found in 80% of the septic patients [11].

In a study of Shazly A.N. El et al. they investigated that there are 34% sepsis cases in which 58.8% had hypophosphatemia while other 41.1% had normophosphatemia. Although a higher percentage of septic patients were hypophosphatemic, this finding did not reach statistical significance. [10]. Another study indicated that out of 8% of sepsis cases, 50% had HP and the other 50% had normophosphatemia, and that there was no significant connection between sepsis and previously recognized variables linked with hypophosphatemia [14]. The inefficiency of the respiratory muscle contraction due to hypophosphatemia makes weaning patients from artificial ventilation difficult [14].

Patients with phosphate derangements primarily focus on the association of hypophosphatemia [19,20] and the critically ill population but in our area, there is scarcity of the studies related to the sepsis and related markers hence we aimed to

conduct the study to find correlation between hypophosphatemia and severity of the sepsis.

2. METHODOLOGY

This cross-sectional study was conducted at department of paediatric, civil hospital Karachi by non-probability consecutive sampling technique from october 2020 to april 2021. The sample size was calculated by taking prevalence of hypophosphatemia in septic children $P=(58.8\%)$.⁶ Using margin of error(d)= 7% the total calculated sample size was 190 patients with the help of WHO software for sample size calculation taking 95% confidence level. Children of either gender presenting with clinical sepsis age 1 month to 15 years were included in the study. Children having chronic renal failure, congenital renal disease and severe malnutrition were excluded from the study. Hypophosphatemia was defined as the serum phosphate level less than 2.5mg/dL. It was categorized as mild: 2-2.5 mg/dL, moderate: 1-2 mg/dL and severe : < 1 mg/dL. Patients were labeled as clinical sepsis if two or more of the following criteria were fulfilled: Axillary Temperature $>38^{\circ}\text{C}$ or $<36^{\circ}\text{C}$, tachycardia or fast breathing, White blood cell count $>12,000/\text{mm}^2$, or 10% immature (band) forms, on complete blood count report.

Complete history and detailed examination of the septic children were taken at the time of admission. Serum phosphate level, CBC, CRP, and blood culture of all included children were sent within 24 hours of admission. The sepsis was diagnosed as per operational definitions. The final outcome i.e. hypophosphatemia was labeled when the serum phosphate level was less than 2.5mg/dL. It was categorized as mild, moderate and severe showing range of 2- 2.5 mg/dL, 1-2 mg/dL, < 1 mg/dL) respectively. Biasness and confounder were controlled by strictly following the inclusion criteria. All patients were given supportive and definitive management as per hospital protocols. All data was kept strictly confidential. The data was compiled and analyzed using SPSS version 21. For qualitative factors such as gender, culture, hypophosphatemia and levels of hypophosphatemia, frequencies and percentages were calculated.

Age, serum phosphate, axillary temperature, heart rate, respiration rate, white blood cell count, and CRP were all provided as the mean SD quantitative variables. Stratification was used

to control the effect modifiers such as age, gender, axillary temperature, heart rate, respiration rate, white blood cell count, and CRP. The Chi-square test was used to determine whether category variables were related. The significance level was set at $P 0.05$.

3. RESULTS

In this study, 190 patients were included to assess the hypophosphatemia in clinical sepsis children. Mean \pm SD of age was 2.69 ± 3.19 . Mean \pm SD of serum phosphate was 3.79 ± 1.30 , axillary temperature was 41.00 ± 23.27 , heart rate was 133.30 ± 21.88 with respiratory rate was 46.32 ± 14.48 . Mean \pm SD of CBC was 14.13 ± 8.65 and CRP was 56.71 ± 86.89 . Positive blood culture was found in 38 (20%) patients while negative culture was found in 72 (80%) patients. Hypophosphatemia was found in 36 (18.9%) patients. Mild and moderate hypophosphatemia was noted in 18 (50%) patients of each category. Stratification of age group, gender, axillary temperature, heart rate, respiratory rate, white blood cell (CBC) and CRP were done with respect to hypophosphatemia in order to find significant difference between the groups

Table 1. Descriptive statistics (mean) of the study participants n=190

Age (Years)	2.69 \pm 0.23
Serum phosphate level (mg/dl)	3.79 \pm 0.94
Axillary temperature (C)	41.00 \pm 1.68
Hear rate (bpm)	133 \pm 1.58
Respiratory rate (brpm)	46.32 \pm 1.05
Complete blood count	14.13 \pm 0.62
C reactive protein	56.71 \pm 6.30

Table 2. Characteristics of study participants n = 190

Characteristics	Number (%)
Gender	
Male	106 (55.8)
Female	84 (44.2)
Blood C/S	
Positive	38 (20)
Negative	72 (80)
Hypophosphatemia	
Yes	36 (18.9)
No	154 (81.1)
Severity of hypophosphatemia	
Mild	18 (50)
Moderate	18 (50)

Table 3. Stratification of various factors with Hypophosphatemia n = 190

Variables	Hypophosphatemia		P-value
	Yes	No	
Age (years)			0.152
0-2	27 (14.3%)	96 (50.5%)	
>2	9 (4.7%)	58 (30.5%)	
Gender			0.975
Male	20 (10.5%)	86 (45.3%)	
Female	16 (8.4%)	68 (35.8%)	
Temperature (C)			0.682
38 - 40	35 (18.5%)	149 (78.4%)	
>40	1 (0.5%)	5 (2.6%)	
Heart Rate			0.136
90 - 120	11 (5.8%)	68 (35.7%)	
>120	25 (13.2%)	86 (45.3%)	
Respiratory Rate			0.343
25 - 40	13 (6.8%)	69 (36.4%)	
>40	23 (12.1%)	85 (44.7%)	
CBC			0.980
1 - 10	12 (6.3%)	51 (26.8%)	
>10	24 (12.6%)	103 (54.3%)	
CRP			0.573
0 - 50	24 (12.6%)	110 (57.9%)	
>50	12 (6.3%)	44 (23.2%)	

4. DISCUSSION

The mean age in our study findings was 2.69 ± 3.19 years. Shazly AN EI, et al. [9] noted, a mean age of 14.3 years while Yoshimatsu S, noted as 18.6 ± 12.3 months [14]. In distribution of gender, 106 (55.8%) were male while 84 (44.2%) were female. Shazly reported more dominance of hypophosphatemia/sepsis and so on more in males in comparison with females [21] whereas Brotfain E, et al noted to have 48 (64%) males and 27 (36%) females [22].

Hypophosphatemia was found in 36 (18.9%) patients in our study. The incidence of hypophosphatemia was noted in 21 (42%) cases in the findings of Shazly AN EI, et al. [9] and 80% in Barak V, et al [10]. The study of Riedler CF noted in 41% cases [13]. A total of 5 children had hypophosphatemia throughout their PICU stay, more than half of our patients had hypophosphatemia at the time of admission [15] while Goldstein J noted in 56% [23] and Polderman KH reported the prevalence of 61% [24]. The findings of present study showed in comparison of previously conducted studies. In our study, stratification of confounders / effect modifiers with respect to hypophosphatemia, insignificant difference was noted in age group ($P=0.152$), gender ($P=0.975$), axillary temperature ($P=0.682$), heart rate ($P=0.136$),

respiratory rate ($P=0.343$), CBC ($P=0.980$) and CRP ($P=0.573$). Hypophosphatemia is linked to a worse prognosis in critically ill adults and children, according to various studies (prolonged length of stay and the need for respiratory support, and higher mortality). However, there was inadequate information as to when hypophosphatemia becomes essential and requires treatment [25].

In an international investigation, 98 (46.4%) of the females had the average age of 10.51 days. The average phosphorus concentration was 4.39 ± 0.67 mg/dL. Phosphorus levels in girls and boys were substantially different. The difference in mean phosphorus between positive and negative blood culture patients was likewise significant. However, phosphorus levels were not linked with type of feeding, ESR, urine culture, or CRP status [26].

5. CONCLUSION

Hypophosphatemia was documented in the considerable number of the children in the clinical sepsis. Serum phosphate levels should be checked in children who present with sepsis. To avoid the issues of a double burden of disease in these patients, the strategies to prevent these diseases should be supported.

6. LIMITATIONS

The sample population represents a single institutional experience; but the study sample can be generalized as the sample came from various areas of Pakistan. Serum procalcitonin should also be done in suspected sepsis children.

ETHICAL APPROVAL

Permission was taken from the ethical review committee of the institute.

CONSENT

Informed consent was taken from the care takers.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Asghar A, Hashmi M, Rashid S, Khan FH. Incidence, outcome and risk factors for sepsis-a two year retrospective study at surgical intensive care unit of a teaching hospital in Pakistan. *J Ayub Med Col Abbottabad*. 2016;28(1):79-83.
2. Martin GS, Mannino DM, Eaton S, Moss M. The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med*. 2003;348(16):1546-54.
3. Angus DC, Linde-Zwirble WT, Lidicker J, Clermont G, Carcillo J, Pinsky MR. Epidemiology of severe sepsis in the United States: analysis of incidence, outcome, and associated costs of care. *Crit Care Med*. 2001;29(7):1303-10.
4. Friedman G, Silva E, Vincent JL. Has the mortality of septic shock changed with time. *Crit Care Med*. 1998;26(12):2078-86.
5. Padkin A, Goldfrad C, Brady AR, Young D, Black N, Rowan K. Epidemiology of severe sepsis occurring in the first 24 hrs in intensive care units in England, Wales and Northern Ireland. *Crit Care Med*. 2003;31(19):2332-8.
6. Miller CJ, Doepker BA, Springer AN, Exline MC, Phillips G, Murphy CV. Impact of serum phosphate in mechanically ventilated patients with severe sepsis and septic shock. *J Intens Care Med*. 2018;1(1):1-9.
7. Haider DG, Lindner G, Wolzt M, Ahmad SS, Sauter T, Leichtle AB, et al. Hyperphosphatemia is an independent risk factor for mortality in critically ill patients: results from a cross-sectional study. *PLoS one*. 2015; 10(8):1-8.
8. Brotfain E, Schwartz A, Boniel A, Koyfman L, Boyko M, Kutz R, et al. Clinical outcome of critically ill patients with thrombocytopenia and hypophosphatemia in the early stage of sepsis. *AnaesthesiolIntensTher*. 2016;48(5):294-9.
9. Shazly AN EI, Soliman DR, Assar EH, Behiry EG, Ahmed IA. Phosphate disturbance in critically ill children: incidence, associated risk factors and clinical outcomes. *Ann Med Surg*. 2017; 21:118-23.
10. Barak V, Schwartz A, Kalickman I, Nisman B, Gurman G, Shoenfeld Y. Prevalence of hypophosphatemia in sepsis and infection: the role of cytokines. *Am J Med*. 1998;104(1):40-7.
11. Bone RC, Fisher JC, Clemmer TP, Slotman GJ, Metz CA, Balk RA. Sepsis syndrome: a valid clinical entity. *Critic care Med*. 1989;17(5):389-93.
12. Shoenfeld Y, Khoury M, Ben-Yehuda O. Hypophosphatemia in sepsis. *Eur J Intern Med*. 1991; 2:67-71.
13. Riedler CF, Scheitlin WA. Hypophosphatemia in septicemia: higher incidence in gram negative than in gram-positive infections. *Br Med J*. 1969; 1(5646):753-6.
14. Yoshimatsu S, Chisti MJ, Hossain M, Islam MM, Fukushima T, et al. Hypophosphatemia among severely-malnourished children: case series. *J Health Pop Nutr*. 2012;30(4):491.
15. Rady HI, Abdel K, Mohamed K. Prevalence and risk factors of hypophosphatemia in pediatric intensive care unit. *J AnesthCrit Care Open Access*. 2014;1:00033.
16. Worley G, Claerhout SJ, Combs SP. Hypophosphatemia in malnourished children during refeeding. *Clin Pediatr*. 1998;37:347-52.
17. Subramanian R, Khardori R. Severe hypophosphatemia. Pathophysiologic implications, clinical presentations, and treatment. *Med*. 2000;79(1):1-8.
18. Manary MJ, Hart CA, Whyte MP. Severe hypophosphatemia in children with kwashiorkor is associated with increased mortality. *J Pediatr*. 1998;133(6):789-91.

19. Heames RM, Cope RA. Hypophosphataemia causing profound cardiac failure after cardiac surgery. *Anaesthes.* 2006;61(12):1211-3.
20. Gaasbeek A, Meinders AE. Hypophosphatemia: an update on its etiology and treatment. *Am J Med.* 2005;118(10):1094-101.
21. Barak V, Schwartz A, Kalickman I, Nisman B, Gurman G, Shoenfeld Y. Prevalence of hypophosphatemia in sepsis and infection: the role of cytokines. *Am J Med.* 1998;104(1):40-7
22. Brotfain E, Schwartz A, Boniel A, Koyfman L, Boyko M, Kutz R, et al. Clinical outcome of critically ill patients with thrombocytopenia and hypophosphatemia in the early stage of sepsis. *Anaesthesiol Intensive Ther.* 2016;48(5).
23. Goldstein J, Vincent JL, Leclerc JL, Vanderhoeft P, Kahn RJ. Hypophosphatemia after cardiothoracic surgery. *Intensive Care Med* 1985;11:144-8.
24. Polderman KH, Bloemers FW, Peerdeman SM, Girbes AR. Hypomagnesemia and hypophosphatemia at admission in patients with severe head injury. *Crit Care Med* 2000;28:2022-5.
25. Blaser AR, Gunst J, Ichai C, Casaer MP, Benstoem C, Besch G, Dager S, Fruhwald SM, Hiesmayr M, Joannes-Boyau O, Malbrain ML. Hypophosphatemia in critically ill adults and children—A systematic review. *Clinical Nutrition.* 2021 Apr 1;40(4):1744-54.
26. Firouzi M, Sherkatolabbasieh H, Garmsiri M. A cross-sectional study on the correlation between blood phosphorus level with sepsis and associated prognostic factors in neonates. *Annals of Medicine and Surgery.* 2022 Apr 7;103582.

© 2022 Qadri et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/87473>