



## **Influence of Nitrogen on Growth Indices of Different Wheat Varieties in Late Sown Condition**

**Kuldeep Singh <sup>a\*</sup>, Anil Kumar Singh <sup>a#</sup>, Ram Pratap Singh <sup>a†</sup>, Vishal Singh <sup>a<sup>o</sup></sup>,  
Ankit Kumar <sup>a<sup>o</sup></sup> and Divya Singh <sup>a<sup>o</sup></sup>**

<sup>a</sup> Department of Agronomy, ANDUAT, Ayodhya, India.

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/IJPSS/2022/v34i2031218

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/89092>

**Original Research Article**

**Received 22 April 2022**

**Accepted 30 June 2022**

**Published 05 July 2022**

### **ABSTRACT**

Growth indices changes from sowing to harvesting of the crop were monitored under the influence of four varieties (PBW- 373, HD- 2285, K-7903, HUW-234) and four Nitrogen levels (0 kg ha<sup>-1</sup> (Control), 120 kg ha<sup>-1</sup>, 150 kg ha<sup>-1</sup> and 180 kg ha<sup>-1</sup>) during rabi season of 2019-20 and 2020-21. Analysis of the two years average data indicated that PBW-373 along with 150 kg N ha<sup>-1</sup> resulted in the highest significant values of Crop Growth Rate (CGR) and Net Assimilation Rate (NAR) up to 60 and 90 DAS respectively after which it declines whereas Relative Growth Rate (RGR) show a declining trend from sowing till harvesting. Net Assimilation Rate (NAR) was found to be unaffected by the varieties but was significantly affected by different levels of nitrogen being highest with 150 kg N ha<sup>-1</sup>. However, it was found to declines from sowing to harvesting. Leaf area duration (LAD) was also found to increases from sowing till harvest being highest with 150 kg N ha<sup>-1</sup>. No interaction effect was found among different treatments. The overall findings concluded that variety PBW-373 along with 150 kg N ha<sup>-1</sup> could be more beneficial in the study area.

*Keywords: Wheat; harvesting; staple food; Growth indices; Nitrogen level.*

<sup>o</sup> Research Scholar,

<sup>#</sup> Associate Professor,

<sup>†</sup> Assistant Professor,

\*Corresponding author: E-mail: [skd5799@gmail.com](mailto:skd5799@gmail.com);

## 1. INTRODUCTION

Wheat (*Triticum aestivum* L.) belongs to the Poaceae family. It is a staple food of the planet. India is one of the main wheat-producing and consuming countries in the world. In India, it is cultivated on an area of 31.62 million hectares having productions of 109.52 million tones with average productivity of 3460 kg ha<sup>-1</sup> [1]. Uttar Pradesh ranks first with reference to the production of 35.50 million tones contributing about 32.42% of the country's production followed by Punjab 17.62 and Madhya Pradesh 17.14 tones ha<sup>-1</sup> [1]. Wheat is very sensitive and responsive to nitrogen fertilization. Nitrogen plays a vital role in increasing the yield of the crop. Application of proper amount of nitrogen is considered key to obtain bumper crop of wheat. High nitrogen supply favours the conversion of carbohydrates into proteins, which in turn promotes the formation of proto-plasm [2]. Nitrogen comprises 7% of total dry matter of plants and is a constituent of many fundamental cell components such as nucleic acids, amino acids, enzymes, and photosynthetic pigments [3].

In addition, nitrogen is also found in chlorophyll, the green coloring agent of plant's leaf. Excessive nitrogen causes succulent growth, resulting in lodging, delayed maturity, and greater susceptibility to diseases and pests. Nitrogen application at the proper dose has had by far the most important effect in terms of increasing crop production.

## 2. MATERIALS AND METHODS

The present research work entitled "Response of nitrogen to different varieties of late sown wheat (*Triticum aestivum* L.) in eastern U.P." was carried out during two successive *Rabi* seasons of 2019-20 and 2020-21 at the Agronomy Research Farm of Acharya Narendra Deva University of Agriculture & Technology, Kumarganj, Ayodhya (U.P.). Ayodhya is situated at about 26°47' North latitude and 82°12' East longitudes with an altitude of location is 113 meters above the Mean Sea Level. The total average rainfall received during crop season was 158 and 0.00 mm during 2019-20 to 2020-21, respectively. The weekly mean maximum temperature ranged from 14.0°C to 36.8°C with an average of 25.4°C during 2019-20 and from 10.1°C to 38.5°C with an average of 24.3°C during 2020-21. The weekly mean relative humidity ranged from 42.8 to 86.1 per cent with an average of 64.4 per cent during 2019-20 and

34.5 to 84.5 per cent with an average 59.4 per cent during 2020-21. The location of the experimental field was same for both the years of investigation. The soil of the experimental site was homogeneous in fertility status with uniform textural makeup. The alluvial soils of Indo-Gangetic plains in general are deep, flat, well drained with low available nitrogen and medium in available phosphorus and potassium. The soil of experimental field was moderate alkaline in reaction (pH) 8.32 and 8.26, low in organic carbon (0.31% and 0.32%), low in available nitrogen (189 kg ha<sup>-1</sup> and 185 kg ha<sup>-1</sup>), phosphorus (16.2 kg ha<sup>-1</sup> and 16.3 kg ha<sup>-1</sup>) and medium in potassium (282 kg ha<sup>-1</sup> and 284 kg ha<sup>-1</sup>) during 2019-20 and 2020-21, respectively. The experiment was laid out in split-plot design with 4 Varieties i.e., PBW- 373, HD- 2285, K- 7903, HUW-234 allocated to main plot and four nitrogen level i.e., 0 kg ha<sup>-1</sup> (Control), 120 kg ha<sup>-1</sup>, 150 kg ha<sup>-1</sup>, 180 kg ha<sup>-1</sup> allocated to sub plot which were replicated thrice. The time of sowing of wheat varieties are 20 december, 2020 and 21 december, 2021 respectively. Half of the nitrogen was applied at the time of sowing while the remaining half applied during the first irrigation. A basal 60 kg/ha dose of phosphorus, 40 kg/ha dose of potassium was applied to all treatment plots at the time of sowing. However, dimensions of the gross plot and net plot size is 4.0 x 5.0 cm<sup>2</sup> and 3.6 x 4.5 cm<sup>2</sup> respectively. The crop received four irrigations in all i.e., CRI, tillering, jointing and flowering stage. Growth indices were calculated based on the formula given by different scientists.

### 2.1 Crop Growth Rate (g m<sup>-2</sup> day<sup>-1</sup>)

The rate of dry matter production per unit land area per unit time or crop growth rate (CGR) was worked out by using the formula proposed by Watson (1947) and expressed as g m<sup>-2</sup> day<sup>-1</sup>.

$$CGR = \frac{W_2 - W_1}{t_2 - t_1} \times \frac{1}{P}$$

Where,

W<sub>1</sub> and W<sub>2</sub> are dry matter of crop (g) at time t<sub>1</sub> and t<sub>2</sub> respectively.

P = Ground area covered by crop (m<sup>2</sup>)

### 2.2 Relative Growth Rate (g g<sup>-1</sup> day<sup>-1</sup>)

The rate of increase in dry weight per unit dry weight of crop expressed in g g<sup>-1</sup> day<sup>-1</sup> was calculated using the formula suggested by Blackman (1919).

$$RGR = \frac{\text{Log } W_2 - \text{Log } W_1}{t_2 - t_1}$$

Where,

$W_1$  and  $W_2$  are dry weight (g) of crop at time  $t_1$  and  $t_2$  respectively.

### 2.3 Net Assimilation Rate (g cm<sup>-2</sup> day<sup>-1</sup>)

It indirectly indicates the rate of net photosynthesis. It is expressed as g of dry matter produced per m<sup>2</sup> of leaf area in a day. For calculating NAR, the total leaf area of the crop has to be used but not the leaf area index. It was calculated at 30, 60 and 90 DAS intervals as per the formula given by Beadle (1987) and expressed in g cm<sup>-2</sup> day<sup>-1</sup>.

$$NAR = \frac{(W_2 - W_1) (\text{Log } LA_2 - \text{Log } LA_1)}{(t_2 - t_1)(LA_2 - LA_1)}$$

Where,

$W_1$  and  $W_2$  are the dry matter accumulation (g) at time  $t_1$  and  $t_2$  respectively.

$LA_1$  and  $LA_2$  are leaf area index at time  $t_1$  and  $t_2$  respectively.

### 2.4 Leaf Area Duration (Day)

To correlate the dry matter yield with LAI, integrated the LAI with time and called as Leaf Area Duration. LAD takes account, both the duration and extent of photosynthetic tissue of the crop canopy. The LAD expresses in days.

$$LAD = (LA_1 + LA_2) \times (T_2 - T_1)/2$$

Where,

$LA_1$  = Leaf Area at time  $T_1$

$LA_2$  = Leaf Area at time  $T_2$

The details of the procedures adopted for raising the crop and criteria used for treatment evaluation and methods adopted during the course of investigation are presented in this chapter.

## 3. RESULTS AND DISCUSSION

A perusal of data reveals that crop growth rate remained unaffected by treatments. However, maximum crop growth rate was recorded under PBW 373 at all stages which is at par with HD-2285 and HUW-234 during both years. Minimum crop growth rate was recorded in variety K 7903. In case of nitrogen level, 150 kg N ha<sup>-1</sup> was recorded significantly higher CGR which is at par

with 180 kg N ha<sup>-1</sup>. The data reveal that crop growth rate was higher during second year of experiment as compare to first year. From the mean data of CGR it can observed that maximum CGR was recorded up 60 DAS after which it declines till harvest. Patra and Ray, [4] reported that high nitrogen levels increased leaf area, leaf number and vegetative growth of plants thus increasing the photosynthetic capacity; consequently, the higher dry matter produced increasing crop growth rate (CGR).

Relative growth rate was found to be higher during initial growth stages and gradually decline further. It was found to be maximum with PBW 373 which is at par with HD 2285 and HUW 234. Variety K 7903 recorded minimum relative growth rate at all stages. However varietal treatment failed to show any significant effect between 60-90 DAS during second year. Different nitrogen level found significant effect on RGR at all growth stages being highest with 150 kg N ha<sup>-1</sup> which is at par with application of 180 kg N ha<sup>-1</sup>. RGR was relatively maximum in second year than first year. However, minimum RGR was recorded under control treatment. These results are confirmatory to those revealed by Ali et al. [5].

A perusal of data reveals that net assimilation rate remained unaffected among varietal treatment. However, maximum NAR was recorded under K 7903 at all stages followed by HUW-234 and HD-2285 during both years. Minimum NAR was recorded in variety PBW 373. Whereas in case of nitrogen level, control treatment recorded significantly higher NAR followed by 120 kg N ha<sup>-1</sup>. The data reveal that NAR was lower during second year of experiment as compare to first year. From the mean data of NAR, it can observe that maximum NAR was recorded upto 60 DAS after which it declines till harvest. NAR decreased significantly with decreases in nitrogen level from 0 kg ha<sup>-1</sup> to 180 kg ha<sup>-1</sup>. This result was in close conformity with Alemu [6].

Leaf area duration found to be lower during initial growth stages and gradually increased further. It was found to be maximum with PBW 373 which is at par with HD 2285. Variety K 7903 recorded minimum LAD at all growth stages. However, varietal treatment failed to show any significant effect during initial days i.e., 0-30 DAS. Different nitrogen level found significant effect on LAD at all growth stages being highest with 150 kg N ha<sup>-1</sup> which is at par with application of 180 kg N

**Table 1. Effect of Nitrogen levels and varieties on crop growth rate (g/m<sup>2</sup>/ day) × 10<sup>-3</sup>**

Treatments	0-30 DAS		30-60 DAS		60-90 DAS	
	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21
<b>Varieties</b>						
HD 2285	2.25	2.34	14.69	15.09	8.47	8.56
PBW 373	2.27	2.36	15.13	15.57	8.71	8.78
K 7903	2.19	2.27	12.53	12.89	7.36	7.43
HUW 234	2.21	2.32	13.99	14.37	8.10	8.17
SEm±	0.04	0.04	0.43	0.46	0.22	0.22
CD at 5%	NS	NS	1.49	1.58	0.76	0.75
<b>Levels of nitrogen</b>						
0 (Control)	2.15	2.22	12.40	12.76	7.28	7.35
120 kg ha <sup>-1</sup>	2.25	2.34	13.99	14.39	8.12	8.19
150 kg ha <sup>-1</sup>	2.45	2.49	17.33	17.85	8.67	8.75
180 kg ha <sup>-1</sup>	2.38	2.44	17.24	17.79	8.57	8.65
SEm±	0.04	0.04	0.35	0.36	0.17	0.17
CD at 5%	NS	NS	1.03	1.06	0.49	0.50

**Table 2. Effect of Nitrogen levels and varieties on relative growth rate (g g<sup>-1</sup> day<sup>-1</sup>) × 10<sup>-3</sup>**

Treatments	30-60 DAS		60-90 DAS	
	2019-20	2020-21	2019-20	2020-21
<b>Varieties</b>				
HD 2285	39.94	40.29	23.03	23.48
PBW 373	41.16	42.33	23.67	23.90
K 7903	34.08	35.07	20.02	21.37
HUW 234	38.06	39.08	22.03	22.66
SEm±	1.17	1.17	0.59	0.80
CD at 5%	4.12	4.13	2.09	NS
<b>Levels of nitrogen</b>				
0 (Control)	33.72	33.94	19.79	21.12
120 kg ha <sup>-1</sup>	38.04	39.12	22.08	22.61
150 kg ha <sup>-1</sup>	40.94	42.11	23.58	23.73
180 kg ha <sup>-1</sup>	40.53	41.60	23.31	23.94
SEm±	0.95	0.96	0.45	0.48
CD at 5%	2.80	2.83	1.34	1.41

**Table 3. Effect of Nitrogen levels and varieties on Net assimilation rate (g/m<sup>2</sup>/ day) × 10<sup>-3</sup>**

Treatments	30 -60 DAS		60 -90 DAS	
	2019-20	2020-21	2019-20	2020-21
<b>Varieties</b>				
HD 2285	54.07	52.49	22.92	23.16
PBW 373	53.58	53.15	23.61	22.66
K 7903	55.82	55.32	25.60	24.18
HUW 234	54.71	54.24	24.78	23.51
SEm±	1.91	1.30	1.54	1.68
CD at 5%	NS	NS	NS	NS
<b>Levels of nitrogen</b>				
0 (Control)	59.70	50.14	29.04	33.59
120 kg ha <sup>-1</sup>	53.17	55.98	23.94	20.65
150 kg ha <sup>-1</sup>	52.36	54.42	22.15	19.43
180 kg ha <sup>-1</sup>	52.96	54.66	21.78	19.84
SEm±	1.49	1.48	1.32	1.31
CD at 5%	4.37	NS	3.88	3.86

ha<sup>-1</sup> and 120 kg N ha<sup>-1</sup>. LAD was relatively more during second year of investigation than first year. However, minimum LAD was recorded under control treatment. These findings are supported by Islam and Jahan [7] in wheat, LAD describes the total amount of leaf area

present over a particular period of growth and is directly correlated with dry matter production. It has been observed that any practice that increases the longevity of green leaves should increase the dry weight of plants.

**Table 4. Effect of tillage options, residue and weed management practices leaf area duration (Days)**

Treatments	0-30 DAS		30-60 DAS		60-90 DAS	
	2019-20	2020-21	2019-20	2020-21	2019-20	2020-21
<b>Varieties</b>						
HD 2285	25.31	26.10	88.20	90.71	135.08	138.30
PBW 373	25.61	26.36	89.66	92.99	137.93	142.64
K 7903	24.60	25.35	81.23	83.96	121.76	125.59
HUW 234	24.86	25.61	84.23	86.63	127.09	130.80
SEm±	0.54	0.71	1.41	1.59	2.06	2.12
CD at 5%	NS	NS	4.87	5.51	7.13	7.34
<b>Levels of nitrogen</b>						
0 (Control)	23.78	24.53	72.15	74.33	101.89	104.81
120 kg ha <sup>-1</sup>	25.35	26.10	88.24	90.60	132.49	136.13
150 kg ha <sup>-1</sup>	25.73	26.48	91.65	94.91	144.49	148.76
180 kg ha <sup>-1</sup>	25.54	26.33	91.28	94.45	142.99	147.63
SEm±	0.47	0.59	1.20	1.92	1.74	2.69
CD at 5%	1.37	NS	3.49	5.59	5.07	7.85

#### 4. CONCLUSION AND RECOMENDATIONS

After summarizing the overall results of this study, it can be concluded that variety PBW 373 in combination with application of nitrogen 150 kg ha<sup>-1</sup> accelerated the growth indices hence growth of the crop.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Anonymous; 2002. Available:www.populationreview.com, Ranking wheat production by country 2022
- Arnon I. Plant Population and Distribution Patterns in Crop Production in Dry Regions. Islamabad, Pakistan: National BookFoundation; 1972;(1)359–360,443–456.
- Bungard RA, Wingler A, Morton JD, Andrews M. Ammonium can stimulate nitrate and nitrite reductase in the absence of nitrate in *Clematis vitalba*. Plant Cell and Environment. 1999;22:859–866.
- Patra B, Ray PK. Response of wheat to various nitrogen levels under late sown condition. Journal of Experimental Agriculture International. 2018;21(1):1-5.
- Ali A, Ahmad A, Syed WH, Khaliq T, Asif M, Aziz M, Mubeen M. Effects of Nitrogen on Growth and yield components of wheat. Sci.Int. (Lahore). 2011;23(4):331-332
- Alemu S. The effect of Nitrogen on the Growth and Dry matter Accumulation of Wheat (*Triticum aestivum* L.) at Hawassa University College of Agriculture Farm. Int. J. Adv. Multidiscip. Res. 2018;5 (3):13-19.
- Islam S, Jahan N. Growth analysis of wheat var. BARI gom-26 as affected by foliar spray of NAA at different nitrogen fertilizer levels. Dhaka Univ. J. Biol. Sci. 2016;25(2):131-138.

© 2022 Singh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/89092>