



The Impact of Pumpkin Seed Extracts on the Histology of the Hypothalamus and Testosterone Level of Alloxan Induced Diabetic Male Rats

Udi Onoriode Andrew ^a, Okoro Ogheneybrorue Godswill ^{a*},
Lucky Omamuzo Ogagayere ^a, Ahama Endurance Efe ^b,
Enaohwo Taniyowho Mamerhi ^b,
Igben Vincent-Junior Onoriode ^b,
Chris-Ozokor LiLian Ebeye ^b and Ogba Grace Oreva ^b

^a Department of Human Anatomy, College of Basic Health Sciences, Achievers University, Owo, Ondo State, Nigeria.

^b Department of Human Anatomy and Cell Biology, Faculty of Basic Medical Sciences, Delta State University, Abraka, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJMAH/2022/v20i12763

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93007>

Original Research Article

Received: 17/08/2022

Accepted: 21/10/2022

Published: 09/11/2022

ABSTRACT

Objective: Epidemiologic studies and randomized clinical trials demonstrate that type 2 diabetes can be substantially avoided by making dietary and lifestyle changes rather than relying heavily on medications. This study evaluates the ameliorating effect(s) of *Telfairia occidentalis* seed

*Corresponding author: E-mail: thomasgodswill23@gmail.com;

administration on the histology of the Hypothalamus and testosterone level in normal and Alloxan-induced diabetes in Wistar rats.

Methods: Twenty male Wistar rats, weighing between 120-180 g, were divided into five groups of 4 rats each. Groups 1 and 2 representing Normal Control (NC) and Diabetic Control (DC) received only feed and water. Group 3 normal rats received 300 mg/kg *Telfairia occidentalis* seed extract; groups 4 and 5 received 50 mg/kg of Metformin and 300 mg/kg of *Telfairia occidentalis* seed extract respectively.

Results: The result revealed that the administration of *Telfairia occidentalis* seed extract moderately increased the body weight ($16.76 \pm 1.75\%$) significantly at ($P < 0.05$) when compared to normal control ($14.06 \pm 2.93\%$) and diabetes significant decrease ($-27.68 \pm 1.92\%$) body weight when compared to metformin-treated ($-6.11 \pm 1.85\%$) and extract ($-5.33 \pm 2.18\%$). The levels of FBG with the extract ($60.70 \pm 4.88\%$) and metformin ($61.24 \pm 6.19\%$) significantly decreased at ($p < 0.05$) when compared to diabetic control ($-64.78 \pm 12.51\%$). The FSH and LH level significantly increased at ($p < 0.05$). Histological changes were also seen in animals of the treated groups.

Conclusion: It is evident that *Telfairia occidentalis* seed extract contains anti-hyperglycemic agents capable of lowering blood glucose levels.

Keywords: Pumpkin seed; hypothalamus; testosterone; diabetic.

1. INTRODUCTION

“Diabetes mellitus (DM) is a complex and multifarious group of disorders that disturb the metabolism of carbohydrates, fat, and protein. It results from a shortage or lack of insulin secretion or reduced sensitivity of the tissue to insulin” [1].

“Type 1 diabetes mellitus is an autoimmune disease caused by the destruction of pancreatic beta cells and characterized by a defect in insulin secretion while Type 2 diabetes mellitus results from abnormalities in insulin secretion and or insulin action or both” [2]. “Dyslipidemia, often presents in diabetic patients and is the main risk factor for cardiovascular and cerebrovascular diseases” [3].

“Diabetes mellitus is currently one of the most costly and burdensome chronic diseases and is a condition that is increasing in epidemic proportions throughout the world. It is a serious illness with multiple complications and premature mortality, accounting for at least 10% of total healthcare expenditure in many countries” [4]. “The prevalence of diabetes in all age groups worldwide is projected to rise from 171 million in 2000 to 366 million in 2030” [5]. “Reasons for this increase include an increase in sedentary lifestyle, consumption of energy-rich diet, obesity, and higher life span” [6].

One of the most commonly used diabetogenic substances in experimental diabetes research is alloxan. It's a cyclic urea analog made up of 2,4,5,6-tetraoxo-hexahydro pyrimidine in its

chemical makeup [7]. Alloxan induces diabetes in animals and inhibits glucose-induced insulin production by β -cells in the pancreas' Islets of Langerhans. Alloxan accumulates rapidly and selectively in β -cells compared to non β -cells, according to research.

“Many traditional plant treatments for diabetes have also been in use but most of the evidence for their beneficial effects has been anecdotal” [8]. Traditional anti-diabetic herbs may give new oral hypoglycemic chemicals, which could help rural communities in underdeveloped nations overcome the high cost and limited availability of current medications.

This study, therefore seeks to evaluate the ameliorating effect(s) of *Telfairia occidentalis* seed administration on the histology of the Hypothalamus and testosterone level in normal and Alloxan-induced diabetes in Wistar rats.

2. METHODS

Telfairia occidentalis seeds were purchased from fruit vendors in Abraka, Ethiope East Local Government Area, Delta State, Nigeria, and was authenticated and identified to the species level by Dr. (Mrs.) Edema Noyo E. of the Department of Botany, Faculty of Science, Delta State University, Abraka. A voucher specimen of *Telfairia occidentalis* was kept in the laboratory. The powder was weighed (300 g) and soaked in 1500ml of ethanol respectively, for 72 hrs [8]. An electrical evaporator extraction unit was used

to obtain the extract (rotary evaporator). The solvent was extracted at 45 degrees Celsius and 60 cm of water pressure. A paste-like extract was obtained and oven-dried to a solid state before being milled into a fine powder.

Metformin (500 mg) as the standard drug was obtained from Omena Pharmacy, Abraka, Ethiope-East Local Government Area, Delta State. The powdered metformin was weighed (0.05 g) in 1000 ml of distilled water and the extract of seed was weighed (3 g) in 10 ml of distilled water to get the appropriate concentrations that were used to administer to the matured healthy adult male rats.

For the diabetic study, hyperglycaemia was induced using Alloxan-monohydrate was dissolved in sodium chloride buffer. Before induction, their fasting blood glucose of the animals was checked after an overnight fast. Alloxan was prepared by dissolving 2 g of Sodium Chloride in 50 ml of water to yield 0.9% of chloride buffer; 3 g of Alloxan was dissolved in 10 ml of chloride buffer to yield 40mg/kg of Alloxan. 1 ml of the resultant solution was injected into the animals through the intraperitoneal and their fasting blood glucose was assessed using ACCU-CHEK active blood glucometer, 72hours after induction. A 200 mg/dl increase in pre-induction fasting Blood glucose level was considered to be diabetic. Diabetes was not induced in animals for the normoglycaemic study.

The research was divided into two phases; the normoglycaemic experimental study with 10 animals divided into 2 groups (n=5).

Group 1 which served as Control.

Group 2 was treated with 300 mg/kg of Crude *Telfairia occidentalis*.

The diabetic study which comprised of 15 animals divided into three (3) groups (n=5).

Group 3 (negative control) induced diabetes but untreated.

Group 4 was induced diabetes and treated with 50 mg/kg of Metformin.

Group 5 was induced diabetes with 40 mg of Alloxan and treated with 300 mg/kg of Crude *Telfairia occidentalis*.

The results were expressed as Mean \pm SEM (Standard Error of the Mean). Data obtained were statistically analyzed using the Student's *t*-Test statistics, one-way analysis of variance (ANOVA), followed by post-hoc Fisher's test for multiple comparisons, using the software, Statistical Package for Social Science (SPSS) version 20 windows software. Significance level was at p values < 0.05 , while p values > 0.05 was considered to be statistically non-significant.

3. RESULTS

From Table 1 the level of body weight in normal and diabetic rats after treatment was reported to be significantly lower in the diabetic group as compared to normal rats treated with 300mg/kg *Telfairia occidentalis* seed and control rats. When diabetic rats were given *Telfairia occidentalis* seed extract (300 mg/kg) and Metformin (50 mg/kg), their bodyweight was considerably higher ($P<0.05$) than diabetic control rats.

The diabetic group had a significantly higher blood glucose level than the normal control rats, as seen in Tables 2 and 3. When diabetic rats were given *Telfairia occidentalis* seed extract (300 mg/kg) and Metformin (50 mg/kg), blood glucose levels were considerably lower ($P<0.05$) than in diabetic control rats. When compared to normal control rats, the diabetic group had significantly lower FSH and LH levels. In diabetic rats, *Telfairia occidentalis* seed extract (300 mg/kg) plus Metformin (50 mg/kg) supplementation significantly increased FSH and LH levels ($P<0.05$) as compared to diabetic control rats.

Plates 1-3 revealed vacuolization of dark cells and neuropil in the hypothalamus of diabetic rats, hypothalamus further revealed lesser vacuolization of dark cells and neuropil in the granular layer and molecular layer. Hypothalamus with *Telfairia occidentalis* seed revealed lesser vacuolization of dark cells and neuropil in the granular layer and molecular layer.

Table 1. The effect of *Telfairia occidentalis* extract on the body of the experimental rat

Groups	Pre-treatment weight	After-treatment weight	% change in body weight
Control	122.77±4.84	142.77±0.77	14.06±2.93a
Diabetic control	197.00±4.26	142.25±1.80	-27.68±1.92b
Diabetic+Metformin	172.75±8.44	162.00±7.07	-6.11±1.85c
TOS extract alone	120.60±3.44	144.85±2.40	16.76±1.75a
Diabetic+TOS extract	160.50±3.28	151.75±1.60	-5.33±2.18c

Values are expressed as mean±SEM. n=4

Table 2. The effect of *Telfairia occidentalis* extract on fasting blood glucose level of experimental rat

Groups	Post-induction Glucose level	Post-treatment Final Glucose level	% change in fasting blood glucose
Diabetic control	247.50±22.72	4400.25±16.82	-64.78±12.51a
Diabetic+Metformin	348.00±31.02	129.50±10.69	-61.24±6.19b
Diabetic+TOS extract	357.75±22.36	139.25±17.71	60.70±4.88b

Values are expressed as mean±SEM. n=4

Table 3. The effect of *Telfairia occidentalis* seed extract on reproductive hormones of experimental rat

S/N	Testosterone
Control	2.33±0.79
Diabetic control	0.60±0.00
TOS extract alone	2.80±0.74
Diabetic+Metformin	1.35±0.72
Diabetic+TOS extract	3.97±0.16

Histopathological Findings of the Hypothalamus:




Plate 1. Negative control




Plate 2. Metformin




Plate 3. *Telfairia occidentalis* seed

4. DISCUSSION

In this study, we discovered a significant reduction in body weight of the diabetic rat agrees with the fact that diabetes causes alteration in the metabolism of macromolecules resulting in changes in body composition and body weight. Significant weight loss is usually evident with the progression of the disease. In this study, *Telfairia occidentalis* seed extract treatment lead to a significant improvement in body weight of diabetic rats compared to the control groups. This is inconsistent with the findings of Ignacimuthu et al. [9] who observed an increase in the body weight of alloxan-induced diabetes in rats treated with *Telfairia occidentalis* seed extract.

The findings from the present investigation showed that 300 mg/kg of the *Telfairia occidentalis* seed extract revealed a significant ($p<0.005$) decrease in the blood glucose level after 14 days compared to diabetic control. Similar to *Telfairia occidentalis* seed extract, Metformin (50 mg/kg) showed a significant ($p<0.005$) decrease in the blood glucose level after 14 days compared to diabetic control, as diabetic rats' glucose levels, progressively increase throughout the experiment. The extract produced $60.70\pm 4.88\%$ glucose reduction while metformin had $61.24\pm 6.19\%$ showing that *Telfairia occidentalis* seed extract reduced glucose levels as the standard drug (metformin) [10]. This result agreed with the finding of Kochhar et al. [10] which stated that "pumpkin

seed extracts had a high content of total phenolics and antioxidant activity coupled with moderate to high alpha-glucosidase and angiotensin converting enzyme inhibitory activities and has the potential to reduce hyperglycemia-induced pathogenesis and associated complications linked to cellular oxidative stress" [10,11,12].

Another important finding from the histology of the hypothalamus in this study is that with the *Telfairia occidentalis* seed extract (300 mg/kg BW) used in the study, there was resurgence in the Hypothalamus which revealed lesser vacuolization of dark cells and neuropil in the hypothalamus in the granular layer and molecular layer. This might explain the poorer blood glucose control observed for diabetic rats treated with that dose of *Telfairia occidentalis* seed extract. Though it is not clear by which way *Telfairia occidentalis* seed extract at the dose of 300 mg/kg BW produced a suboptimal control of blood glucose level, the observation from the study will suggest that beyond a certain dose of *Telfairia occidentalis* seed extract, the control of blood glucose level in the Alloxan-induced diabetic rats will be suboptimal.

5. CONCLUSION

It is evident that *Telfairia occidentalis* seed extract contains antihyperglycemic agents capable of lowering blood glucose level and enhancing testosterone levels. The *Telfairia occidentalis* seed extract showed its ability to restore and reverse the already damaged tissues of alloxan-induced rats.

CONSENT

It is not applicable.

ETHICAL APPROVAL

Permission for the use of the animals and animal protocol was obtained from the Animal Ethics Committee of Delta State University, Abraka with Ethical number: DELSU/CHS/ANA/2021/78 before the commencement of the experiments.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Badifu GI, Akpapunan MA, Mgbemere VM. The fate of beta-carotene in processed leaves of fluted pumpkin, a popular vegetable in Nigerian diet plant foods. *Hum. Nutri.* 1995;48:141-147.
2. Balkau B. The DECODE study, Diabetes epidemiology: Collaborative analysis of diagnostic criteria in Europe. *Diabetes Metab.* 2000;26:282-286.
3. Edem DO, Ekanem IS, Ebong PE. Effect of aqueous extracts of alligator pear seed (*Persea americana* Mill) on blood glucose and histopathology of pancreas in alloxan-induced diabetic rats. *Pak J Pharm Sci.* 2009;22(3):272-276.
4. Dhasarathan P, Hemalatha N, Theriappan P, Ranjitsingh A. Antibacterial activities of extracts and their fractions of leaves of *Tridax procumbens* Linn. *AJBS.* 2011;1: 13-17.
5. Amos AF, Carty MC, Zimmet P. The rising global burden of diabetes and its complications: Estimates and projections to the year. *Diabet Med.* 2010;14:1e85.
6. Zibula SM, Ojewole JA. Hypoglycemic effects of *Hypoxis hemerocallidea* corm 'African potato' methanolic extract in rats. *Medical Journal of Islamic Academy of Sciences.* 2000;13:75-78.
7. Syamsudin I, Winarno H. The effect of inai (*Lawsonia inermis*) leaves extract on blood glucose level: An experimental study. *Research Journal of Pharmacology.* 2008; 2(2):20-23.
8. Deore SL, Khadabadi SS, Daulatkar VD, Deokate UA, Farooqui IU. Evaluation of hypoglycemic and antidiabetic activity of bark of *Butea monosperma*. *Phcog Mag.* 2008;4:134-138.
9. Ignacimuthu S, Ayyanar M, Sivaraman SK. Ethnobotanical investigations among tribes in Madurai district of Tamil Nadu (India). *Journal of Ethnobiology and Ethnomedicine.* 2006;2:1-7.
10. Kochhar A, Nagi M. Effect of supplementation of traditional medicinal plants on blood glucose in non-insulin-dependent diabetics: A pilot study. *Journal of Medicinal Food.* 2008;8:545-549.
11. John CO, Lilian EC, Mamerhi TE, Francisca OO, Vera AO, Onoriode AU. Antioxidative properties of *Ocimum gratissimum* alters lead acetate induced oxidative damage in lymphoid tissues and

- hematological parameters of adult wistar rats. Journal of Toxicological Report. 2021; 215-222.
12. Udi OA, Oyem JC, Ebeye OA, Chris-Ozoko LE, Igbigbi SI, Olannye DU. The effects of aqueous extract of *Ocimum gratissimum* on the cerebellum of male wistar rats challenge by lead acetate. Clinical Nutrition Open Science. 2022;44: 28-41.

© 2022 Andrew et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/93007>