



Effect of Potassium and Gibberellic Acid on the Growth and Yield of Cowpea (*Vigna unguiculata* L.)

Musiboyina Ramya Krishna^{a++*}, Biswarup Mehera^{b#}
and Prateek Kumar^a

^a Department of Agronomy, Naini Agricultural Institute, SHUATS, Prayagraj, Uttar Pradesh - 211007, India.

^b Naini Agricultural Institute, SHUATS, Prayagraj, Uttar Pradesh - 211007, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author MRK designed the study and performed the statistical analysis. Author BM guided the author MRK. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i52818

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97445>

Original Research Article

Received: 03/01/2023

Accepted: 08/03/2023

Published: 11/03/2023

ABSTRACT

The field experiment was conducted at experimental field of the Crop Research Farm, Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, Uttar Pradesh, India. The soil of experimental field is Sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.36%) available N (171.48 kg/ha), available P (27.0 kg/ha) and K (291.2 kg/ha). The Experiment was laid out in Randomized Block Design with nine treatments and one control plot replicated thrice on the basis of one year experimentation. To determine the Effect of Potassium and Gibberellic Acid on the growth and yield of Cowpea (*Vigna unguiculata* L.). The treatments consisted of three levels of Potassium – 20

⁺⁺ M.Sc. Scholar;

[#] Dean;

*Corresponding author: E-mail: mramkri05@gmail.com;

kg/ha, 25 kg/ha and 30 kg/ha and three levels of Gibberellic acid spray – 100 ppm, 120 ppm and 150 ppm. The results showed that treatment with the application of Potassium – 30 kg/ha + GA₃ - 100 ppm was recorded significantly higher Plant height (68.8 cm), no. of nodules/plant (40.9), dry weight (39.32 g). Whereas, maximum Number of pods per plant (16.94), seeds per pod (13.46), seed index (18.36 g), higher seed yield (1459.70 kg/ha) and Haulm yield (2952.30 kg/ha).

Keywords: Potassium; gibberellic acid; cowpea; growth and yield.

1. INTRODUCTION

“Cowpea [*Vigna unguiculata* (L.) Walp] is of immense importance, as it is a multipurpose grain legume extensively cultivated in arid and semi-arid tropics. The cowpea is used as grain, green pods and fodder. Cowpea is grown as a catch crop, weed smothering crop, intercrop, mixed crop and green manure crop. It has ability to fix atmospheric nitrogen in soil at the rate of 56 kg per hectare in association with symbiotic bacteria under favourable conditions” (Yadav, 2007) and Umadevi, G.D. (2019). “Cowpea is the most versatile *kharif* legume because of its drought tolerant character, soil restoring properties and multipurpose use” Patel Hirenkumar et al. [1].

“Potassium has a direct and indirect impact on the plant growth. Using potassium directly causes the reduced transpiration, increasing water absorption or creating internal conditions in order to endure the dryness. The indirect effects take place when using potassium has no value in the plant water relations but based on feeding grounds, it causes the growth increasing. Therefore, the amount that is needed for producing each dry material is being reduced. Studies show that potassium ion gathering in plants before the stresses likes water shortage, coldness, and salinity is insurance for plant survival” Mansouri and Shokoohfar [2].

“Plant growth regulators may be defined as any organic compounds, which are active at low concentrations in promoting, inhibiting or modifying growth and development. The naturally occurring (endogenous) growth substances are commonly known as plant hormones, while the synthetic ones are called growth regulator Range”, V. K. and Giri (2020). “Gibberellins are numerous groups of plant hormones that in addition to auxins are one of the main groups of plant regulators” [3]. “They all differ in physiological activity and structure, and the first identified gibberellins was Gibberellic acid (GA₃). Gibberellins are extensively involved in all phases of plant growth and development, from seed germination to senescence. They promote

seed germination, stimulate stem elongation, leaf expansion, flowering, pollen, and seed development, delay ripening, and inhibit senescence” Mshelmbula et al. (2021).

2. MATERIALS AND METHODS

The present examination was carried out during *Zaid* 2022 at latitude 25° 30'42" N and longitude 81° 00'56" E at Crop Research Farm, Department of Agricultural Sciences, Naini Agricultural Institute, SHUATS, Prayagraj (UP). On average, the elevation is 98 meters above sea level. On 04 April 2022, the crop was sown with the Gomti (V.U-89) variety. The treatments used was Potassium and Gibberellic acid in which Potassium is applied as basal dose and gibberellic acid is applied as foliar spray at pre-flowering and pod filling stages. T₁: Potassium – 20 kg/ha + GA₃ -100 ppm, T₂: Potassium – 20 kg/ha + GA₃ -120 ppm, T₃: Potassium – 20 kg/ha + GA₃ -150 ppm, T₄: Potassium – 25 kg/ha + GA₃ -100 ppm, T₅: Potassium – 25 kg/ha + GA₃ -120 ppm, T₆: Potassium – 25 kg/ha + GA₃ -150 ppm, T₇: Potassium – 30 kg/ha + GA₃ -100 ppm, T₈: Potassium – 30 kg/ha + GA₃ -120 ppm, T₉: Potassium – 30 kg/ha + GA₃ -150 ppm and T₁₀: Control (RDF – N:P:K- 25:50:25 kg/ha). All fertilizers were applied to the soil using urea, single superphosphate (SSP), and potash muriate (MOP). During the 15, 30, 45, and 60 DAS intervals, five randomly selected plant growth symptoms were assessed in each treatment. The mean was compared with a 5% chance of significant results and statistical analysis was performed.

3. RESULTS AND DISCUSSION

3.1 Growth Parameters

The effect of Potassium and Gibberellic Acid on growth of Cowpea growth attributes was shown in Table 1.

3.1.1 Plant height (cm)

Significantly higher plant height (68.8 cm) was recorded with application of Potassium - 30 kg/ha

+ GA₃ - 100 ppm whereas, the treatment Potassium - 25 kg/ha + GA₃ - 100 ppm (67.8 cm) was found to be statistically at par to the treatment with the application of Potassium - 30 kg/ha + GA₃-100 ppm.

“The GA₃ induced at all the growth stages influenced in plant height was attributed to the role of gibberellins in increasing cell elongation and division and internodal elongation” [4,5] reported that “gibberellins increased both cell elongation and division, as evidenced by increase in cell length”. “Gibberellins might have increased cowpea plant height by increasing cell wall extensibility, Behringer et al. [6] reported that in peas, gibberellins decreased the minimum force that will cause cell wall extension. Increased plant height with GA₃ spray may be due to rapid cell elongation in apical region of the plant”. Hirenkumar and Emongor [1] reported same findings who indicated that “exogenous application of GA₃ increased plant height and number of trifoliolate leaves”.

3.1.2 Number of Nodules/plant

Significantly higher nodules/plant (40.9) was recorded with application of Potassium - 30 kg/ha + GA₃ - 100 ppm whereas, the treatment Potassium - 25 kg/ha + GA₃ - 100 ppm (39.5) was found to be statistically at par to the treatment with the application of Potassium - 30 kg/ha + GA₃ - 100 ppm.

“Legume plants develop a symbiotic interaction with rhizobia by forming root nodules in which the bacteria fix atmospheric nitrogen. Nodule formation integrates several developmental processes, such as induction of cortical and pericycle cell division and rhizobia invasion, which are coordinated in time and space” [7]. “Exogenous application of GA₃ has been reported to induce the formation of nodule-like structures on the roots of *Lotus japonicus*” [8].

3.1.3 Dry weight (g/plant)

At 60 DAS, significantly maximum dry weight (39.32 g/plant) was recorded with application of Potassium - 30 kg/ha + GA₃ - 100 ppm whereas, the treatment Potassium – 25 kg/ha + GA₃ - 100 ppm (38.86 g/plant) was found to be statistically at par to the treatment with the application of Potassium - 30 kg/ha + GA₃ - 100 ppm.

Total dry matter was found to be increase gradually from 15 DAS to 60 DAS. The increase of plant dry weight in GA₃ treated plants was

attributed to the increase in leaf area and number of leaves per plant, dry matter accumulation and ultimately induced yield components. These findings are in accordance with those of Deotale et al. [9] in soyabean, in various physiological processes governed under plant levels which ultimately reflected in increased leaf area index, chlorophyll content and net assimilation rate, thus increased of dry weight. These results are in conformity with the results obtained by Fattah [10] in broad bean; Mohandoss and Rajesh (2003) and Emongor [11] in cowpea.

3.2 Yield and Yield Attributes

The effect of Potassium and Gibberellic Acid on growth of Cowpea yield attributes was shown in Table 2.

Significantly Maximum Number of Pods/plant (16.94), Number of seeds/pod (13.6), seed index (18.36 g), Seed yield (1459.70 kg/ha) and Haulm yield (2952.3 kg/ha) were recorded in the treatment application of Potassium - 30 kg/ha + GA₃ - 100 ppm over all the treatments. However, the treatment with (16.57) Potassium - 25 kg/ha + GA₃ - 100 ppm which was found to be statistically at par with Potassium - 30 kg/ha + GA₃ - 100 ppm [12-17].

Significantly highest harvest index (34.15%) was recorded with the treatment application of Potassium - 30 kg/ha + 150 ppm GA₃ over all the treatments. However, the treatments with application of Potassium - 25 kg/ha + GA₃ - 100 ppm (34.04%) and Potassium - 30 kg/ha + GA₃ – 120 ppm (33.58%) in were found to be statistically at par with RDF- N:P:K 25:50:25 kg/ha. Harvest index is directly correlated with seed: straw ratio and is the result of increased in seed and straw yield. Application of urea at pre-flowering and pod initiation stages recorded significantly higher harvest index than the other remaining treatments.

Gibberellic acid can stimulate rapid stem and root growth, induce mitotic division in the leaves of some plants, and increase seed germination and ultimately crop production. Crop yield depends on the accumulation of photo-assimilates during the growing period and the way they are partitioned between desired storage organs of plant. Similar results are reported by Ferdowski Noor et al. (2017) revealed that significantly higher number of pods/plant, seeds/pod, test weight, seed yield and stover yield were recorded under application of gibberellic acid respectively [18-22].

Table 1. Effect of potassium and gibberellic acid on growth of cowpea

S. No.	Treatment Combinations	Plant Height(cm)	No. of nodules	Dry weight (g/plant)
1	Potassium 20 kg/ha + 100 ppm GA ₃	60.9	36.4	35.11
2	Potassium 20 kg/ha + 120 ppm GA ₃	59.8	35.2	34.43
3	Potassium 20 kg/ha + 150 ppm GA ₃	59.2	35.1	33.85
4	Potassium 25 kg/ha + 100 ppm GA ₃	67.8	39.5	38.86
5	Potassium 25 kg/ha + 120 ppm GA ₃	64.8	38.0	36.99
6	Potassium 25kg/ha + 150 ppm GA ₃	62.2	36.9	35.98
7	Potassium 30 kg/ha + 100 ppm GA ₃	68.8	40.9	39.32
8	Potassium 30 kg/ha + 120 ppm GA ₃	65.9	38.5	37.69
9	Potassium 30 kg/ha + 150 ppm GA ₃	63.6	37.2	36.57
10	Control-N:P:K-25:50:25 kg/ha	58.7	34.9	33.03
	F test	S	S	S
	S.E.m (±)	0.34	0.44	0.17
	CD (5%)	1.04	1.31	0.52

Table 2. Effect of potassium and gibberellic acid on yield of cowpea

S. No	Treatments	At Harvest					
		No. of Pods/plant (No)	No. of seeds /pod	Seed index (g)	Seed yield (kg/ha)	Haulm yield (kg/ha)	Harvest index (%)
1	Potassium 20 kg/ha + 100 ppm GA ₃	14.04	11.03	16.80	1208.2	2464.4	32.90
2	Potassium 20 kg/ha + 120 ppm GA ₃	13.49	10.87	16.68	1150.3	2387.9	32.51
3	Potassium 20 kg/ha + 150 ppm GA ₃	12.58	10.39	16.31	1097.0	2285.9	32.43
4	Potassium 25 kg/ha + 100 ppm GA ₃	16.57	13.17	18.17	1433.0	2899.0	33.15
5	Potassium 25kg/ha + 120 ppm GA ₃	15.73	12.72	17.21	1377.4	2669.0	34.04
6	Potassium 25kg/ha + 150 ppm GA ₃	14.53	11.46	17.08	1248.3	2486.9	33.42
7	Potassium 30 kg/ha + 100 ppm GA ₃	16.94	13.46	18.36	1459.7	2952.3	33.09
8	Potassium 30 kg/ha + 120 ppm GA ₃	16.16	12.94	17.43	1403.1	2775.5	33.58
9	Potassium 30 kg/ha + 150 ppm GA ₃	15.17	11.98	17.06	1336.2	2577.0	34.15
10	Control-N:P:K-25:50:25 kg/ha	12.04	10.15	16.12	1029.0	2464.4	31.45
	F test	S	S	S	S	S	S
	S.E.m (±)	0.14	0.09	0.08	9.10	22.10	0.25
	CD (5%)	0.43	0.29	0.27	27.07	65.67	0.75

4. CONCLUSION

On the basis of the results in this study, it is concluded that the application of potassium 30 kg/ha in combination with 100 ppm GA₃ (T₇) was found to be more desirable to be adopted as it recorded higher performance in growth parameter, yield and yield attributes during *zaid* season under Uttar Pradesh conditions.

ACKNOWLEDGEMENT

I express thankfulness to my advisor Dr. Biswarup Mehera and all the faculty members of Department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj -211007, Uttar Pradesh. For providing us essential facilities to undertake the studies.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Hirenkumar Patel D, Patel HC, Sitapara HH, Nayee DD. Influence of plant growth regulators on growth and green pod yield of cowpea [*Vigna unguiculata* (L.) Walp.] cv. Anand veg. Cowpea-1. The Asian J Hort. 2011;6(2):491-5.
2. Mansouri, S, Shokoohfar A. Effect of potassium fertilizer and irrigation intervals levels on yield and yield components of cowpea (*Vigna unguiculata*) in Ahvaz condition. Indian J Fundam Appl Life Sci. 2015;5(1):2231–6345.
3. Bethke PC, Jones RL. Gibberellin signaling. Curr Opin Plant Biol. 1998; 1(5):440-6. DOI: 10.1016/s1369-5266(98)80270-7, PMID 10066618.
4. Taiz L, Zeiger E. Plant Physiology. 3rd ed; Sinauer Associates, inc. Publishers Sunderland Massachusetts. 2002;690.
5. Sauter M, Kende H. Gibberellin-induced growth and regulation of cell division cycle in deep water rice. Plant J. 1992;188: 362-8.
6. Behringer FJ, Cosgrove DJ, Reid JB, Davies PJ. Physical basis for altered stem elongation rates in internode length mutants of *Pisum*. Plant Physiol. 1990; 94(1):166-73. DOI: 10.1104/pp.94.1.166, PMID 11537473.
7. Lievens S, Goormachtig S, Den Herder JD, Capoen W, Mathis R, Hedden P et al. Gibberellins are involved in nodulation of *Sesbania rostrata*. Plant Physiol. 2005;139(3):1366-79. DOI: 10.1104/pp.105.066944, PMID 16258018.
8. Kawaguchi M, Imaizumi-Anraku H, Fukai S, Syono K. Unusual branching in the seedlings of *Lotus japonicus*: gibberellins reveal the nitrogen-sensitive cell divisions within the pericycle on roots. Plant Cell Physiol. 1996;37(4):461-70. DOI: 10.1093/oxfordjournals.pcp.a028968
9. Deotale RD, Maske VG, Sorte NV, Chimurkar BS, Yerne AZ. Effect of GA₃ and NAA on morphophysiological parameters of soyabean. J Soils Crops. 1998;8(1):91-4.
10. Fattah MA, El A. Effect of phosphorus, boron, GA₃, and their interaction on growth, flowering, pod setting, abscission and both green pod and seed yields of broad bean (*Vicia faba* L.) plants. Alexandria Journal of agricultural research. 1997;42(3):311-32.
11. Emongor V. Gibberellic acid (GA₃) influence on vegetative growth, nodulation and yield of Cowpea (*Vigna unguiculata* L. Walp.). J Agron. 2007;6(4):509-17.
12. Deotale RD, Chinmalwar Y, Kalamkar VB, Kamdi SR, Jaybhaye VR. Effect of foliar sprays of cow urine and NAA on chemical, biochemical parameters and yield of pigeon pea. J Soils Crops. 2017;27(2): 159-64.
13. Fisher RA. Discussion of statistical problems in agricultural experimentation; 1950.
14. Khan HR, Bhuiyan MA, Azim F, Rahman MK. Growth, nodulation and uptake of nutrients by cowpea (*Vigna unguiculata* L.) as influenced by phosphorus and potassium fertilization. Current. Agric Res J. 2001;25(1-2):61-6.
15. Kherawat BS, Lal M, Agarwal M, Yadav HK, Kumar S. Effect of applied potassium and manganese on yield and uptake of nutrients by cluster bean (*Cyanosis teragonobola*). J Agric Phys. 2013; 13(1):22-6.
16. Olsen SR, Watanable PS, Dean LA. Estimation of available phosphorus in soil by extraction with sodium bicarbonate. USDA Circular No. 939; 1954.

17. Piper CS. Soil and Plant analysis. Bombay: Hans publisher; 1966.
18. Richards LA. Diagnosis and improvement of saline and alkali soils. USDA Hand book No. 60. Soil Science. 1954;78(2). DOI: 10.1097/00010694-195408000-00012
19. Sirajo et al. Impact of gibberellic acid (GA3) on growth, yield and nodulation on two accessions of cowpea (*Vigna unguiculata* (L.) Walp.). J Appl Sci Environ Manag. 2021;25(8):1435-9.
20. Soil Survey Staff. Soil Taxonomy, a Basic System of Soil Classification for making and interpreting Soil Surveys. USDA Handbook No. 436; 1975.
21. Subbaiah BV, Asija GL. A rapid procedure for the determination of available nitrogen in soils. Curr Sci. 1956;25:259-60.
22. Swaroop K. Effect of phosphorus, potash and rhizobium on pod yield, nutrient uptake and residual available soil NPK of vegetable cowpea. Annals Agric Res. 2006;27(3):250-6.

© 2023 Krishna et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/97445>