



Study of Productivity and Economics of Pearl Millet (*Pennisetum glaucum* L.) Variety NBH-5658 in Relation to Potassium and Plant Growth Regulators

Rathod Sandeep^{a++*}, Rajesh Singh^{a#},
Guggilla Akhil Kumar^{a++} and Addanki Tejaswi^{a++}

^a Department of Agronomy, Naini Agriculture Institute, SHUATS, Prayagraj, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i72866

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97260>

Original Research Article

Received: 17/01/2023
Accepted: 21/03/2023
Published: 28/03/2023

ABSTRACT

Background: Pearl millet can thrive in a wide range of local conditions. In areas where sorghum cannot thrive, pearl millet is a viable alternative. Sorghum and maize can't compete with pearl millet when it comes to soil moisture utilisation and heat tolerance.

Objectives: Effects of Potassium and plant growth regulators on yield and economics of pearl millet.

Methods: With the goal of studying the effect of Potassium and plant growth regulators on yield and economics of Pearl millet (*Pennisetum glaucum* L.) Var. NBH - 5658 under a Randomized block

⁺⁺ M.Sc. Scholar;

[#] Assistant Professor;

*Corresponding author: E-mail: sandeepathod621@gmail.com;

design with 10 treatments (T1-T10). The experimental results revealed that Potassium 60 kg/ha + NAA-100ppm,+Traicontanol-500ppm, produced ear head length (20.50 cm), Number of grains/ear head (1970.0), grain yield (2.72 t/ha), Stover yield (3.78 t/ha) harvest index (41.84 %).T9 Potassium 60 kg/ha + NAA-100ppm,+Traicontanol-500ppm produced greater gross returns (1,36,000.00 INR/ha), net returns (92,840.00 INR/ha), , and benefit-to-cost ratios (2.14).

Keywords: Pearl millet; potassium; plant growth regulators; NAA; triacontanol; yield; economics; attributes; net return; B:C ratio.

1. INTRODUCTION

Pearl millet [*Pennisetum glaucum* (L.)] is a cereal crop, after rice, wheat, maize, and sorghum, that is essential. On 30 million acres of land in the tropical dry and semi-arid region of Asia and Africa, it is a staple meal for 90 million poor people and accounts for half of the world's millet output. After rice, wheat, and maize, it is the fourth most grown crop in India. Several rural communities utilise the bajra crop as a feed grain for their cows and for thatching rooftops. Most bajra is grown in dry, arid regions [1]. A cereal crop, after rice, wheat, maize, and sorghum, that is essential. On 30 million acres of land in the tropical dry and semi-arid region of Asia and Africa, it is a staple meal for 90 million poor people and accounts for half of the world's millet output. After rice, wheat, and maize, it is the fourth most grown crop in India. Several rural communities utilise the bajra crop as a feed grain for their cows and for thatching rooftops. Most bajra is grown in dry, arid regions [2].

Protein (11.6%), iron in particular (8.8%), fat (5%) and carbs (67%) are all abundant in its grains. Most of the world's pearl millet is cultivated in North West India, which also produces 24% of the world's coarse grains and covers 42% of the total area under pearl millet cultivation [2]. Concerns regarding agriculture's capacity to meet the demands of a population that is expanding at an exponential rate have increased due to a lack of new land available for food cultivation and deteriorating soil fertility. When compared to solitary cropping, intercropping increases overall production per unit area per unit time by making efficient use of resources. Planting short-lived crops like pearl millet alongside cluster beans and green gramme crops may increase economic returns per unit of land since there will be less rivalry due to their temporally varying peak resource demand (Bishan Rawat et al, 2018).

A vital main nutrient for plants, potassium serves a variety of purposes. During the grain filling process, potassium turns sugar into starch and

improves water usage efficiency. As a result of its crucial role in controlling stomatal function and the activation of enzymes, potash is frequently depleted in soils when long-term continuous cropping is practised without the addition of additional potash, leaving little exchangeable potassium. In order to complete its life cycle, millet, one of the crops with a high potassium need, needs a significant amount of potassium. The main source for supplying soil with K in the absence of external potassium administration is weathering of the soil [3].

Under environmental stress, plant growth regulators (PGRs) have the potential to boost crop yield. Chemicals known as "growth regulators" can change how organisms grow and develop, which might enhance output, improve grain quality, or make harvesting easier [4]. The amount of nutrients and the use of plant growth regulators had a big impact on Pearl millet's growth metrics. The exogenous use of NAA to boost yield and growth in the face of diverse stresses as salt, drought, extremely high or low temperatures, and heavy metal toxicity. Also, they participate in very significant agronomic developmental processes such as seed germination, leaf angle, blooming duration, and seed yield.

2. MATERIALS AND METHODS

During the Zaid season of 2022, a field experiment was conducted out at the C.R.F of the wing of Agronomy in Shaits Prayagraj, which is located at 25° 24' 42" N latitude, 81° 50' 56" E longitude, and 98 m altitude over the mean sea level (MSL) to see how Pearl millet's growth and productivity are affected by potassium and plant growth regulators (*Pennisetum glaucum* L.). Ten treatments, each replicated three times, were used in the experiment, which was set up using a randomised block design. The size of each plot is 3 metres. The suggested doses for the medication include potassium through, nitrogen via urea, and phosphorus via DAP. Potassium via Muriate of Potash. (T1) Potassium

40 kg /ha + NAA 100 ppm (T2) Potassium 40 kg /ha +Triacontanol 500 ppm, (T3) Potassium 40 kg /ha + NAA 100 ppm + Triacontanol 500 ppm, (T4) Potassium 50 kg /ha + NAA 100 ppm, (T5) Potassium 50 kg /ha + Triacontanol 500 ppm, (T6) Potassium 50 kg/ha + NAA 100 ppm + Triacontanol 500 ppm, (T7) Potassium 60 kg/ha + NAA 100 ppm, (T8) Potassium 60 kg/ha + Triacontanol 500 ppm, Treatment (T9) Potassium 60 kg/ha + NAA 100 ppm + Triacontanol 500 ppm (T10) N-P-K (RDF 80;40;40) /ha. The Stover production from each online plot was measured and expressed in tonnes per hectare after 10 days of full drying in the sun. Using, the statistics were computed and examined. The Gomez and Gomez [5] statistical approach.

3. RESULTS AND DISCUSSION

3.1 Yield and Yield Attributes

3.1.1 Ear head length (CM)

The statistical analysis of ear head length revealed the enormous impact of ear head period. The treatment of 60 kg K/ha+100 PPM NAA + 500 PPM Triacontanol /ha. resulted in a significant and maximal ear head length (20.50 cm). However, with 60 kg K/ha+100 PPM NAA + 500 PPM Triacontanol /ha., no other treatment achieved statistical parity.. might due to the presence of Potassium in that application enhances the development of strong cell walls [6] and improves germination of pollen in the florets which leads to high spikelet fertility [7]. The application of triacontanol and NAA was attribute to an increased rate of photo synthetic activity accelerated transport and efficiency of utilization photosynthetic products thus result resulting cell elongation and rapid cell growing portion, increase grain yield the plant" [8].

3.2 Number of Grains in the Ear

The statistical analysis of the amount of grains per ear head revealed a significant influence. Significant and the largest number of grains per ear head were recorded in the treatment of 60 kg K/ha+100 PPM NAA + 500 PPM. The statistical parity between Potassium 60 kg/ha + Triacontanol 500 ppm and 60 kg K/ha+100 PPM NAA + Triacontanol 500 PPM was achieved, The improvement of number of grains per ear head with the application of Potassium 50 kg/ha and Sulphur 20 kg/ha was might due to The presence

of potassium might be attributed to better filling of grains [9] The application of triacontanol and NAA reported As a result of increased PGR and other nutrient supplies to plants, the overall improvement in plant development can be linked to application. Plants may have benefited from an earlier supply of nutrients during the floral primordial initiation stage, resulting in a greater number of functional tillers and ultimately more grains/ear heads. These findings are also consistent with those [8].



3.3 Grain Yield

Different combinations of Potassium & PGR can have a significant effect on grain production. A grain yield of 2.72 ta/ha was obtained with a treatment of Potassium 60 kg/ha + NAA 100 ppm + Triacontanol 500 ppm however, Potassium 60 kg/ha + NAA 100 ppm + Triacontanol 500 ppm yielded results statistically equivalent to those of (Potassium 60 kg/ha + NAA 100 ppm + Triacontanol 500 ppm). might due to the presence of potassium in that application stimulates the cumulative effect of improvement in yield attributes viz., number of effective tillers per plant, ear head length and thickness and test weight [10] and increased availability, absorption, and translocation of K nutrient [11]. Increased availability of nutrients to plants at the flower primordial initiation stage which might have helped in greater effective tillers formation and ultimately increased number of grains/ear head and ear head length [12]. A significant improvement in biological yield can therefore be attributed to the better grain production characteristics. These findings are also consistent with those o Sivakumar et al. [13].



3.4 Stover Yield

The stover yield output of the pearl millet crop had also been greatly altered by the treatment of Nitrogen & Phosphorus. In terms of stover yield (3.78 t/ha), the highest was observed, Potassium 60 kg/ha + NAA 100 ppm + Triacantanol 500 ppm. however, (Potassium 60 kg/ha + Triacantanol 500 ppm) was shown to be statistically equivalent to , Potassium 60 kg/ha + NAA 100 ppm + Triacantanol 500 ppm) With the addition of potassium and PGR , pearl millet

yielded substantially the application of Potassium 60 kg/ha + NAA 100 ppm + Triacantanol 500 ppm might due to the presence of potassium in that application enhances the development of strong cell walls and therefore stiffer straw which might be resulted into profuse tillering. These results are agreement with those reports by [10] and increased availability, absorption, and translocation of K nutrient [11]. The application of triacantanol and NAA reported As increased growth parameters and yield related features. Stover yield result accelerated nutrient uptake which helped the plants to put optimum growth, Stew yield are improved [14].

3.5 Economics

Among the different combination of nutrient source Potassium 60 kg/ha + NAA 100 ppm + Triacantanol 500 ppm recorded higher gross return (1,36,000.00 INR/ha) net return (92,840.00 INR /ha), and benefit: cost ratio (2.14) [15].



Table 1. Effect of potassium and plant growth regulators on ear head length (cm), no of grains/ ear head (g) and stover yield (t/ha) and harvest index(%) of pearl millet

S. No	Treatments	Ear head length (cm)	No of Grains/ ear head	Grain yield (t/ha)	Stover yield (t/ha)	Harvest Index (%)
1	Potassium 40 kg/ha + NAA 100 ppm	16.15	1621.00	2.33	3.25	41.75
2	Potassium 40 kg/ha + Triacantanol 500 ppm	16.75	1681.00	2.18	3.44	38.79
3	Potassium 40 kg/ha + NAA 100 ppm + Triacantanol 500 ppm	17.53	1759.00	2.24	3.37	39.75
4	Potassium 50 kg/ha + NAA 100 ppm	17.07	1783.00	2.40	3.58	40.14
5	Potassium 50 kg/ha + Triacantanol 500 ppm	18.30	1856.00	2.56	3.67	41.09
6	Potassium 50 kg/ha + NAA 100 ppm + Triacantanol 500 ppm	19.50	1868.00	2.42	3.64	39.93

S. No	Treatments	Ear head length (cm)	No of Grains/ ear head	Grain yield (t/ha)	Stover yield (t/ha)	Harvest Index (%)
7	Potassium 60 kg/ha + NAA 100 ppm	18.20	1945.00	2.67	3.72	41.78
8	Potassium 60 kg/ha + Triacantanol 500 ppm	20.40	1964.00	2.69	3.74	41.80
9	Potassium 60 kg/ha + NAA 100 ppm + Triacantanol 500 ppm	20.50	1970.00	2.72	3.78	41.84
10	Control 80:40:40 N:P:K	15.91	1521.00	2.20	3.15	41.12
	F test	S	S	S	S	S
	SEm±	0.06	5.16	8.35	6.81	0.47
	CD (P = 0.05)	0.19	15.34	0.02	0.11	0.74

Table 2. Effect of potassium and plant growth regulators on economics of pearl millet

S.No	Treatments	Cost of Cultivation (INR/ha)	Gross return (INR)	Net return (INR/ha)	B:C ratio
1	Potassium 40 kg/ha + NAA 100 ppm	42,790.00	1,16,500.00	73,710.00	1.72
2	Potassium 40 kg/ha + Triacantanol 500 ppm	42,840.00	1,09,000.00	66,160.00	1.54
3	Potassium 40 kg/ha + NAA 100 ppm + Triacantanol 500 ppm	42,940.00	1,12,000.00	69,060.00	1.60
4	Potassium 50 kg/ha + NAA 100 ppm	42,950.00	1,20,000.00	77,050.00	1.79
5	Potassium 50 kg/ha + Triacantanol 500 ppm	43,000.00	1,28,000.00	85,000.00	1.97
6	Potassium 50 kg/ha + NAA 100 ppm + Triacantanol 500 ppm	43,100.00	1,21,000.00	77,900.00	1.80
7	Potassium 60 kg/ha + NAA 100 ppm	42,950.00	1,33,500.00	90,400.00	2.10
8	Potassium 60 kg/ha + Triacantanol 500 ppm	43,160.00	1,34,500.00	91,550.00	2.12
9	Potassium 60 kg/ha + NAA 100 ppm + Triacantanol 500 ppm	43,260.00	1,36,000.00	92,840.00	2.14
10	Control 80:40:40 N:P:K	42,050.00	1,10,000.00	67,950.00	1.61

4. CONCLUSION

Treatment Potassium 60 kg/ha + NAA 100 ppm + Triacantanol 500 ppm. produced the highest grain yield (2.72 ta/ha), gross return (1,36,000.00 INR/ha), net return (92,840.00 INR/ha), and benefit: cost ratio (2.14), which may be more preferable for farmers because it is more economically profitable and thus can be recommended to farmers.

ACKNOWLEDGEMENTS

I'd like to thank my advisor, Dr. RAJESH SINGH, for his unwavering support, direction, and insightful suggestions for increasing the quality of this work Mr. Guggilla Akhil Kumar, Miss Addanki Tejaswi., who has been a continual source of inspiration, as well as all of the faculty members of the Department of Agronomy, SHUATS, Prayagraj, Uttar Pradesh (U.P), India, are to be thanked for providing required facilities, as well

as their collaboration, encouragement, and support.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Mannepu Venkata Sai Ganesh, Rajesh Singh, Thakur Indu, Kalyani D. Pearl millet (*Pennisetum glaucum* L.,) Var ABV-04 influenced by nitrogen & phosphorus effects on growth parameters and yield. International Journal of Plant and Soil Science. 2022;34(14):2320-7035.
2. Agricultural Statistics at a Glance; 2021.
3. Srinivasarao Ch, Sumanta Kundu Bk. Ramachandrappa, Sharanhoopal Reddy, Rattan Lal, Venkateswarlu B, Sahrawat KL, Prakash Naik R. Potassium release

- characteristics, potassium balance and finger millet yield sustainability in a 27 – year long experiment on an (*Pennisetum glaucum*) - Alfisol in the semi-arid tropical India; 2013.
4. Espindula MC, Rocha VS, Grossi JAS, Souza MA, Souza LT, Favarato LF. Use of growth retardants in wheat. *Planta Daninha*. 2009;27:379-387.
 5. Gomez KA, Gomez AA. Statistical procedures for agricultural research. 2nd edn. John Wiley and Sons, Inc. London, UK; 1984.
 6. Chaudhry NN, Khaif HR, Raj AD, Yadav V, Yadav P. Effect of nutrients (K and S) on growth, yield and economics of pearl millet [*Pennisetum glaucum* (L.)]. *International Journal of Forestry and Crop Improvement* 2014;5(1):9-12.
 7. Chouhan S, Naga SR, Bhadru P, Koli DK, Kumar A, Jaiswal. Effect of bioorganics and potassium on yield attributes of Pearl millet [*Pennisetum glaucum* (L.) R. Br. emend Stuntz]. *International Journal of Chemical Studies*. 2018;6(2):2038-2041
 8. Suresh G, Guru G, Subbalakshmi L. Effect of plant growth regulators and nutrient levels and yield attributes and yield of pearl millet. *Indian Journal of Pharmacognosy and Phytochemistry*. 2020; 9(5):311–306.
 9. Heidari M, Jamshid P. Interaction between salinity and potassium on grain yield, carbohydrate content and nutrient uptake in pearl millet. *ARPJN Journal of Agricultural and Biological Science*. 2010;5(6):39-46.
 10. Kacha DJ, Khafi HR, Mehta AC, Shekh MA, Jadav RP. Effect of potassium and zinc on yield and quality of rabi pearl millet (*Pennisetum glaucum*). *Crop Research*. 2011;41(1, 2 & 3):31-34.
 11. Jat G, Majumdar SP, Jat NK, Mazumdar SP. Effect of potassium and zinc on crop yield, nutrient uptake and distribution of potassium and zinc fractions in Typic Ustipsamment. *Indian journal of Agricultural Sciences*. 2014;84(7):832-8.
 12. Kunduru Manikanteswara Reddy, Umeha, C, Meshram MR. Impact of potassium and sulphur levels on pearl millet (*Pennisetum glaucum* L. biological forum. *An International Journal*. 2021;13(1):92-97.
 13. Sivakumar R, Pathmanaban G, Kalarani MK, Mallika V, Srinivasan PS. Effect of foliar application of growth regulators on biochemical attributes grain yield in pearl millet. *Indian Journal of Plant Physiology*. 2002;7(1):79-82.
 14. Adey Mourya, Shikha Singh. Effect of bio-fertilizers and plant growth regulators on growth and yield of Pearl millet (*Pennisetum glaucum* L.). *The Pharma Innovation Journal*. 2022;11(4): 1791-1794.
 15. Hussain N, Khan AZ, Akbar H, Bangash NG, Khan ZH, Idrees M. Response of maize varieties to phosphorus and potassium levels. *Sarhad Journal of Agriculture*. 2007;23(4):881-887.

© 2023 Sandeep et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/97260>