



Electron Transfer Reaction of Theobromine and Permanganate Ion in Aqueous Acidic Media

Frank Jones^a, Idongesit Bassey Anweting^{b*}
and Idongesit Edem Okon^{a,c}

^a Department of Chemistry, Ahmadu Bello University, Zaria, Nigeria.

^b Chemistry Department, University of Uyo, Uyo, Nigeria.

^c Transport Technology Centre, Nigerian Institute of Transport Technology, Zaria, Kaduna State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJACR/2023/v13i2242

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98168>

Original Research Article

Received: 01/02/2023

Accepted: 03/04/2023

Published: 15/04/2023

ABSTRACT

Electron transfer reaction of theobromine (hereafter referred to as TB) and permanganate ion has been studied using spectrophotometric technique under pseudo first-order conditions at $T = 24 \pm 0.1^\circ\text{C}$, $[\text{H}^+] = 3.2 \times 10^{-1} \text{ mol dm}^{-3}$ and $I = 1.0 \text{ mol dm}^{-3}$ (Na_2SO_4). Stoichiometric study depicted that two moles of TB were consumed by one mole of permanganate ion. The rate of the reaction is first order in both $[\text{TB}]$ and $[\text{MnO}_4^-]$ and second order overall. It was observed that, $[\text{H}^+]$ and ionic strength respectively enhanced the rate of the reaction in the range studied. Michaelis-Mentens analysis showed no evidence of intermediate complex formation. Added anions and cations were observed to catalyze the reaction rate. The reaction conforms to the rate law as shown below:

$$-d[\text{MnO}_4^-]/dt = (a+b[\text{H}^+])[\text{TB}][\text{MnO}_4^-]$$

*Corresponding author: Email: idongesitanweting@uniuyo.edu.ng;

where $a = 0.041 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ and $b = 0.028 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$.

The reaction thus occurs by an outer-sphere pathway and plausible mechanism is proposed for the reaction.

Keywords: Electron transfer; theobromine; permanganate ion and spectrophotometric.

1. INTRODUCTION

“Electron transfer (ET) is one of the unique chemical processes which have received considerable attention due to its role in physical and biochemical systems” [1]. “It occurs when an electron moves from an atom or a chemical species (e.g. a molecule) to another atom or chemical species. The development of the field, experimentally and theoretically, as well as its relation to the study of other kinds of chemical reactions, presents an intriguing history, one in which many threads have been brought together” [2]. “The process is a mechanistic description of the thermodynamic concept of redox, wherein the oxidation states of both reaction partners changed. Numerous biological processes involve ET reactions, these processes include oxygen binding, photosynthesis, respiration, and detoxification. ET reactions commonly involve transition metal complexes, but there are now many examples of ET in organic chemistry and other discipline or field of study” [3-5]. “Moreover, many reactions in bioinorganic systems involve the electron transfer at one stage or the other and proper understanding of these electron transfer processes would help in the understanding development and eventual effective control of a wide area of science and technology” [6].

“Theobromine also known as xantheose is an alkaloid of the methylxanthine family which can be found in chocolate, kolanuts, tea leaves and other food substances. In modern medicine, it is used as a vasodilator, a diuretic and as a heart stimulant” [7]. “It is also known to exhibit antioxidant and pro-oxidant properties” [8]. “It is slightly water soluble but more fat soluble” [9]. “Theobromine is a purine alkaloid which can be derived from a nucleoside called xanthosine. Cleavage of the ribose and N-methylation yields 7-methylxanthosine. 7-methylxanthosine in turn is the precursor to theobromine” [10].

The electron transfer reactions of permanganate ion have been reported by several authors [11-15]. Permanganate undergoes several electron transfer reactions with organic and inorganic compounds in both acidic and alkaline

medium [16]. The extent of oxidation by permanganate ion depends on the nature of the reductant and the pH of the reaction medium. In acidic medium Mn VII exists as HMnO_4 , H_2MnO_4^+ , HMnO_3 , and Mn_2O_7 . There exist paucity of data on electron transfer reaction between permanganate ion and theobromine and it is hopeful that the data that shall be generated from this study will provide more information about theobromine (alkaloid).

2. MATERIALS AND METHODS

2.1 Materials

Accurately, 3.60g (BDH) of theobromine was dissolved in 200 cm^3 of distilled water to give $1.0 \times 10^{-1} \text{ mol dm}^{-3}$ solution. Standard solutions of potassium permanganate (KMnO_4) and sulfuric acid (H_2SO_4) were also prepared, and standardized. Standard solutions of sodium sulfate (Na_2SO_4), sodium formate (HCOONa), magnesium sulfate (MgSO_4), calcium sulfate (CaSO_4) and sodium nitrate (NaNO_3) were prepared by dissolving accurately weighed quantity of the solute in a given volume of distilled water.

2.2 Methods

2.2.1 Stoichiometry

The stoichiometry of the reaction was determined spectrophotometrically using the mole ratio method [17-21]. Permanganate concentration was kept constant while that of the TB was varied in the range, $[\text{TB}] = (1.0 - 3.8) \times 10^{-2} \text{ mol dm}^{-3}$ at $[\text{MnO}_4^-] = 2.0 \times 10^{-4} \text{ mol dm}^{-3}$, $[\text{H}^+] = 1.0 \times 10^{-2} \text{ mol dm}^{-3}$, $l = 0.50 \text{ cm}$ (Na_2SO_4) and $\lambda_{\text{max}} = 530 \text{ nm}$. The reactions were allowed to go to completion, when the absorbances of the solutions of the reaction mixture remained constant. The stoichiometry was indicated by the point of inflexion, was evaluated from the plots of absorbance versus mole ratio ($[\text{TB}] / [\text{MnO}_4^-]$) [22].

2.2.2 Kinetic studies

The rate of the reaction was monitored by measuring decreased in absorbances of the

permanganate at 530nm using Corning Colorimeter 252 model. Kinetic measurements were made under pseudo-first order conditions with [TB] at least 10-folds excess of $[\text{MnO}_4^-]$. The pseudo first order rate constant (k_{obs}) in each case was obtained from the slope of the plots of $(\log A_t - A_\infty)$ versus time [23, 19, 24]. The second order rate constant (k_2) was obtained as $k_{\text{obs}}/[\text{TB}]$ at $[\text{TB}] = 4.0 \times 10^{-3} - 2.8 \times 10^{-2} \text{ mol dm}^{-3}$, $[\text{H}^+] = 3.2 \times 10^{-1} \text{ mol dm}^{-3}$, $I = 1.0 \text{ mol dm}^{-3}$ (Na_2SO_4) and $[\text{MnO}_4^-] = 2.0 \times 10^{-4} \text{ mol dm}^{-3}$.

2.2.3 Effect of change in hydrogen ion concentration, $[\text{H}^+]$ on the reaction rate

The effect of $[\text{H}^+]$ on the rate of the reaction was investigated using sulfuric acid in the range $[\text{H}^+] = 0.24 - 0.64 \text{ mol dm}^{-3}$, while the concentrations of other reactants were kept constant at, $I = 1.0 \text{ mol dm}^{-3}$ (Na_2SO_4) and $\lambda_{\text{max}} = 530\text{nm}$ [19, 20].

2.2.4 Effect of ionic strength (I) on the rate

The effect of ionic strength of the reaction medium was investigated by keeping the concentration of TB and MnO_4^- constant and varying the ionic strength of the reaction medium in the range $I = 0.2-1.4 \text{ mol dm}^{-3}$ (Na_2SO_4) and $\lambda_{\text{max}} = 530\text{nm}$ [25-27, 24].

2.2.5 Effect of added cations and anions on the rate of reaction

The effect of added ions in the reaction medium was carried out by adding various concentrations of Mg^{2+} , Ca^{2+} , NO_3^- and HCOO^- to the reaction medium while all other conditions remained constant [28, 25, 19], at $I = 1.0 \text{ mol dm}^{-3}$ (Na_2SO_4), $[\text{MnO}_4^-] = 2.0 \times 10^{-4} \text{ mol dm}^{-3}$, $[\text{TB}] = 2.0 \times 10^{-2} \text{ mol dm}^{-3}$ and $[\text{H}^+] = 3.2 \times 10^{-1} \text{ mol dm}^{-3}$.

2.2.6 Free radical test

About 5cm^3 of acrylamide was added to partially oxidized reaction mixture containing various concentrations of solution of theobromine, permanganate ion and hydrogen ion. This was followed by a large excess of methanol [29-30]. The acrylamide was also added to the solution of theobromine and permanganate ion separately, serving as control.

2.2.7 Test for intermediate complex formation


Spectroscopic test was carried out by comparing the electronic spectrum of the reaction mixture

20 minutes after the start of the reaction with that of the permanganate ion within a wavelength of 400-700nm. Kinetic test was also investigated using Michaelis-Mentens plot of $1/k_{\text{obs}}$ versus $1/[\text{TB}]$ [12, 13].

3. RESULTS AND DISCUSSION

3.1 Stoichiometry and Product Analysis

The stoichiometry studies showed that one mole of MnO_4^- was consumed by two moles of TB as represented by the equation 1.




Similar stoichiometry has been reported for the oxidation of DL-alanine by permanganate ion [9].

However, in the reaction of permanganate ion with malachite green, nicotinic acid, and L-tryptophan one mole of permanganate was consumed by one mole of each reductant [22, 11, 12, 14]. Mn^{2+} , a product of oxidation of MnO_4^- was confirmed qualitatively by reacting the product with indole [31]. Formation of white precipitate is an indication of the presence of Mn^{2+} .

3.2 Kinetic Study

The plots of $(\log A_t - A_\infty)$ versus time were linear to more than 80% extent of reaction indicating that the reaction is first order dependence on $[\text{MnO}_4^-]$. A typical pseudo-first order plot is presented in Fig. 1. The pseudo-first order rate constants were determined at different initial concentrations of TB. Plot of $\log k_{\text{obs}}$ versus $\log [\text{TB}]$ at, $I = 1.0 \text{ mol dm}^{-3}$, $[\text{H}^+] = 3.2 \times 10^{-1} \text{ mol dm}^{-3}$ gave a slope of 0.94 (Fig. 2). This suggests a first order dependence on [TB]. The values of second order rate constant $k_2 = k_{\text{obs}}/[\text{TB}]$ were constant (Table 1) suggesting that the reaction is also first order in [TB] and that the reaction is second order overall, therefore conforming to the rate equation below:



Similar order kinetics have been reported in the reduction reactions of permanganate ion by aliphatic aldehydes [32], acetaldelydes [16], DL-alanine [33], L- asparaginase [34]) and L-tryptophan [14].

3.3 Effect of $[H^+]$ on the Rate of the Reaction

The rate of the reaction was enhanced by increase in $[H^+]$ at $[H^+] = 0.24 - 0.64 \text{ mol dm}^{-3}$, $[TB] = 2.0 \times 10^{-2} \text{ mol dm}^{-3}$, $I = 1.0 \text{ mol dm}^{-3}$ (Na_2SO_4), $[\text{MnO}_4^-] = 2.0 \times 10^{-4} \text{ mol dm}^{-3}$ and $\lambda_{\text{max}} = 530\text{nm}$ and $T = 24 \pm 0.1^\circ\text{C}$. Plot of $\log k_{\text{obs}}$ versus $\log[H^+]$ gave a slope of 1.31, indicating that the reaction is first order with respect to $[H^+]$. Plot of k_2 versus $[H^+]$ was linear with positive intercept. The rate equation of the reaction as a function of H^+ can be represented by equation 3

$$-d[\text{MnO}_4^-]/dt = (a+b[H^+]) [TB] [\text{MnO}_4^-] \quad (3)$$

where $a = 0.041 \text{ dm}^3\text{mol}^{-1} \text{ s}^{-1}$ and $b = 0.028 \text{ dm}^3\text{mol}^{-1} \text{ s}^{-1}$.

The rate of the reaction showed two parallel pathways of reaction; one involving the protonated and the other the unprotonated species both reacting to give a product [35]. Similar results were reported in the reaction of permanganate ion with L- ascorbic acid [36], and malachite green [22].




Fig. 1. Typical pseudo-first order plot for the redox reaction of TB with MnO_4^- , $[TB] = 1.0 \times 10^{-2} \text{ mol dm}^{-3}$, $[\text{MnO}_4^-] = 2.0 \times 10^{-4} \text{ mol dm}^{-3}$, $[H^+] = 1.0 \times 10^{-2} \text{ mol dm}^{-3}$, $I = 0.5 \text{ mol dm}^{-3}$, $\lambda_{\text{max}} = 530\text{nm}$ and $T = 23.0 \pm 0.1^\circ\text{C}$




Fig. 2. Plot of $\log k_{\text{obs}}$ versus $\log [TB]$ at $[\text{MnO}_4^-] = 2.0 \times 10^{-4} \text{ mol dm}^{-3}$, $[TB] = (2.0 - 3.8) \times 10^{-2} \text{ mol dm}^{-3}$, $I = 1.0 \text{ mol dm}^{-3}$ (Na_2SO_4) and $T = 24 \pm 0.1^\circ\text{C}$

Table 1. Pseudo-first order and second order rate constants for the reaction of TB and MnO_4^- at $[\text{TB}] = 2.0 \times 10^{-2} \text{ mol dm}^{-3}$, $[\text{MnO}_4^-] = 2.0 \times 10^{-4} \text{ mol dm}^{-3}$, $T = 24.0 \pm 0.1^\circ\text{C}$, $I = 1.0 \text{ mol dm}^{-3}$ and $\lambda_{\text{max}} = 530 \text{ nm}$

| $10^3 [\text{TB}]$, mol dm^{-3} | $10^1 [\text{H}^+]$, mol dm^{-3} | $10 [\text{I}]$, mol dm^{-3} | $10^3 k_{\text{obs}}$, s^{-1} | k_2 , $\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ |
|--|---|---|--|--|
| 4.00 | 3.20 | 1.00 | 0.380 | 0.095 |
| 8.00 | 3.20 | 1.00 | 0.736 | 0.092 |
| 12.0 | 3.20 | 1.00 | 1.100 | 0.092 |
| 16.0 | 3.20 | 1.00 | 1.470 | 0.092 |
| 20.0 | 3.20 | 1.00 | 2.020 | 0.101 |
| 24.0 | 3.20 | 1.00 | 2.200 | 0.091 |
| 28.0 | 3.20 | 1.00 | 2.820 | 0.100 |
| 20.0 | 2.40 | 1.00 | 1.590 | 0.079 |
| 20.0 | 3.20 | 1.00 | 2.300 | 0.110 |
| 20.0 | 4.00 | 1.00 | 3.070 | 0.153 |
| 20.0 | 4.80 | 1.00 | 3.720 | 0.186 |
| 20.0 | 5.60 | 1.00 | 4.790 | 0.239 |
| 20.0 | 6.40 | 1.00 | 5.860 | 0.293 |
| 20.0 | 3.20 | 0.20 | 1.220 | 0.060 |
| 20.0 | 3.20 | 0.60 | 1.480 | 0.074 |
| 20.0 | 3.20 | 0.80 | 1.660 | 0.083 |
| 20.0 | 3.20 | 1.00 | 2.490 | 0.120 |
| 20.0 | 3.20 | 1.40 | 3.260 | 0.160 |




Fig. 3. $\text{Log}k_2$ versus $I^{1/2}$ plot for the oxidation of TB by MnO_4^- at $[\text{TB}] = 1.0 \times 10^{-2} \text{ mol dm}^{-3}$, $[\text{MnO}_4^-] = 2.0 \times 10^{-4} \text{ mol dm}^{-3}$, $[\text{H}^+] = 3.2 \times 10^{-1} \text{ mol dm}^{-3}$, $I = 0.2 - 1.4 \text{ mol dm}^{-3}$ (Na_2SO_4) and $T = 24 \pm 0.1^\circ\text{C}$

3.4 The Effect of Ionic Strength on the Rate of the Reaction

The increase in ionic strength of the reaction medium resulted in increase in the rate of the

reaction within the concentration range studied, the positive salt effect observed for the reaction showed the interaction of like charges (Bronsted-Debye positive salt effect) in the activated complex. Similar results were obtained by [17,

36)]. Linear plot was obtained for $\log k_2$ versus $I^{1/2}$ (Fig. 3).

3.5 Effect of Added Ions on the Rate of the Reaction

The added ions NO_3^- , HCOO^- , Mg^{2+} and Ca^{2+} increased the rate of the reaction as shown on Table 2. "Catalysis of the reaction by these added anions and cations is possibly due to the coulombic bridging in which the closeness of approach of the reactant ions in the activated complex is such as to allow added ions to come in between them, thereby affecting the rates of reaction. This is when the reaction proceeding through the outer-sphere pathway in in which the reactant ions maintain their coordination integrity in the activated complex prior to and during electron transfer" [37].

3.6 Test for Intermediate Complex Formation

The Michaelis-Menten plot of $1/k_{\text{obs}}$ versus $1/[\text{TB}]$ showed no intercept on the $1/k_{\text{obs}}$ axis (Fig. 4),

suggesting the absence of an intermediate complex formation. Spectrophotometric measurement showed no clear shift in the λ_{max} from 530nm when reactants were mixed and the electronic spectrum was recorded after 20 minutes of mixing. This further suggests the absence of intermediate complex formation [13, 15].

3.7 Reaction Mechanism

The mechanism can be explained as follows: protonation of MnO_4^- by H^+ occurred at equation 4 to form HMnO_4 . TB reacted with HMnO_4 to form to form Mn^{2+} and other product that were not determined in equation 5. TB further reacted with MnO_4^- in equation 6, moreover equation 5 and 6 are the rate determining steps. The mechanism is as follows:



Table 2. Rate constants for the effect of cations and anions in the oxidation reaction of theobromine and permanganate ion at $[\text{TB}] = 2.0 \times 10^{-2} \text{ mol dm}^{-3}$, $T = 24.0 \pm 0.1^\circ\text{C}$, dm^{-3} , $\lambda_{\text{max}} = 530 \text{ nm}$, $[\text{MnO}_4^-] = 2.0 \times 10^{-4} \text{ mol dm}^{-3}$ and $I = 1.0 \text{ mol dm}^{-3} (\text{Na}_2\text{SO}_4)$

| Zn | $10^3 [\text{Z}], \text{ mol dm}^{-3}$ | $10^3 k_1, \text{ s}^{-1}$ | $k_2, \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ |
|--------------------|--|----------------------------|---|
| Mg^{2+} , | 0.00 | 2.26 | 010 |
| | 5.00 | 2.68 | 0.13 |
| | 10.0 | 3.07 | 0.15 |
| | 20.0 | 3.83 | 0.19 |
| | 30.0 | 4.20 | 0.21 |
| | 40.0 | 4.79 | 0.25 |
| Ca^{2+} | 0.00 | 2.11 | 0.10 |
| | 5.00 | 3.45 | 0.13 |
| | 10.0 | 3.64 | 0.16 |
| | 20.0 | 3.83 | 0.19 |
| | 30.0 | 4.03 | 0.22 |
| | 40.0 | 4.41 | 0.23 |
| HCOO^- | 0.00 | 2.03 | 0.10 |
| | 5.00 | 3.26 | 0.14 |
| | 10.0 | 3.45 | 0.17 |
| | 20.0 | 3.83 | 0.19 |
| | 30.0 | 4.41 | 0.22 |
| | 40.0 | 4.98 | 0.24 |
| NO_3^- | 0.00 | 2.73 | 0.11 |
| | 5.00 | 3.64 | 0.15 |
| | 10.0 | 4.03 | 0.18 |
| | 20.0 | 4.41 | 0.21 |
| | 30.0 | 4.60 | 0.23 |
| | 40.0 | 5.37 | 0.26 |




Fig. 4. Michealis–Mentens plot of $1/k_{obs}$ versus $1/[TB]$ at $[MnO_4^-] = 2.0 \times 10^{-4} \text{ mol dm}^{-3}$, $[TB] = 4.0 \times 10^{-3} - 2.8 \times 10^{-2} \text{ mol dm}^{-3}$, $I = 1.0 \text{ mol dm}^{-3}$ (Na_2SO_4) and $T = 24 \pm 0.1^\circ C$

$$\text{Rate} = k_1[TB][HMnO_4] + k_2[TB][MnO_4^-] \quad (7)$$

$$\text{But, } [HMnO_4] = K[H^+][MnO_4^-] \quad (8)$$

$$\text{Rate} = Kk_1[TB][MnO_4^-][H^+] + k_2[TB][MnO_4^-] \quad (9)$$

$$\text{Rate} = (Kk_1[H^+] + k_2)[TB][MnO_4^-] \quad (10)$$

$$\text{Rate} = (Kk_1 + [H^+])[TB][MnO_4^-] \quad (11)$$

4. CONCLUSION

The electron transfer reaction of permanganate ion and theobromine in aqueous sulfuric acid media showed a stoichiometry of 1:2. The order of one was observed with respect to both the obromine and permanganate ion. The rate of the reaction was enhanced by added NO_3^- , $HCOO^-$, Mg^{2+} and Ca^{2+} . The Michaelis-Mentens plot of $1/k_{obs}$ versus $1/[TB]$ showed no intercept on the $1/k_{obs}$ axis, suggesting the absence of an intermediate complex formation. From the above reasoning it can be inferred that the kinetic and spectroscopic evidences are in support of the occurrence of outer-sphere mechanism in the theobromine–permanganate reaction.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Marcus RA, Sutin N. Electron transfer in chemistry and biology. Acta. 1985;811:265 - 322.
- Marcus RA. Electron transfer reactions in chemistry. Theory and experiment. Pure Appl. Chem. 1997;63:13-29.
- Greenwood NN, Earnshaw A. Chemistry of the Elements 2nd edition. Oxford Butherworth Heinemann. 1997;182-189.
- Holleman AF, Wiberg E. Inorganic Chemistry. Academic Press San Diego;2001. pp. 67- 69.
- Osunlaja AA. Kinetics and mechanisms of the oxidation of some thioureas by μ -superoxo binuclear cobalt(III) complex and 3,7- bis(dimethylamino)phenazothionium chloride in aqueous acidic media. Unpublished Ph.D dissertations, Ahmadu Bello University, Zaria, Nigeria and the References Therein; 2014.
- Iyun JF. The Kinetics and mechanisms of the oxidation-reduction and substitution reactions of some metal ion complexes in acid solutions. Unpublished Ph.D Thesis, Ahmadu Bello University, Zaria, Nigeria;1982.
- William MM. Dictionary of Biochemistry and related subjects. Philosophical library. 1943; 311,530,573.
- Brij B, Tewari GB, Larsen A. Kengne-Momo, A, Auclair, K and Lan S. Butler. An overview of molecular spectroscopic

- studies on theobromine and related alkaloid. *Appl. Spectrosc. Rev.* 2012; 47(3):163-179.
9. Mumford GK, Benowitz NL, Evans SM, Kaminski BJ, Preston KL, Sannerud CA, et al. Absorption rate of methylxanthines following capsules, cola and chocolate. *Eur. J. Clin. Pharmacol.* 1996;51 (3-4):319-325.
DOI:10.1007/s002280050205. PMID 9010706. S2CID 8405909.
 10. Ashihara H, Yokota T, Crozier A. Biosynthesis and catabolism of purine alkaloids. *New Light on Alkaloid Biosynthesis and future prospects. Adv. Bot. Res.* 2013;68:111-138.
DOI:10.1016/B978-0-12-408061-4.00004-3. ISBN 9780124080614.
 11. Ishaq A. Kinetics and mechanism of oxidation of nicotine by permanganate ion in acid per chlorate solution. *Int. J. Chem.* 2010;2(2):193-200.
 12. Anweting IB, Iyun JF, Idris SO. Kinetics and mechanism of oxidation of nicotinic acid by permanganate ion in aqueous acidic medium. *Arch. Appl. Sci. Res.* 2012a;4(4):1628-1635.
 13. Anweting IB, Iyun JF, Idris SO. Kinetics and mechanism of oxidation of L- cysteic acid by permanganate ion in aqueous acidic medium. *Arch. Appl. Sci. Res.* 2012b;4(5):2002-2011.
 14. Anweting IB, Iyun JF, Idris SO. Kinetics and mechanistic approach to the oxidation of L- tryptophan by permanganate ion in aqueous acidic medium. *Adv. Appl. Sci.* 2012c ;3(6):3401-3409.
 15. Anweting IB, Iyun JF, Idris SO. Electron transfer reaction of L-aspartic acid and permanganate ion in aqueous acidic medium. *Adv. Appl. Sci.* 2012d;3(6):3421-3429.
 16. Ajibade F, Iyun JF. The kinetics and mechanism of oxidation of acetaldehyde by dichromate and permanganate ions in aqueous acidic medium. *J. Chem. Soc. Nig.* 1992;15:80.
 17. Adetoro A, Idris SO, Onu AD, Okibe FG. Kinetics and mechanistic steps to electron transfer reaction of peroxo-bridged binuclear cobalt(II) complex of succinimide by glycine in aqueous acidic medium. *Bull. Chem. Soc. Ethiop.* 2021;35(2):425-434.
 18. Ukoha PO, Ibrahim E. Mechanism of the oxidation of β -mercaptoacetic acid by trioxiodate(V) in aqueous acid medium. *Chemcl. J.* 2004;138-141.
 19. Anweting IB, Idris SO, Onu AD. Kinetics and mechanism of oxidation of thiourea by tetrakis (2, 2' - bipyridine)- μ - oxodiiron (III) complex in aqueous acidic medium. *FUW Trends Sci.Technol. J.* 2017;2(2):850-860.
 20. Anweting IB, Idris SO, Onu AD. Kinetics and mechanism of redox reaction of tetrakis (2, 2'- Bipyridine)- μ - oxodiiron(III) complex and glutathione in aqueous hydrochloric Acid. *FUW Trends Sci. Technol. J.* 2021;1(6):161-165.
 21. Insauti M, Meta-Perez F, Alvarez-Macho MP, Kinetic study of the oxidation of L-phenylalanine by potassium permanganate in acidic medium. *Int. J. Chem. Kinetic.* 1995;27:507.
 22. Mohammed Y, Iyun JF, Idris SO. Kinetic approach to the mechanism of the redox reaction of malachite green and permanganate ion in aqueous acidic medium. *Afr. J. Pure Appl. Chem.* 2009;3(12):269- 274.
 23. Iyun JF, Shehu AR. Kinetics and mechanism of the oxidation of ethanol and propanol by chromium(VI) in acidic medium. *Chemcl. J.* 2004 ;55 -58.
 24. Busari A, Idris SO, Onu AD, Abdulkadir I. Reaction of $S_2O_5^{2-}$ ion and μ -oxo-tetrakis(1, 10-phenanthroline)diiron(III) complex ion in aqueous phenanthroline buffer:a kinetic study. *FUW Trends Sci.Technol. J.* 2019. 4(3) :651-656.
 25. Ukoha PO, Iyun JF. Oxidation of L-ascorbic acid by $enH_2[(FeHEDTA)_2O].6H_2O$ in aqueous medium. *J. Chem. Soc. Nig..* 2002. 27 (2):119 - 122.
 26. Ukoha PO, Ibrahim E. Mechanism of the oxidation of β -mercaptoacetic acid by trioxiodate(V) in aqueous acid medium. *Chemcl. J.* 2004;138-141.
 27. Adetoro A, Iyun JF, Idris SO. Bromate ion oxidation of pyrocatechol violet in aqueous hydrochloric acid. kinetics and mechanism. *Arch. Appl. Sci. Res.* 2010. 2(6):177- 184.
 28. Ukoha PO, Iyun JF. Kinetics of reduction of an Iron(III) complex ion by mercaptoethanol and mercaptoethylamine in perchloric acid medium. *J. Chem. Soc. Nig..* 2001. 26(2):163-168.
 29. Osunlaja AA, Idris SO, Iyun JF. Mechanism of the reduction of methylene blue by thiourea in aqueous acidic medium..*Int. J. Chem. Technol. Res.* 2012; 4(2):609 - 617.

30. Osunlaja AA, Idris SO, Iyun JF, Uzairu A. Reaction of 1,3-dimethyl-2-thiourea with coordinated superoxide in $[(\text{NH}_3)_5\text{CO}(\mu\text{-O}_2)\text{CO}(\text{NH}_3)_5]^{5+}$. A mechanistic study. Int. J. Chem. Stud. 2013;3(1):201-208.
31. Iyun JF. The study of the Kinetic and mechanism of the oxidation of catechol by permanganate and bromate ions in acid medium. J. Chem. Soc. Nig... 1990;15:80-86.
32. Banerji KK. Kinetics and mechanism of oxidation of aliphatic aldehydes by permanganate. Indian J. Chem. 1973; 11:242.
33. Hassan R.M. Kinetics and mechanism of oxidation of DL-Alanine by acid perchlorate ions in media solution. Can. J. Chem. 1991;69:2018.
34. Zahedi M, Bahrami H.. Kinetics and mechanism of the autocatalytic oxidation of L- asparagines in a moderately concentrated sulfuric acid medium. Kinet. Cat. 2004;45 (3):351-358.
35. Babatunde OA, Iyun JF. Kinetics and mechanism of the reaction of di- μ -oxo-tetrakis (1,10-phenanthroline)-dimaganese (III, IV) perchlorate by sulphite ion in acidic solution. J. Sci. Res. 2009;26 (3) :406-416.
36. Babatunde OA. A Study of the Kinetics and mechanism of oxidation of L-ascorbic by permanganate ion in acidic medium. World J. Chem. 2008;3(1):27 – 31.
37. Sharpe AG. Inorganic Chemistry. First edition Longman, London. 1982;518-534.

© 2023 Jones et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/98168>