



# Study of Genetic Diversity in Indian Barnyard Millet Genetic Resources [*Echinochloa frumentacea* (L.)]

G. Nehru <sup>a++\*</sup>, A. Trivikrama Reddy <sup>b#</sup>,  
C. V. Chandra Mohan Reddy <sup>c†</sup> and K. N. Sreenivasulu <sup>d‡</sup>

<sup>a</sup> Department of Genetics and Plant Breeding, Agricultural College, Mahanandi, A.P., India.

<sup>b</sup> Department of Genetics and Plant Breeding, A.P.S.S.C.A, Lam Guntur, A.P., India.

<sup>c</sup> ARS, Rekulakunta, A.P., India.

<sup>d</sup> Department of Statistics & Computer Applications, Agricultural College, Bapatla, A.P., India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2023/v13i102978

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/105870>

Original Research Article

Received: 28/06/2023

Accepted: 05/09/2023

Published: 13/09/2023

## ABSTRACT

Sixtyfour Indian barnyard millet genetic resources were studied to assess the extent of genetic divergence for 18 quantitative traits using Mahalanobis D2 Statistic. Sixtyfour genetic resources of Indian barnyard millet were evaluated during kharif, 2020 in a lattice square design, replicated twice at Regional Agricultural Research. Based on Tocher's method, the entire genetic materials were separated into eight distinct and non-overlapping clusters suggesting availability of genetic diversity. Cluster I has highest number of accessions (57) and the remaining were mono genotypic clusters II, III, IV, V, VI, VII and VIII containing only one accession each indicating high degree of

<sup>++</sup> P.G. Scholar;

<sup>#</sup> Director;

<sup>†</sup> Senior Scientist (Small Millets);

<sup>‡</sup> Assistant Professor;

\*Corresponding author: E-mail: [nehruna2015@gmail.com](mailto:nehruna2015@gmail.com);

heterogeneity among the genotypes. The widest inter-cluster distance was recorded between clusters IV and VII. Therefore, it is concluded that the genotypes to these clusters has to be intercrossed in order to create more diversity and improving grain yield in barnyard millet. The traits, grain yield per plant and panicle weight contributed maximum towards total divergence has scope of crop improvement through those traits. Diversity of parent paly main role for the successful of any breeding programme. So the crosses can be made among the parents having widest divergence would more likely to yield desirable recombination in segregating generations.

**Keywords:**  $D^2$  statistic; genetic diversity; genetic resource; Indian barnyard millet.

## 1. INTRODUCTION

Barnyard millet serves as dual purpose for feed and fodder crop [1]. It is mainly grown in India, China, Japan, and Korea. It is the fourth most produced minor millet, providing food security to many poor people across the world. Globally, India is the biggest producer of barnyard millet, both in terms of area (0.146 m ha<sup>-1</sup>) and production (0.147 mt) with average productivity of 1034 kg/ha during the last 3 years (IIMR, 2018). The major barnyard millet producing states in India are Uttarakhand, Madhya Pradesh, Karnataka, Uttar Pradesh and North east region of India. The cultivated species of barnyard millet viz., Indian barnyard millet (*Echinochloa frumentacea*) and second Japanese barnyard millet (*Echinochloa esculenta*) Sood et al. [2].

Prevalence of genetic diversity is a prime requisite in any crop improvement programme. Multivariate analysis through  $D^2$  statistic, provides a quantitative estimate of association between geographic and genetic diversity based on the generalized distance. It enables to discriminate between different cultivars according to the diversity prevalent in the germplasm collections and thus assist in the selection of genetically divergent parents for their exploitation in hybridization programme. In addition, principal component analysis may be used to classify the relationships between the traits in multi-trait systems and for identifying the patterns of data by reducing the number of dimensions. It also provides an insight into the process contributing differences in yield among the genotypes, a vital aspect in identification and selection of top ranking genotypes out of diverse germplasm base.

Now a days the crop gained importance due to its rich nutritional profile and high dietary fibre content. Watanable [3] stated that the importance of barnyard millet has increased due to its highly rich nutritious grains and presence of strong

antioxidative compounds. The barnyard millet contains 10.5% protein, 3.6% fat, 68.8% carbohydrate and 398 kcal/100 g energy. The total dietary fiber content was high (12.6%) including soluble (4.2%) and insoluble (8.4%) fractions. Indian barnyard millet contains between 51.5% to 59.5% starch Ugare et al. [4]. The genetic divergence studies enable to assess the extent of genetic diversity prevalent in the genetic resources utilized for study. Material and Methods

## 2. MATERIALS AND METHODS

Sixtyfour genetic resources of Indian barnyard millet were evaluated during kharif, 2020 in a lattice square design, replicated twice at Regional Agricultural Research Station, Nandyal. The details of 64 genotypes are given in the Annexure I. Data was collected on five randomly selected plants of each genotypes per replication for 18 quantitative traits like Days to 50% flowering, basal tillers, Days to maturity, Peduncle length (cm), Sood et al. [2] in barnyard millet and Amarnath et al. [5] in foxtail millet. flag leaf blade length(cm), flag leaf blade width(cm), Plant height (cm), Panicle length (cm), 1000 seed weight (gm), Panicle weight /plant (gm), Harvest Index(%), Lower raceme length (cm), Grain yield/plant (gm), Phosphorus, Calcium, Iron, Zinc mg/100g, Protein %. The analysis was done through software - WINDOWSTAT 9.2 version. The experimental design used for the study was lattice square design in two replication with randomization and the statistical analysis method used are ANOVA,  $D^2$  statistics and Principal component analysis or canonical root analysis.

## 3. RESULTS AND DISCUSSION

The ANOVA (Table 1) revealed significant differences among the genetic material for all the traits studied indicating considerable extent of variability in the germplasm accessions, which could be used in further breeding programmes.

**Table 1. Analysis of variance for 18 morpho-physiological and biochemical characters studied in barnyard millet germplasm accessions**

	DF	Days to 50% flowering	basal tillers	Days to maturity	peduncle length (cm)	FLB Length(cm)	FLB Width(cm)
Replications	1	6.498	0.17	6.216	0.028	1.123	0.023
Treatments (unadjusted)	63	106.721 **	0.788 **	106.202 **	12.791 **	12.886 **	0.122 **
Error (r.c.b.)	63	8.680 **	0.057 **	30.618 **	1.217 **	3.146 **	0.018 **
Blocks within Replicated (adj)	14	10.312	0.073	35.729	1.354	3.791	0.022
Among (a)	7	11.495	0.034	39.991	1.771	3.003	0.013
Among (b)	7	9.129	0.112	31.468	0.937	4.579	0.031
Intrablock Error	49	8.213	0.053	29.157	1.178	2.962	0.017
Total	127	57.297 **	0.421 **	67.920 **	6.949 **	7.962 **	0.070 **
	DF	Plant height (cm)	Panicle length (cm)	1000 seed weight (gm)	panicle weight / plant (gm)	Harvest Index (%)	lower raceme length (cm)
Replications	1	5.53	0.025	0.057	0.512	3.073	0.107
Treatments (unadjusted)	63	389.058 **	18.084 **	0.459 **	6.365 **	10.119 **	1.707 **
Error (r.c.b.)	63	51.494 **	1.366 **	0.033 **	0.596 **	4.204 **	0.071 **
Blocks within Replicated (adj)	14	50.592	1.833	0.032	0.765	5.529	0.066
Among (a)	7	37.696	2.954 *	0.033	0.622	7.957	0.043
Among (b)	7	63.487	0.712	0.03	0.908	3.1	0.09
Intrablock Error	49	51.751	1.233	0.034	0.547	3.826	0.073
Total	127	218.585 **	9.649 **	0.245 **	3.457 **	7.129 **	0.883 **
	DF	Grain yield/ plant (gm)	Phosphorus (mg/ 100g)	Calcium (mg/ 100g)	Iron (mg/ 100g)	Zinc (mg/ 100g)	Protein %
Replications	1	3.971	7.681	0.296	4.423	0.972	0.997
Treatments (unadjusted)	63	5.024 **	192.381 *	11.996 **	8.228 **	86.895 **	3.414 **
Error (r.c.b.)	63	1.138 **	106.989 **	1.660 **	1.297 **	10.480 **	0.476 **
Blocks within Replicated (adj)	14	1.45	151.448	3.416 **	1.355	6.091	0.637
Among (a)	7	0.405	52.24	3.616 **	1.621	5.139	1.127 *
Among (b)	7	2.494 *	250.656 *	3.216 *	1.089	7.043	0.146
Intrablock Error	49	1.049	94.287	1.159	1.281	11.734	0.430
Total	127	3.088 **	148.567 *	6.777 **	4.760 **	48.312 **	1.937 **

**Table 2. Clustering pattern of 64 barnyard millet genotypes by Tocher' method**

Cluster No.	No.of genotypes	Genotype(s)
I	57	GEch-8, GEch-45, GEch-20, GEch-60,GEch-94, GEch-186, GEch-191, GEch-360, GEch-365, GEch-372, GEch- 416, GEch-418, GEch-393,GEch-383, GEch-567, GEch-568, GEch-566, GEch- 570,GEch-571, GEch-572, GEch-573, GEch-374, GEch-578, GEch-579, GEch-206, GEch-581, GEch-997, GEch-634, TNEf-190, TNEf-191, TNEf-192, TNEf-193, TNEf-194, TNEf-195, TNEf-196, TNEf-197, TNEf-198, TNEf-199, TNEf-201, TNEf-203, TNEf-204, TNEf-205, TNEf-206, TNEf-207, TNEf-208, TNEf-209, TNEf-210, TNEf-211, TNEf-212, TNEf-213, TNEf-214, TNEf-216, TNEf-217, TNEf-219, BM NDL-1, BM NDL-2,BM NDL-3, TNEf-318.
II	1	GEch-633
III	1	TNEf-200
IV	1	TNEf-216
V	1	TNEf-202
VI	1	TNEf-215
VII	1	TNEf-218
VIII	1	GEch-995

**Table 3. Average intra and inter cluster distance for the barnyard millet accessions studied**

Clusters	Cluster Distances							
	I	II	III	IV	V	VI	VII	VIII
I	92.23	171.93	174.5	152.98	140.92	244.29	213.76	148.2
II		0	248.36	286.75	168.04	268.23	127.71	163.12
III			0	80.2	217.92	107.77	277.34	158.92
IV				0	287.28	62.03	360.53	168.53
V					0	328.08	126.78	113.43
VI						0	401.29	152.7
VII							0	207.48
VIII								0

**Table 4. Cluster means with respect to yield and yield attributes among 64 barnyard millet germplasm accession**

Clusters	Cluster Means: Torcher method																	
	DFF	BT	DTM	PL (cm)	FLB L (cm)	FLBW (cm)	PH (cm)	PAL (cm)	TSW (g)	PW (g)	HI (%)	LRL (cm)	GYP (gm)	Ph (mg/ 100g)	Ca (mg/ 100g)	Fe (mg/ 100g)	Zn (mg /100g)	Pr (%)
I	48.17	1.94	88.24	17.56	26.92	2.2	118.25	18.46	2.48	10.1	33.66	3.23	8.69	276.13	19.69	16.72	45.12	8.75
II	42.5	2.7	82.5	20.23	24.35	2	118.45	25.72	2.95	11.2	31.43	6.02	9	284	18.05	18.3	54.1	8.53
III	69	1.4	109	23.25	29.17	2.38	121.66	18.3	2.7	14.4	34.25	2.7	12.15	279.5	16.1	16.7	55.55	7.53
IV	68.99	2	108.98	18.9	26.49	2.48	116.05	16.37	2.75	8.1	32.22	2.27	7.25	266	17.9	17.35	38.23	7.5
V	43.5	2.3	84	14.74	24.65	1.88	98.33	14.02	2.45	14.7	34.64	3.4	12.9	266	17.05	16.35	56.55	7.53
VI	72.5	2.5	112.5	18.95	27	2.53	122.17	17.29	2.3	10.9	34.89	3.82	8.75	267	18.35	16.95	41.95	6.83
VII	41	2.3	81	23.97	21.95	2.21	90.91	15.05	3.65	14.9	30.23	5	12.85	280	22.7	15.1	43.6	7
VIII	55.5	3.9	95.5	19.5	30	1.79	121.5	18.38	2.2	12.6	31.55	4.25	10.4	281.5	21.55	16.5	52.25	7.68

DFF Days to 50% Flowering, BT Basal tillers, DTM Days to maturity, PL Peduncle length, FLBL Flag leaf blade length, FLBW Flag leaf blade width, PH Plant height, PAL Panicle length, TSW 1000 seed weight, PW Panicle weight, HI Harvest index, LRL Lower raceme length, GYP Grain yield per plant, Ph Phosphorus mg/100g, Ca Calcium mg/100g, Fe Iron mg/100g, Zn mg/100g, Pr Protein.

**Table 5. Contribution of yield and its attributes towards total diversity in 64 barnyard millet germplasm accessions**

S. No.	Characters	Contribution %	Times ranked 1st
1	Days to 50% flowering	0.4	7.1
2	Basal tillers	7.4	149.0
3	Days to maturity	2.3	46.4
4	Peduncle length (cm)	4.8	97.0
5	Flag Leaf Blade Length (cm)	0.2	4.0
6	Flag Leaf Blade Width (cm)	4.2	84.1
7	Plant height (cm)	1.8	36.1
8	Panicle length (cm)	0.2	3.0
9	1000 seed weight (gm)	6.3	127.0
10	Panicle weight /plant (gm)	15.7	317.0
11	Harvest Index (%)	13.5	272.0
12	lower raceme length (cm)	7.3	147.0
13	Grain yield/plant (g)	16.6	335.1
14	Phosphorus (mg/100g)	0.2	3.8
15	Calcium (mg/100g)	3.7	74.6
16	Iron (mg/100g)	2.3	46.4
17	Zinc (mg/100g)	6.1	122.0
18	Protein (%)	7.2	144.8

**Table 6. Canonical vectors for 18 characters in 64 barnyard millet germplasm accessions**

Parameters	Z <sub>1</sub>	Z <sub>2</sub>	Z <sub>3</sub>	Z <sub>4</sub>	Z <sub>5</sub>	Z <sub>6</sub>	Z <sub>7</sub>
Eigene Value (Root)	3.70473	2.73658	1.8845	1.75198	1.47025	1.16972	0.9
% Var. Exp.	20.58185	15.20319	10.46943	9.73324	8.16807	6.49842	5.2
Cum. Var. Exp.	20.58185	35.78505	46.25448	55.98772	64.15578	70.65421	75.9
<b>Characters</b>							
Days to 50% flowering	0.40553	0.14318	0.34532	0.02626	0.01383	0.17468	0
basal tillers	-0.18463	0.20613	-0.0076	-0.31914	-0.0538	0.55239	0
Days to maturity	-0.39136	-0.17318	-0.33946	0.00617	-0.00573	-0.23601	0
peduncle length (cm)	-0.013	0.27683	0.13999	-0.08948	0.63295	-0.09399	0
FLB Length(cm)	0.11125	-0.14059	-0.2546	-0.17546	0.30118	0.42982	0.2
FLB Width(cm)	0.23477	0.01766	0.04979	0.4873	-0.21787	0.15969	0
Plant height (cm)	0.31174	0.03393	-0.26736	-0.19041	-0.10658	-0.18571	0.3
Panicle length (cm)	0.21917	0.22722	-0.32112	0.23562	-0.02406	-0.13507	0.3
1000 seed weight (g)	-0.28102	-0.02969	-0.1274	0.39129	0.22562	-0.16856	0.1
panicle weight /plant (g)	-0.21117	0.22571	0.18775	-0.06614	-0.28809	0.05128	0.3
Harvest Index(%)	0.39348	-0.11278	0.08662	-0.00875	0.17041	-0.2252	0.1
lower raceme length (cm)	-0.08338	0.46686	-0.24556	0.01618	0.10394	-0.06371	-0.1
Grain yield/plant (g)	-0.1167	-0.09104	0.03925	0.55332	0.174	0.39394	0
Phosphorus mg/100g	-0.1728	0.18516	0.43724	-0.00856	0.32679	-0.18849	0.1
Calcium mgs/100g	-0.03164	-0.46873	-0.0194	-0.23863	0.1569	0.01741	0.1
Iron mg/100g	0.04763	0.22991	-0.39911	0.00839	0.18985	0.21672	0
Zinc mg/100g	-0.25998	-0.19603	0.17271	0.05257	-0.04862	0.08949	0.5
Protein %	0.18902	-0.3485	-0.0497	0.07461	0.2689	0.07093	-0.1

**Table 7. Canonical root values, per cent of variation and cumulative variation explained for 64 barnyard millet germplasm accessions**

<b>Eigene Value (Root)</b>	3.70473	2.73658	1.8845	1.75198	1.47025	1.16972	0.9
<b>% Var. Exp.</b>	20.58185	15.20319	10.46943	9.73324	8.16807	6.49842	5.2
<b>Cum. Var. Exp.</b>	20.58185	35.78505	46.25448	55.98772	64.15578	70.65421	75.9

**Table 8. Mean values of canonical vectors for 64 barnyard millet germplasm accessions**

S. No	Genotype	X Vector	Y Vector	Z Vector
1	Gech-8	3.498	8.382	4.284
2	Gech-45	7.768	4.824	4.366
3	Gech-20	12.916	7.279	3.867
4	Gech-60	6.474	1.984	4.394
5	Gech-94	6.922	7.644	3.136
6	Gech-186	8.306	8.543	4.484
7	Gech-191	9.652	3.384	4.645
8	Gech-360	7.757	5.525	5.016
9	Gech-365	6.646	3.655	6.252
10	Gech-372	3.734	1.387	7.34
11	Gech-416	8.502	2.087	4.779
12	Gech-418	9.684	0.717	3.385
13	Gech-393	9.59	2.765	6.393
14	Gech-383	9.496	4.531	7.518
15	Gech-567	8.537	11.057	3.342
16	Gech-568	9.488	7.851	4.452
17	Gech-566	6.139	8.131	4.711
18	Gech-570	7.292	6.243	5.328
19	Gech-571	4.657	8.791	3.633
20	Gech-572	7.128	2.323	7.98
21	Gech-573	6.775	7.246	5.782
22	Gech-374	7.748	0.971	8.766
23	Gech-578	5.283	5.674	5.295
24	Gech-579	5.856	2.438	7.223
25	Gech-206	10.81	3.714	7.87
26	Gech-581	4.136	5.899	7.341
27	Gech-997	8.876	3.722	9.682
28	Gech-995	6.865	9.463	10.14
29	Gech-633	4.368	13.015	2.946
30	Gech-634	9.227	3.522	6.348
31	TNEf-190	13	7.703	4.216
32	TNEf-191	7.955	0.761	9.631
33	TNEf-192	8.372	0.477	8.543
34	TNEf-193	11.612	5.864	8.537
35	TNEf-194	10.029	8.875	3.463
36	TNEf-195	10.58	1.12	8.753
37	TNEf-196	7.056	4.011	9.506
38	TNEf-197	4.506	0.824	8.462
39	TNEf-198	7.218	2.899	7.302
40	TNEf-199	12.955	9.244	13.684
41	TNEf-200	5.465	3.245	7.604

S. No	Genotype	X Vector	Y Vector	Z Vector
42	TNEf-201	1.957	5.771	9.192
43	TNEf-202	9.821	1.178	6.593
44	TNEf-203	8.739	-1.358	6.474
45	TNEf-204	6.269	-0.664	8.218
46	TNEf-205	13.451	3.924	8.746
47	TNEf-206	5.075	2.972	7.818
48	TNEf-207	8.236	1.036	5.426
49	TNEf-208	12.432	8.324	10.1
50	TNEf-209	13.878	4.078	6.856
51	TNEf-210	6.454	3.584	6.496
52	TNEf-211	14.943	7.961	7.961
53	TNEf-212	10.781	3.091	9.663
54	TNEf-213	11.931	3.239	9.512
55	TNEf-214	16.194	12.923	11.406
56	TNEf-215	15.74	6.724	10.987
57	TNEf-216	4.806	4.074	6.465
58	TNEf-217	-1.178	9.54	9.022
59	TNEf-218	8.693	5.406	8.2
60	TNEf-219	7.336	2.049	5.76
61	BMNDL-1	7.827	4.863	7.876
62	BMNDL-2	7.838	3.764	5.07
63	BMNDL-3	12.781	3.813	7.165
64	TNEf-318	3.498	8.382	4.284







Fig. 1. Clustering pattern of 64 barnyard millet germplasm accessions by Torchers method



**Fig. 2. Cluster diagram showing average intra and inter cluster distances**

Based on relative magnitude of  $D^2$  values, 64 genotypes were grouped into eight distinct and non-overlapping clusters (Table 2, Fig. 1). Cluster I has highest number of accessions (57) and the others were mono genotypic clusters II, III, IV, V, VI, VII and VIII containing only one accession each indicating high degree of heterogeneity among the genotypes. The  $D^2$  values of eight clusters are presented in the Table 3 and Fig. 2. The intra-cluster distances were lower than the inter-cluster distances indicating the presence of widest genetic diversity between the clusters rather than within the clusters. The intra cluster distance (0) observed in clusters II, III, IV, V, VI, VII and VIII indicated the close resemblance between the germplasm accessions present in

that cluster. Therefore, selection of parents which occupying in the same cluster may not be considered effective as it lacks diversity (Kumar et al. (2010). Inter-cluster distance values varied from 62.03 (cluster IV and cluster VI) to 401.29 (cluster VI and cluster VII). Hence, the germplasm accessions between cluster IV (TNEf-216, TNEf-202, TNEf-215) and cluster VII TNEf-218 has high degree of genetic diversity as they possessing maximum inter cluster distance between cluster VI and VII. This genotypes can be utilized under for hybridization programme for obtaining superior segregants. Similar finding were reported by Brunda et al. [6], Devaliya et al. [7], Mahanthesha et al. [8] and Amarnath et al. [5].






Fig. 3. Relative contribution of 18 characters to total genetic diversity in 64 barnyard millet germplasm accession



**Fig. 4. Two dimensional (2D) graph showing relative positions of 64 barnyard millet germplasm based on PCA scores**

Cluster means for 18 traits per cluster were presented in Table 4. The clusters formed were distinct as maximum variation was observed in the clusters for the characters studied. The cluster I, observed maximum for protein content (8.7). Cluster II has maximum values for panicle length (25.7), lower raceme length (6), phosphorus content (284 mg) and iron content (18.3 mg). Cluster III has maximum panicle length (29.1). The mean for grain yield per plant (12.9) and zinc content (56.5) recorded highest in cluster V. The highest mean for days to 50%

flowering (72.5), flag leaf blade width (2.5), plant height (122.1) and harvest index (34.8) are recorded in cluster VI. Cluster VII reported maximum values for peduncle length (23.9), 1000 seed weight (3.65), panicle weight (14.9), calcium content (22.7) and desired means for days to 50% flowering (41), and plant height (90.9) and days to maturity (81.0). Basal tillers (3.9) recorded highest mean in cluster VIII. The least values for days to 50% flowering (41), days to maturity (81) i.e earliness were reported in Cluster VII.



**Fig. 5. Three dimensional (3D) graph showing relative positions of 64 barnyard millet germplasm based on PCA score**

The result shows that no cluster has at least one genotype with all desirable characters, so the direct selection of genotype is not possible for immediate use. Therefore, the genotypes from divergent cluster has to be selected for hybridization programme for combining all characters. Similar reports were recorded by Murugan & Nirmalakumari [9] and Kavya.

The relative contribution of various plant traits towards diversity are presented in the (Table 5 and Fig. 3). All the characters studied, out them the trait grain yield per plant contributed maximum (16.6%) towards genetic diversity and next by panicle weight (15.7%) indicating that these traits can be used for the genetic improvement programme. Similar finding were published by Kumari and Singh (2015), Prabu et al. [10], Manimekalai et al. [11] and Mahanthesha et al. [8] for grain yield / plant.

The canonical root analysis or principal component analysis (PCA) was used to verify the clustering pattern obtained by Mahalanobis  $D^2$  statistic. For 64 barnyard millet germplasm accessions, canonical root analysis was carried out as per the procedure outlined by Rao [12]. Seven canonical roots accounted for 75.9 per cent of total divergence (Table 6). The first,

second, third and fourth roots accounted for 20.58, 15.20, 10.46, and 9.7 per cent respectively to total variability. The remaining canonical roots viz., fifth, sixth and seventh contributed 8.16, 6.49 and 5.2 per cent respectively towards the total variability. The mean values of canonical variates for three roots X, Y and Z were furnished in Table 7. Two dimensional (2D) and three dimensional (3D) picture was constructed by plotting the mean values of vectors as in Fig. 4. and Fig. 5. The amount of contribution of different traits towards canonical vectors total divergence was presented in Table 8.

In the vector Z1, traits contributing towards total divergence positively were Days to 50% flowering (0.405), flag leaf blade length (0.111), flag leaf blade width (0.234), plant height (0.311), panicle length (0.219), harvest index (0.393) and protein (0.189) while the rest of the characters contributed negatively to the total diversity.

For the vector Z2, days to 50% flowering (0.143), basal tillers (0.206), peduncle length (0.276), flag leaf blade width (0.01766), plant height (0.0339), panicle length (0.227), panicle weight per plant (0.225), lower raceme length (0.466), phosphorus (0.185) and iron (0.229) contributed

positively to the genetic diversity. While the traits viz., days to maturity, flag leaf blade length, 1000 grain weight, harvest index, grain yield / plant calcium, zinc and protein had contributed negatively to the diversity.

In the vector Z3, the traits viz., days to 50% flowering (0.345), peduncle length (0.139), flag leaf blade width (0.049), panicle weight per plant (0.187), harvest index (0.086), grain yield per plant (0.039), phosphorus (0.437) and zinc (0.172) had contributed positively to diversity. While rest of the characters showed negative contribution towards divergence.

In the vector Z4, the traits viz., days to 50% flowering (0.026), days to maturity (0.006), flag leaf blade width (0.487), panicle length (0.236), 1000 seed weight (0.391), lower raceme length (0.016), grain yield per plant (0.533), iron (0.008), zinc (0.053) and protein (0.075) had contributed positively to diversity. While rest of the characters showed negative contribution towards divergence.

For the vector Z5, the traits contributed towards total divergence were days to 50% flowering (0.014), peduncle length (0.633), flag leaf blade length (0.301), 1000 seed weight (0.226), harvest index (0.170), lower raceme length (0.104), grain yield per plant (0.174), phosphorus (0.327), calcium (0.157), iron (0.190) and protein (0.269) had contributed positively to diversity. While rest of the characters showed negative contribution towards divergence.

In the vector Z6, the trait days to 50% flowering (0.175), basal tillers (0.552), flag leaf blade length (0.430), flag leaf width (0.160), panicle weight per plant (0.051), grain yield per plant (0.394), calcium (0.017), iron (0.217), zinc (0.089) and protein (0.071) contributed positively to diversity. While rest of the characters showed negative contribution towards divergence.

For the vector Z7, the traits flag leaf blade length (0.200), plant height (0.300), panicle length (0.300), 1000 seed weight (0.100), panicle weight per plant (0.300), harvest index (0.100), phosphorus (0.100) and zinc (0.500) contributed positively to diversity. While rest of the characters showed negative contribution towards divergence.

For the vector Z1 to Z7 similar findings were reported by Prabu et al. [10], Metha et al. [13] in

barnyard millet and Gopikrishana et al. [14] in little millet [15,16].

#### 4. CONCLUSION

Diversity of parent paly main role for the successful of any breeding programme. So the crosses can be made among the parents having widest divergence would more likely to yield desirable recombination in segregating generations. Information on the relative contribution of various plant characters towards divergence (Table 5 and Fig. 3) has also been reported to aid the breeder in choice of parents for hybridization and effective selections in the advance generations. Among all the characters studied, contributed maximum grain yield per plant (16.6%) towards genetic divergence followed by panicle weight per plant (15.7%) indicating feasibility of genetic improvement through those characters. Similar reports were published by Mahanthesha et al. [8] for grain yield per plant.

#### ACKNOWLEDGEMENT

The authors are highly thankful to Acharya N.G. Ranga Agricultural University, Guntur, Andhra Pradesh, India for providing barnyard millet genetic resources and assistance for the conduct of experiment.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Bandyopadhyay BB. Evaluation of barnyard millet cultivars for fodder yield under single and double cut treatments at higher elevation of hills. *Agricultural Science Digest*. 2009;29(1): 66-68.
2. Sood S, Khulbe RK, Kumar A, Agrawal PK, Upadhyaya HD. Barnyard millet global core collection evaluation in the submontane Himalayan region of India using multivariate analysis. *The Crop Journal*. 2015;3(6):517-525.
3. Watanabe M. Antioxidative phenolic compounds from Japanese barnyard millet (*Echinochloa utilis*) grains. *Journal of Agricultural and Food Chemistry*. 1999; 47(11):4500-4505

4. Ugare R, Chimmad B, Naik R, Bharati P, Itagi S. Glycemic Index & Significance of Barnyard Millet (*Echinochloa frumentacea*) in type II Diabetics. Journal of Food Science and Technology. 2014;51(2).
5. Amarnath K, Prasad AD, Reddy CCM. Assessment of genetic diversity in Indian Italian millet genetic resources [*Setaria Italica* (L.) Beauv]. Electronic Journal of Plant Breeding. 2019;10(1):83-91.
6. Brunda SM, Kamatar MY, Naveenkumar KL, Hundekar R. Genetic diversity in the foxtail millet (*Setaria italica*) germplasm as determined by phenotypical traits. The Ecscan. 2015b;8:483-488.
7. Devaliya SD, Singh M, Intawala CG. Genetic divergence studies in finger millet (*Eleusine coracana* (L.) Gaertn.). International Journal of Current Microbiology and Applied Sciences. 2017; 6(11):2017-2022
8. Mahanthesha M, Sujatha M, Pandravada SR, Meena AK. Study of genetic divergence in finger millet (*Eleusine coracana* (L.) Gaertn) germplasm. International Journal of Pure and Applied Bioscience. 2017b; 5(3):373-377.
9. Murugan R, Nirmala kumari A. Genetic divergence in italian millet. (*Setaria italica* (L.) Beauv). Indian Journal of Genetics. 2006;66(4):339-340.
10. Prabu R, Vanniarajan C, Vetrivanthan M, Gnanamalar RP, Shanmugasundaram R, Ramalingam J. Association studies in barnyard millet (*Echinochloa frumentacea* (Roxb.) Link) for early maturity and yield contributing traits at high altitude region. Electronic Journal of Plant Breeding. 2020;11(1):192-196 (2020).
11. Manimekalaia M, Dhasarathanb M, Karthikeyanb MA, Murukarthickc A, Renganathana VG, Thangaraja K, Vellaikumarb S, Vanniarajana C, Senthilb N. Genetic diversity in the barnyard millet (*Echinochloa frumentacea*) germplasms revealed by morphological traits and simple sequence repeat markers. Current Plant Biology. 2018;14:71–78.
12. Rao CR. Advanced Statistical Methods in Biometrical Research. Ed. I, John Wiley and Sons. NelV York; 1952.
13. Mehta H, Tyagi PC, Mohapatra KP. Genetic diversity in barnyard millet (*Echinochloa frumentacea* Roxb.). Indian Journal of Genetics and Plant Breeding. 2005;65 (04):293-295.
14. Gopikrishnan A, Pandiyan M, Thilagam P, Veeramani P, Nanthakumar S. Characterization of little millet (*Panicum sumatrense*) genetic diversity in yelagiri hills of Tamil Nadu. Indian Journal of Plant Genetic Resources. 2021;34(01):59-63.
15. ICAR- Indian Institute of Millets Research. Annual Report: 2017-18. Rajendranagar, Hyderabad. 2018;11-13.
16. Kavya P, Sujatha M, Pandravada SR, Hymavathi TV. Exploiting genetic divergence in italian millet [*Setaria italica* (L.) P. Beauv.] elite germplasm lines: A neglected and a underutilised crop. Environment and Ecology. 2016;35(3C): 2312-2315.

## ANNEXURE

### Annexure I. Details of the 64 Indian barnyard millet genetic resources utilized for study

S. No.	Germplasm accession	Parentage/ Collection/Source
1	GEch-8	Dholi
2	GEch-45	Dholi
3	GEch-20	Dholi
4	GEch-60	NBPGR DS 67/82/22-2-83 (From GE ch 751)
5	GEch-94	Dholi
6	GEch-186	Almora
7	GEch-191	Almora
8	GEch-360	Almora
9	GEch-365	Almora
10	GEch-372	Almora
11	GEch-416	Almora
12	GEch-418	NBPGR Bihar ( from GE ch 759)
13	GEch-393	Almora
14	GEch-383	Almora
15	GEch-567	Bihar
16	GEch-568	Bihar
17	GEch-566	Orissa
18	GEch-570	Orissa
19	GEch-571	Orissa
20	GEch-572	Orissa
21	GEch-573	Orissa
22	GEch-374	Almora
23	GEch-578	Rewa
24	GEch-579	Rewa
25	GEch-206	Almora
26	GEch-581	Rewa
27	GEch-997	India
28	GEch-995	India
29	GEch-633	Madhya Pradesh, Rewa
30	GEch-634	Madhya Pradesh, Rewa
31	TNEf-190	TNAU, Athiyandal
32	TNEf-191	TNAU, Athiyandal
33	TNEf-192	TNAU, Athiyandal
34	TNEf-193	TNAU, Athiyandal
35	TNEf-194	TNAU, Athiyandal
36	TNEf-195	TNAU, Athiyandal
37	TNEf-196	TNAU, Athiyandal
38	TNEf-197	TNAU, Athiyandal
39	TNEf-198	TNAU, Athiyandal
40	TNEf-199	TNAU, Athiyandal
41	TNEf-200	TNAU, Athiyandal
42	TNEf-201	TNAU, Athiyandal
43	TNEf-202	TNAU, Athiyandal
44	TNEf-203	TNAU, Athiyandal
45	TNEf-204	TNAU, Athiyandal
46	TNEf-205	TNAU, Athiyandal
47	TNEf-206	TNAU, Athiyandal
48	TNEf-207	TNAU, Athiyandal
49	TNEf-208	TNAU, Athiyandal
50	TNEf-209	TNAU, Athiyandal
51	TNEf-210	TNAU, Athiyandal
52	TNEf-211	TNAU, Athiyandal



<b>S. No.</b>	<b>Germplasm accession</b>	<b>Parentage/ Collection/Source</b>
53	TNEf-212	TNAU, Athiyandal
54	TNEf-213	TNAU, Athiyandal
55	TNEf-214	TNAU, Athiyandal
56	TNEf-215	TNAU, Athiyandal
57	TNEf-216	TNAU, Athiyandal
58	TNEf-217	TNAU, Athiyandal
59	TNEf-218	TNAU, Athiyandal
60	TNEf-219	TNAU, Athiyandal
61	BMNDL-1	Nandyal local
62	BMNDL-2	PLS from local
63	BMNDL-3	PLS from local
64	TNEf-318	TNAU, Athiyandal

© 2023 Nehru et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/105870>