



# Seroprevalence and Evaluation of IgM and NS1 Ag Detection for Early Diagnosis of Dengue Cases in a Tertiary Care Hospital

Charu Jain <sup>a++</sup>, Nikita Birhman <sup>b#</sup>, Swati Sinha <sup>b†</sup>,  
Anshdha Shah <sup>b†</sup>, Shukla Das <sup>b‡\*</sup>, N. P. Singh <sup>b^</sup>,  
Poonam Mathur <sup>a#</sup>, Narendra Singh Mogha <sup>a##</sup>  
and Divya Kashyap <sup>b#^</sup>

<sup>a</sup> Department of Microbiology, University College of Medical Sciences and GTB Hospital, Delhi, India.

<sup>b</sup> Department of Microbiology, Viral Research and Diagnostic Laboratory, University College of Medical Sciences and GTB Hospital, Delhi, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/JAMB/2023/v23i10753

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/105806>

**Original Research Article**

**Received: 03/07/2023**

**Accepted: 05/09/2023**

**Published: 14/09/2023**

<sup>++</sup> Assistant Professor;

<sup>#</sup> Research Assistant;

<sup>†</sup> Scientist-B;

<sup>‡</sup> Director Professor;

<sup>^</sup> Head & Director Professor

<sup>##</sup> Senior Technical Assistant;

<sup>#^</sup> Laboratory Technician;

\*Corresponding author: E-mail: [cjain@ucms.ac.in](mailto:cjain@ucms.ac.in);

## ABSTRACT

**Introduction:** Dengue fever, a mosquito borne illness, have grown to pose a serious danger to public health systems in developing nations. The Dengue virus (DENV), a member of the Flavivirus genus has four serotypes that cause the disease which range in severity from subclinical infections to severe disease with high morbidity & mortality. This study aims to assess the seroprevalence of dengue fever cases in a tertiary care hospital from East- Delhi region.

**Material & Methods:** This study was a retrospective observational study conducted in the Department of Microbiology at UCMS & GTBH. The data was collected from laboratory records from a period of Jan 2019 to December 2019.

**Results:** A total of 1449 clinically suspected cases of dengue were tested for the presence of NS1 antigen and IgM antibodies by capture ELISA. This study showed 32.22% (467/1449) seropositivity of dengue infection. Out of 1272, a total of 456 samples were positive for NS1 Ag ELISA. Whereas, out of 177, a total 11 samples were positive for IgM ELISA.

**Observation:** The study highlights the importance of early diagnosis and prompt management of dengue cases to prevent severe disease and reduce morbidity and mortality. This underscores the need for increased awareness, early detection, and appropriate management of dengue fever in endemic areas. especially during the peak months of October and November.

**Keywords:** *Aedes mosquito; dengue virus; enteric fever; seroprevalence; IgM.*

## 1. INTRODUCTION

Dengue fever, a mosquito borne illness, have grown to pose a serious threat to public health systems in developing nations. The vector borne viral disease is primarily transmitted by two most common *Aedes* mosquito (*Aedes aegypti* and *Aedes albopictus*) [1]. The Dengue virus (DENV), a member of the *Flavivirus* genus has four serotypes that cause the disease which range in severity from subclinical infections to severe disease with high morbidity & mortality. Despite the fact that the most of dengue fever episodes are mild and self-limiting, it can nevertheless result in deadly complications such as DHF (Dengue Hemorrhagic Fever) and DSS (Dengue Shock Syndrome) [2]. The asymptomatic cases in the earlier stages of infection play a major role in the maintaining transmission of the disease in the community resulting in outbreak [3]. Typical symptoms of dengue include a high-grade fever, headache, discomfort in the muscles and joints, aches, retroorbital pain, and skin rashes [3]. Severe dengue symptoms result from plasma leakage, hemo-concentration leading to hemorrhagic shock, and multiple organ failure with high fatality. The National Centre for Vector Borne Diseases Control (NCVBDC) has provided case definition for the disease based on symptoms & laboratory finding and is a notifiable disease under Integrated Disease Surveillance Project (IDSP) disease survey [4].

Serology, Virus isolation and molecular methods like Nucleic Acid Amplification Test (NAAT) are

some of the diagnostic modalities currently available to detect dengue infection. NAATs and viral culture are expensive & specialized infrastructure is needed therefore have limited diagnostic use. Advance molecular techniques such as RT-PCR, Multiplex RT-PCR, Realtime RT-PCR, and LAMP are available to detect the virus during acute phase of infection [4]. However, for early diagnosis and intervention in resource limited settings, cost effective tests like ELISA are routinely used to identify dengue infection [5].

As per NVBDCP, India reported a total case 157315, 44585 and 193245 during the year of 2019, 2020 and 2021, respectively. The total death reported during the period was 166, 56 and 346, respectively. Delhi reported a total case 5077, 1269 and 13089 during the year of 2019, 2020 and 2021, respectively and only 23 deaths in 2021 [6]. The present study aim to assess the seroprevalence of dengue in the patients treated at a tertiary care hospital from East- Delhi region during 2019.

## 2. MATERIALS AND METHODS

A retrospective observational study was conducted in the Department of Microbiology at UCMS & GTB Hospital. The demographic and laboratory data were retrieved from the department for year 2019. Serological examination based on the days of fever (either NS1 or IgM) was routinely performed in the serum samples which were collected from the

dengue suspected patients. Serology test were performed for NS1Ag and IgM antibodies by using dengue NS1Ag MICROLISA (IR031096) and MACELISA kit (Panbio Dengue IgM Capture ELISA) respectively. Demographic data were entered in MS excel (version 2016). Statistical analysis was performed using SPSS software.

### 3. RESULTS

A total of 1449 clinically suspected dengue cases were tested for the presence of NS1 antigen and IgM antibodies by capture ELISA in the year 2019. This study showed 32.22% (467/1449) seropositivity of dengue infection. Out of 1272, NS1Ag ELISA 456 samples were positive. Whereas, out of 177 sample total 11 samples were positive for IgM ELISA.

Table 1 showed that from the total positive cases, Males (51.82%, 242/467) were more affected than females (48.17%, 225/467). Most common affected age group was 21-40 years (43.89%, 205/467), followed by 6-15 years (25.26%, 118/467) of age group (Table 2). Majority of positive dengue patients were from IPD (93.79%, 438/467) than OPD (6.20%, 29/467) (Table 3). Table 4 depicts the monthly

distribution of positive cases. Maximum cases were recorded in October (30.40%, 142/467) followed by November (27.19%, 127/467).

### 4. DISCUSSION

Dengue NS1 antigen, a highly conserved glycoprotein which is produced in both membrane-associated and secretion forms, is abundant in the serum of patients during the early stages of DENV infection [5]. It's been discovered to be helpful in recognizing the presence of acute dengue infections. It is a simple test that is specific and shows high sensitivity. Early case detection with NS1, or when the infection is in the viremic stage is important from an epidemiological perspective to prevent transmission of the infection. For DENV, the NS1 ELISA-based antigen assay is commercially available, and the sensitivity and specificity of this assay has been assayed previously. Moreover, the specificity of NS1 assay helps in differential diagnosis of various flaviviruses [5]. Considering the cost effectiveness of this test and ease of performance as compared to molecular diagnostic test, the Mac ELISA gives a practical advantage in resource limited settings.

**Table 1. Gender-wise distribution of dengue cases**

GENDER	NS 1 Ag ELISA		IgM Ab ELISA	
	Total number	Positive	Total number	Positive
Male	499	236 (51.75%)	62	6 (54.55%)
Female	773	220 (48.25%)	115	5 (45.45%)
<b>Total</b>	<b>1272</b>	<b>456</b>	<b>177</b>	<b>11</b>

**Table 2. Age-wise distribution of dengue cases**

AGE	NS 1 Ag ELISA		IgM Ab ELISA	
	Total Number	Positive	Total Number	Positive
0-5	53	17 (3.75%)	12	01 (9.09%)
6-15	236	115 (25.21%)	23	03 (27.27%)
16-20	169	66 (14.47%)	24	01 (9.09%)
21-40	658	201 (44.07%)	93	04 (36.36%)
>41	156	57 (12.5%)	21	02 (18.19%)
<b>Total</b>	<b>1272</b>	<b>456</b>	<b>177</b>	<b>11</b>

**Table 3. Incidence of dengue cases from the IPD& OPD wards**

Department	NS 1 Ag ELISA		IgM Ab ELISA	
	Total number	Positive	Total number	Positive
IPD	1175	428	170	10
OPD	97	28	07	01
<b>Total</b>	<b>1272</b>	<b>456</b>	<b>177</b>	<b>11</b>

**Table 4. Month-wise distribution of dengue infection**

MONTHS	IgM Ab ELISA		NS1 Ag ELISA	
	Total Samples	Positive Rate	Total Samples	Positive Rate
Jan	19	06 (31.57%)	132	23 (1.74%)
Feb	00	00	34	02 (5.88%)
Mar	17	00	01	00
Apr	36	00	00	00
May	40	00	00	00
Jun	11	00	44	01 (2.27%)
July	00	00	105	25 (23.8%)
Aug	00	00	140	26 (18.57%)
Sep	00	00	251	105 (41.83%)
Oct	00	00	293	142 (48.46%)
Nov	00	00	253	127 (50.19%)
Dec	54	05 (9.25%)	19	05 (26.31%)
<b>Total</b>	<b>177</b>	<b>11 (6.21%)</b>	<b>1272</b>	<b>456 (35.84%)</b>

IgM-Capture (MAC-ELISA) detect the anti-dengue IgM antibody develops within 5 days of fever though it varies between patients. Detectable IgM may persist in some primary infections for up to 90 days, but in the majority of patients, it declines to undetectable levels by 60 days [5]. Nevertheless, it verifies the presence of infection in the person at some point in the recent past months. MAC-ELISA is now widely used as a potential tool for DF/DHF surveillance. Hospitalized patients, who are frequently admitted at the end of their illness and have detectable IgM levels in their blood, can benefit the most from it [5].

The study shows a high seropositivity rate (32.22%) for dengue infection among clinically suspected cases during 2019. This result was higher compared to the study done by Kumar et al, which shows the seropositivity of dengue infection in 23% cases [7]. This highlights the higher burden and need for accurate and timely diagnosis of dengue fever, especially in endemic areas.

In the present study, the male to female ratio of dengue cases was 1.07:1 (242/225) with a male (51.82%) predominance than female (48.17%). However, the difference between the two groups was not statistically significant and this result is similar to the Indian studies which also showed no difference in the gender distribution of dengue cases [8,9].

The most commonly affected age group was 21-40 years, followed by 6-15 years. Indirectly the need for targeted interventions to prevent and manage dengue fever in these high-risk groups. A study done by Badoni G et al, found that the

most positive cases for DENV are in the 21–30-year age group [10].

The predominance of IPD dengue cases (93.79%) compared to OPD cases (6.20%) raises concerns about the severity of the disease. It suggests that a significant number of patients require specialized care and monitoring. Similarly, a study done by Singh T et al, also reported that out of 108 positive cases, most of the cases from the IPD ward (66.66%, 72/108) and rest of the positive cases from OPD 33.33%,36/108) [11]. The difference in the burden of dengue cases in IPD vs OPD emphasizes the need for an early detection and prompt treatment, in order to reduce its impact on healthcare facilities.

This study also noted that the majority of positive cases occurred during the months of October and November. A seasonal variation as reported from other endemic areas, where transmission peaks during the rainy season. Post-monsoon months are more favorable for the vector to breed due to the stagnation of rainwater. The findings underscore the importance of targeted surveillance and control measures during the peak transmission season. This result was also represented by many other Indian studies done by Badoni G et al, Kalpana S et al and Badoni G, Gupta P et al [10,12,13].

The NVBDCP data reveals the fluctuating nature of dengue cases in Delhi, with a high number of cases in 2019, a decrease in 2020, and a significant increase in 2021 [6]. Total 23 deaths were reported in 2021 as per government data [6], which is lesser than the previous year's and thus it highlights the need for continued efforts to

combat dengue, including mosquito control measures, public awareness campaigns, and accessible healthcare services to further reduce the impact of this mosquito-borne disease on the population.

The death rate associated with dengue in Delhi, India can vary from year to year and depends on various factor such as the severity of the outbreak, access to healthcare, and the effectiveness of preventive measures [6]. It is crucial for the government and healthcare authorities to continuously monitor and respond to outbreaks to minimize the impact on public health.

## 5. CONCLUSION

In conclusion, the study demonstrated a high seropositivity rate for dengue infection among clinically suspected cases, and NS1 antigen ELISA test was more effective than IgM ELISA test for early diagnosis. The findings also suggest that males and younger age groups are more prone to dengue infection. Additionally, the study highlights the importance of early diagnosis and prompt management of dengue cases to prevent severe disease and reduce morbidity and mortality. This implies the need for increased awareness, early detection, and appropriate management of dengue fever in endemic areas, especially during the peak months of October and November.

## CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

## ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

## ACKNOWLEDGEMENT

We would like to acknowledge our heartfelt gratitude to the staff of Virology Research & Diagnostic Laboratory (VRDL) and Virology Laboratory, Department of Microbiology, UMCS & GTB Hospital. We express our sincere thanks to everyone, Mr. Adesh, Mr. Mahabir Mehto, Mr. Rinkesh, Mr. Narendra Mogha, Mr. Jay Bhagwan, Ms. Poonam Mathur for their constant support and encouragement for the successful completion of the study.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Suresh A, Sreedhar KV, Mathew J, Vijayakumar KN, Ajithlal PM, Saini P, Kumar NP, Kumar A. Seroprevalence of dengue in urban and rural settings in Kerala, India. *Current Science*. 2021;121(2):233.
2. Shah PS, Alagarasu K, Karad S, Deoshatwar A, Jadhav SM, Raut T, Singh A, Dayaraj C, Padbidri VS. Seroprevalence and incidence of primary dengue infections among children in a rural region of Maharashtra, Western India. *BMC Infect Dis*. 2019;19(1):296. DOI: 10.1186/s12879-019-3937-z. PMID: 30940086; PMCID: PMC6444844
3. Trends in the Seroprevalence of Dengue in a Tertiary Care Hospital of North Karnataka, India [Internet];2022. [Cited September16, 2022];11(3):MO05-MO08.
4. Sinha S, Gahtori R, Kumari P, Kumar R, Saxena M, Pande V. et al. A Novel Multiplex RT-PCR for Simultaneous Detection of Malaria, Chikungunya and Dengue Infection (MCD- RT-PCR). *Proc. Natl. Acad. Sci., India, Sect. B Biol Sci*. 2023;93:755–761 Available:https://doi.org/10.1007/s40011-023-01447-6
5. National Guidelines for Clinical Management of Dengue Fever, NVBDCP; 2014. Available:https://nvbdcp.gov.in/WriteReadData/1892s/Dengue-National-Guidelines-2014.pdf
6. National Center for Vector Borne Diseases Control (NCVBDC), Ministry of Health & Family Welfare, Government of India. DENGUE/DHF SITUATION IN INDIA Dengue Cases and Deaths in the Country since 2015, updated on; 2022. Available:https://nvbdcp.gov.in/index4.php?lang=1&level=0&linkid=431&lid=3715
7. Kumar M, Verma RK, Mishra B. Prevalence of Dengue fever in western Uttar Pradesh, India: A Gender-Based Study. *Int J Appl Basic Med Res*. 2020 Jan-Mar;10(1):8-11. DOI: 10.4103/ijabmr.IJABMR\_337\_18. Epub 2020 Jan 3. PMID: 32002378; PMCID: PMC6967337.

8. Mehendale SM, Risbud AR, Rao JA, Banerjee K. Outbreak of dengue fever in rural areas of Parbhani district of Maharashtra (India) Indian J Med Res. 1991;93:6–11.
9. Parande A, Shinde R, Parande M, Mulla RI, Chikkaraddi U. Trends in the Seroprevalence of Dengue in a Tertiary Care Hospital of North Karnataka, India. Nat J Lab Med. 2022;11(3):M005-008.
10. Badoni G, Gupta PK, Gupta P, Kaistha N, Mathuria YP, Pai MO. et al. Dengue-chikungunya infection in the tertiary care hospital of northern India: Cross-sectional latent class cluster analysis in viral infection. Heliyon. 2023;9(3):e14019. DOI: 10.1016/j.heliyon.2023.e14019. PMID: 36925523; PMCID: PMC10011203
11. Singh T, Nigudgi A, Tiwari V, Khushwaha P, Garg A. Emergence of dengue as a febrile illness in Rewa and Nearby Districts of Madhya Pradesh during the Year, 2021: A Cross-Sectional Study. Journal of Clinical & Diagnostic Research. 2022;16(5).
12. Kalpana S, Varma G. Comparative evaluation of dengue NS1 antigen rapid test and IgM ELISA in a tertiary care hospital in VIMS, Ballari. IP Int J Med Microbiol Trop Dis. 2021;7(3):175-178.
13. Badoni G, Gupta P, Pai MO, Kaistha N, Ratho R, Sokeechand N. Dengue burden and circulation of Dengue- 2 serotype among children along with clinical profiling in Uttarakhand, India: A cross-sectional study from 2018 to 2020. Cureus. 2023; 15(1):e33913. DOI:10.7759/cureus.33913

© 2023 Jain et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/105806>