



A New Estimator of Shannon Entropy with Application to Goodness-of-Fit Test to Normal Distribution

Mbanefo S. Madukaife ^{a*}

^a Department of Statistics, University of Nigeria, Nsukka, Nigeria.

Author's contribution

The sole author designed, analyzed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/ARJOM/2023/v19i10735

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <http://www.sdiarticle5.com/review-history/104427>

Received: 15/06/2023

Accepted: 22/08/2023

Published: 28/08/2023

Original Research Article

Abstract

In this paper, a new estimator of the Shannon entropy of a random variable X having a probability density function $f(x)$ is obtained based on window size spacings. Under the standard normal, standard exponential and uniform distributions, the estimator is shown to have relative low bias and low RMSE through extensive simulation study at sample sizes 10, 20, and 30. Based on the results, it is recommended as a good estimator of the entropy. Also, the new estimator is applied in goodness-of-fit test to normality. The statistic is affine invariant and consistent and the results show that it is a good statistic for assessing univariate normality of datasets.

Keywords: Entropy estimator; window size spacing; bias of an estimator; root mean square error of an estimator; test for normality.

2010 Mathematics Subject Classification: 62E10, 62E20, 62H25.

*Corresponding author: E-mail: mbanefo.madukaife@unn.edu.ng;

Asian Res. J. Math., vol. 19, no. 10, pp. 130-139, 2023

1 Introduction

The entropy of a random variable, X , which has a distribution function $F(x)$ and a probability density function $f(x)$ is a mathematical function which measures the amount of information in the random variable. It is denoted by $H(f)$ and Shannon [1] defines it as:

$$H(f) = - \int_{-\infty}^{\infty} f(x) \log \{f(x)\} dx \tag{1.1}$$

It has been established that different random variables give different entropy functions obtained from (1.1). For instance, suppose X is a normal random variable with mean, μ and variance σ^2 , the entropy of X has been obtained as $H(f) = \frac{1}{2} + \ln(\sigma\sqrt{2\pi}) = \frac{1}{2}\ln(2\pi e\sigma^2)$ and that of a random variable X which is exponential with probability function $f(x) = \lambda e^{-\lambda x}; x > 0, \lambda > 0$ is $H(X) = -\ln\lambda + (1 - \ln\lambda)$. As a result, entropy function can be used to characterize different random variables (different distributions).

In real life however, it is usually not known the distribution from where datasets are generated. Hence, it is of utmost importance to obtain an estimator of the population parameter. Several researchers such as Vasicek [2], Ebrahimi et al. [3], Wiczorkowski and Grzegorzewski [4], Alizadeh Noughabi [5], Alizadeh Noughabi and Arghami [6], Zamanzede and Arghami [7], Al-Omari [8], Lombardi and Pant [9], Kohansal and Rezakhah [10], and Bitaraf et al. [11], to mention but a few, have devoted research attention to this direction. These estimators are obtained using different nonparametric approaches such as window size (m) spacings, kernel density estimation, nearest neighbour technique, and so on. The window size approach, no doubt, has attracted more researchers the rest of other approaches put together. Let x_1, x_2, \dots, x_n be a random sample of size n from a random variable X whose probability law is $f(x)$ with distribution function $F(x)$. Also, let the sample order statistics obtained from the random sample be $X_{(1)}, X_{(2)}, \dots, X_{(n)}$ such that $X_{(1)} \leq X_{(2)} \leq \dots \leq X_{(n)}$. For a non-negative integer m such that $m = 0, 1, 2, \dots, \frac{n}{2}$, the sample m spacing is defined on j th order statistic by $X_{(j+m)} - X_{(j-m)}$. Using such spacings, Vasicek [2] in his pioneer work in this direction, estimated the derivative of $F^{-1}(p)$ by the slope given as:

$$[X_{(j+m)} - X_{(j-m)}] \frac{n}{2m} \tag{1.2}$$

for $\frac{(j-1)}{n} < p < \frac{j}{n}; j = m + 1, m + 2, \dots, n - m$ after transforming (1) in the form:

$$H(f) = \int_0^1 \log \left\{ \frac{d}{dp} F^{-1}(p) \right\} dp; p \in (0, 1) \tag{1.3}$$

to birth an estimator given by:

$$HV_{mn} = \frac{1}{n} \sum_{j=1}^n \log \left\{ \frac{n}{2m} [X_{(j+m)} - X_{(j-m)}] \right\}; \tag{1.4}$$

where $X_{(j)} = X_{(1)}$ for $j < 1$ and $X_{(j)} = X_{(n)}$ for $j > n$. Although this estimator is biased, like all the other non-parametric estimators, it is conceived to have a theoretically sound background to making a good estimator of $H(f)$.

Soon after the development of the estimator in (1.4), a number of researchers criticised the slope in (1.2) as incorrect when $j \leq m$ or $j > n - m + 1$. This triggered different modifications of the slope to arrive at different estimators of the Shannon entropy with comparatively better results. Examples of such estimators obtained in this light include the Ebrahimi et al. [3], Alizadeh Noughabi and Arghami [6] and Al-Omari [12, 8]. Till present however, there has not been a generally accepted estimator with respect to correctness of the slope when $j \leq m$ or $j > n - m + 1$, hence this research paper. Several other nonparametric estimators of the entropy have also been proposed in the literature which are not based on the window size spacings. For instance, Zhang [13] obtained an estimator of the entropy and Zhang [14] showed that the estimator is asymptotically normal with an exponentially decaying bias.

In this work, a modification is introduced to the Bitaraf et al. [11] approach to obtain a competitively new estimator of the Shannon entropy which is applied to goodness-of-fit test to normality of datasets. In what follows, the new estimator is obtained in Section 2. Section 3 gives the Monte Carlo comparison of the estimator with some other existing ones. The application to testing for normality is carried out in Section 4 while the paper is concluded in Section 5.

2 The New Estimator

Let X_1, X_2, \dots, X_n be a random sample of size n from a continuous distribution $F(x)$ having a probability function $f(x)$. Also, let $X_{(1)}, X_{(2)}, \dots, X_{(n)}$ be the order statistics of the sample such that $X_{(1)} \leq X_{(2)} \leq \dots \leq X_{(n)}$. Using the sample order statistics, Ebrahimi et al. [3] modified the Vasicek estimator to propose a new estimator given by:

$$HEb_{mn}^{(1)} = \frac{1}{n} \sum_{i=1}^n \log \left\{ \frac{n}{c_i m} (X_{(i+m)} - X_{(i)}) \right\}, \tag{2.1}$$

where $c_i = \begin{cases} 1 + \frac{i-1}{m}, & 1 \leq i \leq m, \\ 2, & m+1 \leq i \leq n-m, \\ 1 + \frac{n-i}{m}, & n-m+1 \leq i \leq n, \end{cases}$ $X_{(i-m)} = X_{(1)}$ for $i \leq m$ and $X_{(i+m)} = X_{(n)}$ for $i \geq n-m$.

Then, Bitaraf et al. [11] introduced an internal j th spacings, in addition to the m th spacings, thereby changing the normal slope in (1.2) to have:

$$T_i = \frac{1}{2} \sum_{j=0}^1 T_{ij}; T_{ij} = \frac{n}{w_j(m-j)} \{X_{(i+m-j)} - X_{(i-m+j)}\} \tag{2.2}$$

for $\frac{(i-1)}{n} < p < \frac{i}{n}; i = m+1, m+2, \dots, n-m$, where $w_j = \begin{cases} 1, & 1 \leq i \leq m-j \\ 2, & m-j+1 \leq i \leq n-m+j \\ 1, & n-m+j+1 \leq i \leq n \end{cases}$ and

$j = 0, 1$. Based on (2.2), they obtained an estimator of the Shannon entropy as:

$$HB_{mn} = \frac{1}{n} \sum_{i=1}^n \log \{T_i\} \tag{2.3}$$

and showed empirically that the estimator has smaller RMSE and bias than the Ebrahimi et al. [3]. Due to the interesting properties of the j th internal spacing estimator of Bitaraf et al. [11], a new estimator of the entropy is proposed, which modifies the weighting factor of the Bitaraf et al. [11] estimator. The proposed estimator is given by:

$$HM_{mn} = \frac{1}{n} \sum_{i=1}^n \log \left\{ \frac{1}{2} (T_{i0} + T_{i1}) \right\} \tag{2.4}$$

where $T_{ij} = \frac{n}{w_{ij}(m-2j)} \{X_{(i+m-j)} - X_{(i-m+j)}\}$ and $w_{ij} = \begin{cases} 1 + \frac{1}{3}, & 1 \leq i \leq m-j \\ 2, & m-j+1 \leq i \leq n-m+j \\ 1 + \frac{1}{3}, & n-m+j+1 \leq i \leq n \end{cases}$; $X_{(i-m+j)} = X_{(1)}$ for $i \leq m-j$ and $X_{(i+m-j)} = X_{(n)}$ for $i \geq n-m+j$, $\frac{n}{m} \rightarrow 0, j = 0, 1$

Theorem 2.1: For a random sample X_1, X_2, \dots, X_n from a continuous distribution $F(x)$ with a probability density function $f(x)$, $HM_{mn} \xrightarrow{P} H(f)$ as $n \rightarrow \infty, m \rightarrow \infty$, and $\frac{n}{m} \rightarrow 0$.

Proof: Using a property of the convex function, Bitaraf et al. [11] obtained the inequality:

$$\frac{1}{2n} \sum_{i=1}^n (\log a_i + \log b_i) \leq \frac{1}{n} \sum_{i=1}^n \log \left(\frac{1}{2} (a_i + b_i) \right) \leq \frac{1}{n} \sum_{i=1}^n \max \{ \log a_i, \log b_i \}; \tag{2.5}$$

where $a_i, b_i \in R^+$. Taking T_{i0} for a_i and T_{i1} for b_i and following the same principle with Bitaraf et al. [11], we have the extreme terms of the inequality in (2.5) presented as:

$$\begin{aligned} & \frac{1}{2n} \sum_{i=1}^n \left(\log \left\{ \frac{n(X_{(i+m)} - X_{(i-m)})}{2m} \right\} + \log \left\{ \frac{n(X_{(i+m-1)} - X_{(i-m+1)})}{2(m-2)} \right\} + \log \left\{ \frac{2}{w_{i0}} \right\} + \log \left\{ \frac{2}{w_{i1}} \right\} \right) \\ & \leq \frac{1}{n} \sum_{i=1}^n \max \left(\log \left\{ \frac{n(X_{(i+m)} - X_{(i-m)})}{2m} \right\}, \log \left\{ \frac{n(X_{(i+m-1)} - X_{(i-m+1)})}{2(m-2)} \right\} \right) \end{aligned}$$

Since $w_{ij} \in (1, 2)$, $(2n)^{-1} \sum_{i=1}^n \log \left\{ \frac{2}{w_{i0}} \right\}$ and $(2n)^{-1} \sum_{i=1}^n \log \left\{ \frac{2}{w_{i1}} \right\}$ converges to 0. Also, there exists $X_{ij} \in (X_{(i-m+j)}, X_{(i+m-j)})$ such that

$$\frac{F(X_{(i+m-j)}) - F(X_{(i-m+j)})}{X_{(i+m-j)} - X_{(i-m+j)}} = f(X_{ij}); j = 0, 1.$$

In line with Bitaraf [11], the first and second terms of the inequality above give rise to

$$\frac{1}{2} \left[-\frac{1}{n} \sum_{i=1}^n \log \{f(X_{i0})\} - V_{m,n} + U_{m,n} - \frac{1}{n} \sum_{i=1}^n \log \{f(X_{i1})\} - V_{m-2,n} + U_{m-2,n} \right],$$

where $V_{m-2j,n} = \frac{1}{n} \sum_{i=1}^n \log \left\{ \frac{F(X_{(i+m-j)}) - F(X_{(i-m+j)})}{f(X_{ij})[X_{(i+m-j)} - X_{(i-m+j)}]} \right\}$, and

$U_{m-2j,n} = \frac{1}{n} \sum_{i=1}^n \log \left\{ \frac{n}{2^{(m-2j)}} [F(X_{(i+m-j)}) - F(X_{(i-m+j)})] \right\}$; $j = 0, 1$. According to Vasicek [2] and Bitaraf et al. [11], for a fixed n , the effect of $V_{m-j,n}$ decreases as m decreases and $U_{m-j,n}$ converges in probability to zero. Hence, HM_{mn} converges in probability to $H(f)$ as $n \rightarrow \infty, m \rightarrow \infty$, and $\frac{n}{m} \rightarrow 0$ and the proof is completed.

3 Empirical Comparison of Entropy Estimators

In this section, the performance of some selected three estimators of the Shannon entropy is compared with the proposed estimator through simulation studies. The selected estimators include the Vasicek [2], Ebrahimi et al. [3], and the Bitaraf et al. [11] estimators, which are presented in this work as HV_{mn} , HE_{mn} , and HB_{mn} respectively. Precisely, the empirical root mean square error (RMSE) and the bias of these three competing estimators are compared with those of the proposed HM_{mn} estimator based on 10,000 samples of sizes 10, 20, and 30. The sample sizes of 10, 20 and 30 are chosen for the study since they span through small sample to large sample sizes. As a result, they can represent very well the general behaviour of an estimator across sample sizes. These samples are obtained from three different continuous distributions, namely: the standard normal, uniform in the interval (0, 1), and the standard exponential, which are chosen arbitrarily among all the possible statistical distributions. In each of the sample sizes, the study is conducted for $m = 1, 2, \dots, \frac{n}{2}$. Although Grzegorzewski and Wieczorkowski [15] have obtained an optimal m as $m^* = [\sqrt{n} + 0.5]$, but it has been criticised as being incorrect, hence our choice of using all the possible m for each sample size.

From each sample size under each distribution with a known measure of entropy θ , 10,000 samples are generated and the estimated entropy measure, $\hat{\theta}$, is computed from each of the samples using the four competing estimators. Then, the RMSE and bias of each of the estimators is obtained by:

$$RMSE = \left(\frac{1}{10000} \sum_{j=1}^{10000} \{ \hat{\theta}_j - \theta \}^2 \right)^{\frac{1}{2}}; bias = \frac{1}{10000} \sum_{j=1}^{10000} \hat{\theta}_j - \theta. \tag{3.1}$$

All the simulations and computations are carried out using the R statistical software and the results are presented in Tables 1, 2, and 3 for normal, uniform and exponential distributions respectively.

Table 1. Empirical root mean square errors and biases of some Shannon entropy estimators under the standard normal distribution with $\theta = 1.4189$

n	m	RMSEs				Biases			
		HV_{mn}	HE_{mn}	HB_{mn}	HM_{mn}	V_{bias}	E_{bias}	B_{bias}	M_{bias}
10	1	0.6719	0.5559	–	1.2042	-0.6718	-0.3508	–	-1.1068
	2	0.5955	0.4297	0.4115	–	-0.1791	-0.4041	-0.1882	–
	3	0.6190	0.4045	0.3239	0.2789	-0.6090	-0.0066	-0.1761	-0.2118
	4	0.6630	0.3936	0.2781	0.2747	-0.4730	-0.3832	0.2768	-0.4054
	5	0.7165	0.3852	0.2591	0.2864	-0.2858	-0.1560	0.3527	-0.2146
20	1	0.4859	0.4243	–	1.1013	-0.4881	-0.4188	–	-1.1863
	2	0.3770	0.2957	0.2964	–	-0.2671	-0.1690	-0.6345	–
	3	0.3662	0.2631	0.2220	0.2711	0.2124	0.3404	-0.2391	0.4778
	4	0.3762	0.2485	0.1912	0.1894	-0.4731	-0.3147	-0.0624	0.2219
	5	0.3924	0.2395	0.1800	0.1782	-0.8076	-0.6187	-0.0106	0.0491
	6	0.4198	0.2408	0.1818	0.1781	-0.2402	-0.0207	-0.1301	0.0298
	7	0.4453	0.2375	0.1898	0.1808	-0.7344	-0.4843	0.0316	0.0711
	8	0.4736	0.2389	0.2063	0.1806	-0.1426	0.1381	0.3092	-0.2776
	9	0.5018	0.2380	0.2290	0.1803	-0.5254	-0.2141	0.3924	-0.0604
	10	0.5268	0.2353	0.2549	0.1791	-0.6992	-0.3573	0.0472	0.1024
30	1	0.4130	0.3710	–	1.0550	-0.5199	-0.4737	–	-1.1980
	2	0.2987	0.2441	0.2515	–	-0.3975	-0.3321	-0.3971	–
	3	0.2749	0.2072	0.1879	0.2752	-0.4684	-0.3830	-0.0931	0.5519
	4	0.2734	0.1918	0.1587	0.1735	-0.1492	-0.0436	-0.0146	0.0348
	5	0.2831	0.1854	0.1474	0.1488	-0.0057	0.1202	0.1881	0.2138
	6	0.2904	0.1788	0.1432	0.1443	-0.2438	-0.0972	-0.2126	0.0689
	7	0.3051	0.1770	0.1528	0.1429	-0.3850	-0.1683	0.0965	-0.1639
	8	0.3209	0.1768	0.1620	0.1478	-0.3329	-0.1458	-0.1017	-0.0268
	9	0.3376	0.1753	0.1807	0.1466	-0.6350	-0.4275	0.1052	-0.1991
	10	0.3553	0.1762	0.1977	0.1510	-0.3276	-0.0996	0.0971	-0.3528
	11	0.3728	0.1758	0.2139	0.1506	-0.3077	-0.0593	0.0383	-0.0390
	12	0.3871	0.1733	0.2362	0.1527	-0.4180	-0.1492	0.2273	-0.2422
	13	0.4091	0.1741	0.2597	0.1555	-0.4504	-0.1611	0.2060	-0.0603
	14	0.4274	0.1748	0.2805	0.1552	-0.5827	-0.2730	0.1564	0.0003
	15	0.4479	0.1764	0.3037	0.1579	-0.4534	-0.1233	0.2449	-0.0288

Table 2. Empirical root mean square errors and biases of some Shannon entropy estimators under the standard exponential distribution with $\theta = 1$

n	m	RMSEs				Biases			
		HV_{mn}	HE_{mn}	HB_{mn}	HM_{mn}	V_{bias}	E_{bias}	B_{bias}	M_{bias}
10	1	0.6742	0.5704	—	1.2278	-0.6411	-0.2621	—	-1.0727
	2	0.5777	0.4376	0.4271	—	-0.2224	0.0063	0.2789	—
	3	0.5673	0.3984	0.3619	0.4002	-0.3692	-0.2055	0.2221	0.5587
	4	0.5738	0.3905	0.3633	0.3737	-0.0775	0.2163	0.3187	0.5347
	5	0.5970	0.3794	0.4183	0.3727	-0.5236	-0.1298	0.0118	-0.5849
20	1	0.4874	0.4297	—	1.0949	-0.4684	-0.4558	—	-1.4015
	2	0.3774	0.3107	0.3201	—	-0.3003	-0.2719	0.3241	—
	3	0.3583	0.2790	0.2528	0.3363	-0.3846	-0.0740	-0.3377	0.2598
	4	0.3555	0.2613	0.2418	0.2671	-0.3429	-0.3365	-0.3971	-0.1285
	5	0.3563	0.2529	0.2517	0.2531	-0.1613	-0.3875	0.0975	0.0736
	6	0.3572	0.2496	0.2799	0.2643	-0.4417	0.0610	0.3004	0.1359
	7	0.3675	0.2484	0.3176	0.2679	-0.1151	0.0599	0.2417	0.0084
	8	0.3769	0.2566	0.3605	0.2773	-0.1169	0.0393	0.2599	0.1986
	9	0.3818	0.2628	0.4066	0.2960	-0.0756	-0.0904	0.3443	0.3686
	10	0.3954	0.2719	0.4680	0.3139	-0.1185	0.1554	0.4483	0.5568
30	1	0.4207	0.3802	—	1.0530	-0.5638	-0.2902	—	-0.9760
	2	0.3065	0.2629	0.2720	—	-0.1007	0.0792	-0.1688	—
	3	0.2827	0.2285	0.2150	0.3254	-0.2235	-0.2701	-0.1336	0.4435
	4	0.2736	0.2125	0.1975	0.2283	-0.2235	-0.1100	-0.3329	0.0932
	5	0.2722	0.2048	0.1957	0.2132	-0.5552	-0.2931	0.3241	0.3032
	6	0.2719	0.1996	0.2073	0.2102	-0.2572	-0.1826	-0.1514	-0.0942
	7	0.2752	0.1988	0.2300	0.2128	-0.1886	-0.2609	0.0404	0.1135
	8	0.2802	0.2026	0.2543	0.2162	-0.2200	0.1126	0.0583	-0.1477
	9	0.2777	0.2021	0.2859	0.2305	-0.0324	0.2507	0.0171	-0.0296
	10	0.2861	0.2082	0.3218	0.2388	-0.4808	0.0820	0.1876	-0.1442
	11	0.2896	0.2141	0.3571	0.2492	-0.4216	-0.0712	-0.0542	0.1880
	12	0.2953	0.2211	0.3936	0.2652	0.0961	0.2786	0.5240	0.2406
	13	0.2979	0.2312	0.4384	0.2811	-0.0932	0.1771	0.1063	0.0724
	14	0.3044	0.2422	0.4731	0.2983	-0.2753	0.3789	0.3739	0.6182
	15	0.3031	0.2543	0.5213	0.3179	0.0411	-0.2888	0.3556	0.1823

Table 3. Empirical root mean square errors and biases of some Shannon entropy estimators under the uniform distribution in the interval 0 and 1, with $\theta = 0$

n	m	RMSEs				Biases			
		HV_{mn}	HE_{mn}	HB_{mn}	HM_{mn}	V_{bias}	E_{bias}	B_{bias}	M_{bias}
10	1	0.5698	0.4461	—	1.1567	-0.5854	-0.2439	—	-1.1061
	2	0.4520	0.2839	0.2712	—	-0.2184	0.0144	-0.1331	—
	3	0.4534	0.2308	0.1769	0.2590	-0.5832	-0.2226	0.0942	0.2940
	4	0.4833	0.2165	0.1704	0.1769	-0.3333	-0.0827	0.1686	-0.0642
	5	0.5305	0.2058	0.2255	0.1664	-0.5598	0.1308	0.2110	0.0939
20	1	0.4193	0.3555	—	1.0585	-0.2526	-0.2007	—	-0.9864
	2	0.2884	0.2017	0.2081	—	-0.1701	-0.1565	-0.2188	—
	3	0.2702	0.1556	0.1195	0.2666	-0.4173	-0.0834	-0.1081	0.1787
	4	0.2751	0.1324	0.0878	0.1421	-0.1140	-0.0069	0.0012	0.2632
	5	0.2904	0.1231	0.0990	0.1104	-0.1942	-0.0315	0.1244	0.0452
	6	0.3105	0.1131	0.1298	0.1001	-0.3145	-0.0148	0.0864	0.1174
	7	0.3326	0.1086	0.1677	0.0959	-0.3174	-0.2053	0.1870	0.0574
	8	0.3599	0.1083	0.2084	0.0969	-0.2569	-0.0794	0.1652	0.1306
	9	0.3858	0.1077	0.2479	0.1002	-0.3297	-0.0170	0.2803	-0.1262
	10	0.4115	0.1061	0.2900	0.1038	-0.4326	-0.0760	0.26462	0.1206
30	1	0.3697	0.3245	—	1.0264	-0.3457	-0.3587	—	-1.0021
	2	0.2352	0.1757	0.1883	—	-0.1430	-0.1442	-0.1214	—
	3	0.2080	0.1314	0.1062	0.2758	-0.1615	-0.0652	-0.1184	0.1452
	4	0.2043	0.1094	0.0678	0.1402	-0.1870	-0.0786	-0.1437	0.1991
	5	0.2116	0.0963	0.0619	0.0996	-0.1363	0.0008	0.0485	0.1163
	6	0.2217	0.0895	0.0781	0.0837	-0.3114	-0.0610	-0.0716	0.1146
	7	0.2344	0.0835	0.1019	0.0781	-0.2111	0.0190	0.0827	0.1617
	8	0.2490	0.0784	0.1288	0.0745	-0.2545	-0.0812	0.1460	0.0842
	9	0.2658	0.0764	0.1563	0.0735	-0.2719	-0.2071	0.1949	0.0441
	10	0.2827	0.0756	0.1828	0.0761	-0.3348	-0.0527	0.1564	0.0251
	11	0.3013	0.0727	0.2101	0.0775	-0.3395	-0.0082	0.2102	0.1134
	12	0.3194	0.0724	0.2376	0.0800	-0.3262	-0.0182	0.3247	0.0174
	13	0.3369	0.0732	0.2656	0.0831	-0.3025	-0.0229	0.1635	0.1199
	14	0.3557	0.0728	0.2924	0.0887	-0.3883	-0.1183	0.2449	0.1052
	15	0.3740	0.0726	0.3179	0.0915	-0.3555	-0.0375	0.2767	0.0189

From the results in Tables 1, 2, and 3, a number of important facts about the general behaviour of the Shannon entropy estimators as well as specific estimators are established. Firstly, the efficiency behaviour of the estimators depends on the distribution from where the sample is obtained. This is because the minimum RMSE of each of the competing estimators under a specified sample size varies across the distributions. For instance, at sample size of 10, HV_{mn} , recorded minimum RMSEs of 0.5955, 0.5673, and 0.4520 under normal, exponential and uniform distributions respectively. At a sample size of 20, the observed minimum RMSEs are 0.3662, 0.3563, and 0.2702 under normal, exponential and uniform distributions respectively. In a similar manner, minimum RMSEs of 0.2734, 0.2719, and 0.2043 were obtained for the same HV_{mn} at sample size of 30 under the same respective distributions. For the HE_{mn} , ordered sets of minimum RMSEs, in the manner of the HV_{mn} were obtained as (0.3852, 0.3794, 0.2058); (0.2353, 0.2484, 0.1061); and (0.1733, 0.1988, 0.0724) respectively for sample sizes 10, 20, and 30 respectively. For the HB_{mn} estimator, the ordered sets of the minimum RMSEs are (0.2591, 0.3619, 0.1704); (0.1800, 0.2418, 0.0878) and (0.1432, 0.1957, 0.0619) respectively for sample sizes $n = 10, 20,$ and 30 while those of the new estimator, HV_{mn} , are (0.2747, 0.3727, 0.1664); (0.1781, 0.2531, 0.0959) and (0.1429, 0.2102, 0.0735) respectively for sample sizes $n = 10, 20,$ and 30 .

Secondly, the RMSEs for each of the estimators across the distributions considered decrease with increasing sample size. This is however expected for every good estimator. Again, it is observed that the window size m at which the RMSE is a minimum (also known as the optimal window size) for a specified sample size varies from one distribution to another and from one estimator to another. This is against Grzegorzewski and Wieczorkowski [15], who proposed optimal window size $m^* = [\sqrt{n} + 0.5]$, where $[x]$ is the integral part of x . For each of the estimators, the optimal window size, m^* , is presented in Table 4 for each sample size under different distributions with that of Grzegorzewski and Wieczorkowski [15] denoted by $m^*(GW)$. The results in Table 4 show that all the estimators, except the HV_{mn} , maintained optimal window size generally greater than the $m^*(GW)$ in almost all the distributions considered.

Table 4. Optimal window size of different estimators in comparison with $m^*(GW) = [\sqrt{n} + 0.5]$

Distributions	n	$m^*(GW)$	$m^*(HV_{mn})$	$m^*(HE_{mn})$	$m^*(HB_{mn})$	$m^*(HM_{mn})$
Normal	10	3	2	5	5	4
	20	5	3	10	5	6
	30	6	4	12	6	7
Exponential	10	3	3	5	3	5
	20	5	6	7	4	5
	30	6	6	7	5	6
Uniform	10	3	2	5	4	5
	20	5	3	10	4	7
	30	6	4	12	5	9

In addition to the above, the results in Tables 1, 2, and 3 show that HE_{mn} , HB_{mn} , and HM_{mn} generally have smaller RMSEs and bias values than the HV_{mn} in all the sample sizes and distributions considered. As a result, they can be regarded as bias corrected and generally more efficient estimators than the HV_{mn} . Specifically, the HB_{mn} and HM_{mn} recorded least RMSEs in all the sample sizes and distributions considered except in the standard exponential distribution, where the HE_{mn} recorded lower RMSEs than the HM_{mn} in sample sizes $n = 20$ and 30 . Finally, the new HM_{mn} estimator recorded higher RMSEs than the HB_{mn} in the sample size of $n = 10$ of the normal distribution and $n = 20$ and 30 of the uniform distribution. Based on these results, the proposed estimator can be regarded as a good estimator of the Shannon entropy.

4 Test of Normality Based on The New Estimator

It is well known that among all the statistical distributions that possess a density function $f(x)$ and have a given variance, σ^2 , the entropy, $H(f)$, is maximized by the normal distribution. Based on this, Vasicek [2] obtained a statistic for testing the assumption of normality. Now, following the procedure of Vasicek [2], a new test for testing the normality of datasets is obtained by

$$M_{mn} = \frac{1}{s} \prod_{i=1}^n \left\{ \frac{n(X_{(i+m)} - X_{(i-m)})}{2w_{i0}m} + \frac{n(X_{(i+m-1)} - X_{(i-m+1)})}{2w_{i1}(m-2)} \right\}^{\frac{1}{2}} \quad (4.1)$$

where $s^2 = n^{-1} \sum_{i=1}^n (x_i - \bar{X})^2$. The statistic is consistent and affine invariant.

The M_{mn} statistic is applied to test for the normality of four different datasets. The first three datasets are simulated from the standard normal, standard exponential, and standard lognormal distributions, having sample size of 30 in each case while the last dataset is the petal length of the iris setosa dataset, see Seber [16]. Expectedly, the test rejected the normality of all the datasets at 5% level of significance, except the standard normal distribution. The results therefore show that the statistic is a good test for testing normality of datasets.

5 Conclusion

Several estimators of the Shannon entropy already exists in the literature via window size spacings. All the estimators are however biased with varied degrees of biasedness. Also, the ability of these estimators to be appropriate for the true population entropy depends largely on their relative efficiencies, measured in terms of the MSEs (or RMSEs). In this study, the new bias corrected estimator was obtained by a modification of the Bitaraf et al. [11] estimator. In addition to having relatively low bias, the new estimator also has relatively low RMSE compared to some other estimators. As a result, it can be recommended as a good estimator of the Shannon entropy. Also, the M_{mn} statistic obtained for testing normality which is based on the new estimator can be recommended as a good test for normality of datasets. This is based on the ability of the test to reject the normality of the four datasets which are actually non-normal.

Competing Interests

Author has declared that no competing interests exist.

References

- [1] Shannon CE. A mathematical theory of communications. Bell System Technical Journal. 1948;27:379-423,623-656.
DOI:10.1002/bltj.1948.27.issue-3
- [2] Vasicek O. A test for normality based on sample entropy. Journal of the Royal Statistical Society B. 1976;38:54-59.
- [3] Ebrahimi N, Pflughoeft K, Soofi ES. Two measures of sample entropy. Statistics and Probability Letters. 1994;20:225-234.
- [4] Wiczorkowski R, Grzegorzewsky P. Entropy estimators improvements and comparisons. Communication in Statistics-Simulation and Computation. 1999;28(2):541-567.
DOI:10.1080/03610919908813564
- [5] Alizadeh Noughabi H. A new estimator of entropy and its application in testing normality. Journal of Statistical Computation and Simulation. 2010;80(10):1151-1162.

- [6] Alizadeh Noughabi H, Arghami NR. A new estimator of entropy. *Journal of the Iranian Statistical Society*. 2010;9(1):53-64.
- [7] Zamanzade E, Arghami NR. Testing normality based on new entropy estimators. *Journal of Statistical Computation and Simulation*. 2012;82(11):1701-1713.
- [8] Al-Omari AI. A new measure of entropy of continuous random variable. *Journal of Statistical Theory and Practice*. 2016;10:721-735.
DOI: 10.1080/15598608.2016.1217444
- [9] Lombardi M, Pant S. Nonparametric k-nearest-neighbor entropy estimator. *Physical Review E*. 2016;93:013310.
- [10] Kohansal A, Rezakhah S. Two new estimators of entropy for testing normality. *Communications in Statistics – Theory and Methods*. 2016;45(18):5392-5411.
DOI: 10.1080/03610926.2014.942436
- [11] Bitaraf M, Rezaei M, Yousefzadeh F. Test for normality based on two new estimators of entropy. *Journal of Statistical Computation and Simulation* 2017;87(2):280-294.
DOI: 10.1080/00949655.2016.1208201
- [12] Al-Omari AI. Estimation of entropy using random sampling. *Journal of Computation and Applied Mathematics*. 2014;261:95-102.
DOI:10.1016/j.cam.2013.10.047.
- [13] Zhang Z. Entropy estimation in Turing’s perspective. *Neural Computation*. 2012;24(5):1368-1389.
- [14] Zhang Z. Asymptotic normality of an entropy estimator with exponentially decaying bias. *IEEE Transactions on Information Theory*. 2013;59(1):504-508.
- [15] Grzegorzewski P, Wiczorkowski R. Entropy-based goodness-of-fit test for exponentiality. *Commun Stat - Theory Methods*. 1999;28:1183-1202.
- [16] Seber WAF. *Multivariate observations*, 2nd edition. John Wiley and Sons Inc. New York; 2004.

© 2023 Madukaife; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<http://www.sdiarticle5.com/review-history/104427>