



Diabetes Mellitus Control Including Fruits in Diet: Exhaustive Review and Meta-analysis

**Md. Shawan Ali ^{a*}, Shanawaz Ahmed ^a,
Md. Rockibul Islam ^a, Md. Sazu Ahamed ^a,
Md. Ashikur Rahaman ^a, Mst. Khodeza Khatun ^{a*}
and Md. Ashraful Alam ^{a,b*}**

^a *Department of Applied Chemistry and Chemical Engineering, Islamic University, Kushtia-7003, Bangladesh.*

^b *Institute of Glass and Ceramic Research and Testing (IGCRT), Bangladesh Council of Scientific and Industrial Research (BCSIR), Dhaka-1205, Bangladesh.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111953>

Review Article

Received: 11/11/2023
Accepted: 15/01/2024
Published: 22/01/2024

ABSTRACT

A class of illnesses known as diabetes mellitus impacts how our bodies utilize glucose or blood sugar in oxidation control by human lunch. It serves as a significant source of energy for the cells that comprise our muscles and tissues which are essential to our health. It's also the primary energy source for our brain. Each kind of diabetes has a different underlying aetiology. Type 1 and type 2 diabetes are one types of chronic diabetes diseases. Prediabetes and gestational diabetes are two diabetes disorders that may be treated. When our blood sugar permitted levels are greater than usual but not high enough to be diagnosed with diabetes. Unless the right steps are taken to

*Corresponding author: E-mail: shawanacce@gmail.com, shima_acct@yahoo.com, ashrafulalam@bcsir.gov.bd;

stop the progression, prediabetes frequently precedes diabetes. People with type 2 diabetes can still benefit from fruits but only if they avoid ones with high sugar and carbohydrate content which might raise blood glucose levels by hydrolysis. To manage diabetes 2, they need to include fruits with a low Glycemic Index. Cherries, prunes, grapefruit, dried apricots, raisins, peaches, apples, pears, strawberries, plums, guava, orange, grapes, papaya, banana, kiwi, pineapple, figs and mango are among the fruits that have low glycemic index that is low in carbohydrates or high in protein and fiber which can reduce blood sugar spikes. Focusing on the glycemic index in food intake behaviour is the prime object of this review manuscript.

Keywords: Diabetes mellitus; diet; fruits; insulin; type 1; type 2.

1. INTRODUCTION

Diabetes is a condition of the metabolism of carbohydrates in which the body does not utilize the sugars found in meals as it should. It is typified by elevated blood glucose levels and glycosuria brought on by insulin resistance and malfunctioning beta cells in the pancreas [1]. The body uses glucose or sugar, typically produced when part of the food is broken down as fuel. Insulin is a hormone that permits glucose to enter cells after the blood carries or maintains it. Insulin cannot be used correctly by diabetics or they generate too little or none at all [2]. The metabolism of lipids and proteins is also changed in severe stages or quantities of diabetes. Numerous elements including age, sedentary lifestyle, sex, food, obesity, socioeconomic level, genetics and hypertension, among others by birth or genetically [2]. Reduced β -cell mass, decreased insulin secretion and a course of progressive insulin resistance in the liver and peripheral tissues are related to the aetiology of type 2 diabetes whereas abnormalities in insulin secretion cause type 1 diabetes. Hyperglycemia, glycosuria, hyperlipidemia and atherosclerosis are all consequences of altered protein, fat and carbohydrate metabolism brought on by diabetes mellitus [3]. Long-term harm, organ failure and destruction are linked to hyperglycemia brought on by diabetes [4]. A wide range of conditions or parameters including fatty changes, liver cirrhosis and cancer can affect the liver or other organs as well as tissues or cells [5]. An essential component of lipid and glucose balance the liver is an organ independent of insulin. Furthermore, research has demonstrated that tissue damage brought out by diabetes might be responsible for its activity [6,7,9]. A greater amount of study has been conducted on the delayed sequelae of diabetes which include nephropathy, retinopathy, cardiovascular problems, neuropathy, skin ulcers, high blood pressure and weight gain [9,10,11]. The number of patients affected with diabetes being referred

to physicians is experiencing a substantial increase. Diabetes currently afflicts over 140 million individuals in Western countries and this figure is projected to rise to 300 million by the year 2025 [2,10]. Artificial medications are employed to reduce the onset of diabetes but are accompanied by adverse reactions [13]. The search for natural components that work well has become more popular or famous these days because of this problem and their expensive price or availability. Herbal remedies have been the subject of numerous investigations in recent years and these substances have been suggested as treatments for several incurable illnesses, including cancer, neurological disorders, cardiovascular disease and atherosclerosis. Medicinal plants with antioxidant activity can combat these diseases which include many modifications, including shifts in redox state. These plants have long been recognized as a beneficial source of health promotion [3,4,14]. Patients affected with diabetes employed traditional medicine and medicinal herbs before anti-diabetic medications and insulin were discovered. There are various ways that plants or plant extracts might influence blood sugar levels [5,15,15a,15b]. A portion of them might possess insulin kinase while others might suppress the activity of inulinase and still others might promote the regeneration of pancreatic β -cells. Plant fiber may also obstruct the absorption of carbs hence influencing blood sugar levels [6] and honey, vegetable and peel-enrich food is also responsible [15c-15e].

2. SYMPTOMS

While the symptoms of the two forms of diabetes are similar differ in severity. In type 1 diabetes, symptoms appear more frequently and develop more quickly. Polyurea, polydipsia, polyphagia, weight loss, tiredness, cramps, constipation, hazy eyesight and candidiasis are some of the symptoms. Patients with type 1 diabetes for an extended period are at risk for both

macrovascular illness (heart, peripheral vascular, and coronary artery disorders) and microvascular problems [6,16]. Similar symptoms but a sneakier start characterize type 2 diabetes. The majority of instances are discovered accidentally or as a result of difficulties. Large artery atherosclerosis which is frequently linked to obesity, hypertension and hyperlipidemia is a significant risk factor for type 2 diabetes. End-stage renal disease and cardiovascular problems account for the majority of type 2 diabetes deaths. There are regional variations in the severity of these issues as well as their [6,17].

2.1 Types of Diabetes

2.1.1 Type 1 diabetes

Type 1 diabetes sometimes referred to as insulin-dependent diabetes (IDD), is less prevalent than type-2 diabetes and typically manifests during childhood. The prevalence of type 1 diabetes is about 5–10%. Type 1 diabetes prevents the synthesis or reproduction of insulin. Beta cells which are located in the pancreas produce insulin. There is a possibility of the elimination of beta cells. Multiple factors can lead to the demise of beta cells [5,6,18]. Most persons with type 1 diabetes have an immune system malfunction however, certain individuals have a hereditary predisposition to the condition. The body's immune cells erroneously target and eliminate beta cells instead of fulfilling their intended function. Glucose rises in the blood and the pancreas is unable to make insulin without beta cells. Cell membranes restrict the entry of sugar in the absence of insulin. Diabetes type 1 symptoms appear quickly and are characterized by a multifunctional system at conditioning system [5,6,19].

2.1.2 Type 2 diabetes

About 90–95% of diabetics have type-2 or non-insulin dependent diabetes (NIDD) which is more

frequent than adult-onset type-1 diabetes. Insulin resistance a condition in which the body cells do not respond to insulin or the pancreas does not produce enough insulin is the cause of type 2 diabetes [5,6]. Diabetes type 2 symptoms appear gradually. Many persons with type 2 diabetes are unaware of their condition until they visit their doctor due to a health issue. The patient may have high blood glucose levels for seven to ten years before being diagnosed with type 2 diabetes. Even if the blood contains an abundance of insulin in this form the cells are not receptive to it. Because glucose enters cells more difficultly and backs up in the bloodstream [5,6]. Most persons with Type 2 diabetes are older adults, overweight and have a blood relative who has the condition. Type-2 diabetes typically affects people over 40 while it frequently affects people from South Asia and the African Caribbean before 40. Short-term symptoms of uncontrolled diabetes include thirst, weariness, frequent urination and impaired eyesight [5,6,20]. Over an extended period, people may experience heart illness, kidney issues, visual impairments, nerve damage and further challenges. Insulin injections, diet, exercise and medication are used to introduce it. For most people, reducing weight and upping physical activity will help postpone or even prevent Type 2 diabetes. Diabetes that is not under control can cause kidney failure, stroke, heart disease and blindness among other health problems. On the other hand, with appropriate testing, treatment and lifestyle type 2 diabetics are controlled [5,6,19].

2.1.3 Gestational diabetes

Certain pregnant women develop gestational diabetes, which is the third kind of diabetes. This type of diabetes will have similar symptoms and therapy to that of Type 2 diabetes. It normally disappears after childbirth [6]. The quantity and kind of carbohydrates we eat during the day should be the main emphasis of our diet plan.

Table 1. Ingredients of diet in different foods

Foods	Ingredients of diet
Nuts	Nuts reduce insulin resistance, inflammation and cholesterol because they are high in unsaturated fats, proteins and a variety of vitamins and minerals. The investigation found that to lower high blood sugar and fat (triglycerides) we should consume at least 50 grams of almonds, cashews, chestnuts, walnuts or pistachios each day [6,9].
Banana	A study from University College Dublin in Ireland suggests that resistant starch which is present in foods like potatoes, bananas, grains, and legumes may improve gastrointestinal health, help regulate blood sugar levels and increase feelings of fullness. This type of starch is regarded as dietary fiber since it is not broken down in the small intestine [9,12].

Foods	Ingredients of diet
Amla	The Indian gooseberry tree yields amla which has long been used as a traditional treatment for high blood sugar. Additionally, it has a mineral called chromium which controls how carbohydrates are metabolized and improves our body's sensitivity to insulin [9].
Greens	Leafy greens are rich in minerals such as vitamin A and magnesium, as well as fiber. Blood sugar can be lowered with the use of these foods [9]. Consuming 1.35 portions in place of. A 14 percent lower incidence of type 2 diabetes is linked to eating two servings of leafy greens each day [16].
Garlic	Garlic may be able to assist with blood sugar regulation. Consuming garlic is said to reduce fasting blood glucose or blood sugar levels that occur when you haven't eaten. Studies of a similar nature indicate that onions may also benefit blood sugar levels [9,12].

However, it's also critical to eat things we enjoy. Want to eat enough to feel full and prevent overindulging and making bad decisions. These foods can help control blood sugar levels and improve overall health and happiness which is shown in Table 1.

3. INHIBITION OF DIABETES CONTROL

3.1 Insulin Treatment in Diabetes Mellitus

The introduction of insulin to treat diabetes has saved an estimated 5 million years of life for patients with type 1 diabetes during the year 2000 [12]. The manufacturing, formulation and administration of insulin preparations as well as the creation of insulin treatment regimens that sustain long-term normo-glycemia with minimal risk of hypoglycemia have advanced significantly in recent years [14,15]. In both type 1 and type 2 diabetes, the goal of preventing or delaying the development of chronic microvascular problems has been well demonstrated during the past ten years. Regrettably, glycemic control is consistently worse in patients using insulin than in people receiving other therapy [16,17]. It is widely acknowledged that the only significant factor limiting the dosage of insulin is hypoglycemia, making insulin the most effective glucose-lowering medication. In contrast to all oral medications which have a finite maximum impact, the dose of insulin causes increasingly more side effects. The goal of insulin therapy should be to emulate nature which has remarkable efficacy in preventing hypoglycemia in between meals and minimizing postprandial hyperglycemia [18]. However, pharmacological issues make insulin therapy more difficult and complex. First effective portal insulin concentration is only achieved at the cost of hyper-insulinemia in the systemic circulation because subcutaneous insulin injection sites drain into the peripheral circulation rather than the portal circulation. The second factor is the pharmacokinetic and pharmacodynamic characteristics of therapeutic insulin preparations

[19]. For instance, long-acting insulin preparations do not offer the consistent, low concentrations needed in between meals while short-acting insulins are absorbed too slowly and last too long to resemble the typical prandial peaks [19]. Furthermore, it is impossible to forecast how a given insulin dose would drop blood glucose because subcutaneous insulin absorption varies greatly. Once more the injection site is crucial; it should be closer to the abdomen than the thigh, inject more deeply, intramuscularly, warm the area beforehand or use local massage [10]. Some of the shortcomings have been addressed by the development of insulin analogues have an alternative part in the amino acid sequence of human insulin which changes the rate of insulin absorption or some other structural change like being linked to a fatty acid chain that alters the insulin time action curve [20]. Regular insulin is modified to result in the various short-acting insulin analogies: insulin lispro (Humalog), insulin aspart (Novolog) and insulin glulisine (Apidra); intermediate (Isophane, Lente) long-acting analogues: insulin glargine and insulin detemir [21].

Many insulin preparations are available and are grouped according to their duration of action: a rapid-acting formulation to cover meals and intermediate and longer-acting preparations to provide steady basal levels between meals and overnight. Insulin is prepared either from humans, porcine or bovine or a mixture of bovine and porcine in Table 2. Nowadays, human insulin (Humulin, Novolin) is produced using recombinant DNA technology and is widely accessible [21]. Because the amino acid sequences of human, porcine and bovine insulins differ, so do their physicochemical characteristics. In aqueous solutions, human insulin dissolves more frequently than pig insulin does. To increase its stability, it is administered at a pH of neutral. For almost all patients with type 1 diabetes and many with type 2 diabetes

insulin is the cornerstone of their care. Insulin can be injected intravenously (IV) or intramuscularly (IM) although the subcutaneous (SC) method is recommended for long-term therapy [21].

3.2 Short and Rapid-Acting Insulin Preparations

Exhibit the quickest initiation of effects but the briefest duration. It is recommended to administer short-acting insulin approximately 30 to 45 minutes before meals [21]. Regular insulin can also be administered intravenously or intramuscularly. Following intravenous (IV) administration blood glucose concentration highly decreases within 30-45 minutes (5-15 minutes for lispro, aspart and glulisine insulins) and reaches its maximum level between 1.5 to 4 hours (30-90 minutes for lispro, aspart and glulisine). The duration of its effect lasts for 5-8 hours (2-5 hours for lispro, aspart and glulisine) [22]. Intravenous administration of insulin is beneficial for patients experiencing ketoacidosis or undergoing surgery, labour and delivery or critical care. Regular insulin is found in a solution for injection in the form of a hexamer. To be effectively absorbed into the bloodstream the insulin hexamer must break apart into dimers or monomers [7]. The disintegration time in regular insulin is determined by the dissociation process which typically takes 30-60 minutes. This process also influences the beginning and the shape of the time-action curve. Contrary to ordinary insulin the insulin analogues (Lispro, Aspart and Glulisine) rapidly separate into individual units immediately after being injected. This feature results in fast absorption and a shorter duration of action compared to normal insulin. Lispro has two distinct benefits compared to ordinary insulin: firstly, it decreases the occurrence of hypoglycemia by 20% to 30% secondly, it moderately but considerably enhances glucose management as measured by haemoglobin A1c, by 0.3% to 0.5% [8]. Aspart insulin and Glulisine insulins share similarities with Lispro. Various inhalable preparations of quick-acting human regular insulin in the form of dry powder or liquid suspension are accessible. These preparations exhibit a similar initiation and peak action time as rapid-acting insulin but have a more prolonged duration of action compared to existing rapid-acting insulin analogues [10].

3.3 Intermediate-acting Insulin

Subcutaneous injection of intermediate-acting insulin preparations results in an onset of action

of around 1-2 hours, reaching its maximum effect at 6-12 hours, and lasting for 18-24 hours in Table 2. Intermediate-acting insulins are designed to break down more slowly when injected under the skin resulting in a significantly extended period of action [11]. The two most commonly utilized formulations are neutral protamine Hagedorn (NPH) insulin which is an isophane insulin suspension and lente insulin which is an insulin zinc suspension. NPH insulin is a solution of insulin that is combined with zinc and protamine in a phosphate buffer resulting in a suspension [11]. Lente insulin is a combination of ultralente and semilente insulins that are crystalline and amorphous respectively. It is formulated in an acetate buffer to reduce the solubility of insulin. NPH and lente insulins share comparable pharmacokinetic and pharmacodynamic characteristics in terms of their ability to reduce glucose levels. These insulins reach their maximum effect 4-6 hours after being injected subcutaneously and gradually diminish over 18-24 hours. Consequently, injecting these insulins either separately or in a premixed solution before the evening meal increases the risk of nocturnal hypoglycemia [23]. Therefore, it is recommended to provide intermediate-acting insulin preparations after bedtime to assist in stabilizing fasting blood glucose, especially in type 2 DM. The injections are either administered once (in the elderly) or twice a day. The implementation of intermediate-acting insulin formulations led to a decrease in the frequency of administering numerous injections of standard insulin. Regular and NPH insulin can be combined in a single syringe [23]. When NPH insulin is mixed with regular insulin it does not inhibit the action of regular insulin. In advanced research might be used the metal oxide nanoparticle used for this survey or trial study [23a-23c] In contrast, when lente insulin is mixed with regular insulin part of the lente insulin may form a complex with protamine or Zn^{2+} after several hours which delays the absorption of the fast-acting insulin [23].

3.4 Long-acting Insulin Preparations

Ultralente insulin also known as extended insulin zinc suspension and protamine zinc insulin suspension is classified as a long-acting insulin. They exhibit a delayed onset and a prolonged generally steady peak of action. Still, ultralente as a form of long-acting insulin is given in Table 2. While a certain amount of insulin remains in the body for 24 hours after injection it is

insufficient to adequately satisfy the basal or background insulin needs [24]. Ultralente insulin exhibits a comparable start and peak to intermediate-acting insulin however slightly delayed or extended. Ultralente has a duration that surpasses intermediate insulin but falls short of insulin glargine in terms of length. In most situations, it is taken twice in daily (early morning and at bedtime) or in a regimen where ultralente is provided at breakfast to supply basal insulin throughout the day and evening and NPH is given at bedtime to inhibit hepatic glucose production overnight. Nowadays this regimen of adding ultralente is less typically employed with other longacting insulin formulations accessible. Protamine zinc insulin is now obsolete due to its highly erratic and protracted duration of effect, and it is no longer accessible in the United States [24]. Insulin glargine (Lantus) is now the only available insulin that fulfils the majority of the criteria for long-acting insulin. Insulin glargine is produced via recombinant DNA technology which leads to alterations in the amino acid composition of conventional insulin giving it the designation of an insulin analogue. Glargine is a transparent solution with a pH value of 4.0 [25].

Its acidic pH means that it cannot be combined with the short-acting insulin preparations that are now on the market such as lispro or normal insulin which are made to have a neutral pH. When injected through the skin it has an extended flat activity profile lasting almost 24 hours and thus provides acceptable baseline levels when given once daily. In comparison to NPH insulin glargine administered at bedtime results in fewer overnight hypoglycemia episodes and a less fluctuating fasting glucose concentration in both adults and children [26]. It is recommended to provide rapid-acting insulin with the meal instead of dividing the insulin glargine into two injections since some individuals do not have adequate basal insulin coverage as shown by a rise in blood glucose shortly before the next glargine dose. In circumstances where glucose management is satisfactory but seems to have diminishing basal insulin coverage, then it is beneficial to split the insulin glargine. Insulin glargine is not licensed for use in pregnancy and there is inadequate data on pregnancy to provide a suggestion for clinical application at the same time [27].

Acrylated insulin is produced when a saturated fatty acid is added to an e amino acid. Insulin detemir is a readily available analogue of insulin.

Unlike conventional insulin, insulin detemir lacks threonine at position B30 and contains a fatty acyl chain connected at position B29 of the lysine amino acid [28,29]. The presence of a fatty acyl side chain in the substance causes a delay in the metabolism and elimination from the bloodstream following a subcutaneous injection. As a result, the duration of its effects is greater than that of the intermediate substance, but not as long as insulin glargine [30]. The clinical classification of insulin detemir as either a prolonged-intermediate-acting insulin which typically requires two injections per day and has a moderate peak or long-acting insulin which usually requires one injection per day with a minimal peak will ultimately depend on future clinical trials and clinical experience [30].

3.5 Premixed Insulin

Premixed insulin was created to enhance convenience and simplicity for patients when it comes to ordering and adjusting their dosage as well as to facilitate the process for physicians. Currently, premixed insulin is offered in various combinations of short-acting or rapid-acting insulin (soluble insulin) to intermediate-acting insulin (NPH or a protamine solution) [31,32]. The majority of insulin currently used consists of premixed formulations, primarily in ratios of 30/70 (soluble/NPH). However, these formulations are gradually being supplemented by newer combinations of rapid-acting analogues (such as lispro or aspart) with protamine-retarded counterparts [33]. Examples of these combinations include lispro/neutral protamine lispro in ratios of 25/75, 50/50 and 75/25 as well as aspart/neutral protamine aspart in ratios of 30/70, 50/50 and 70/30. Traditional premixed insulin formulations are administered twice a day before breakfast and the evening meal. Although widely favoured several experts argue that a predetermined ratio of short to intermediate insulin does not permit precise adjustment of each insulin component to match self-monitoring blood glucose (SMBG) patterns [34]. Currently, there have been no direct evaluations comparing the use of premixed insulin to the practice of individually mixing and modifying each insulin. Empirical evidence has demonstrated that premixed insulin is unsuitable for individuals with type 1 diabetes as it necessitates frequent modifications to attain optimal blood sugar regulation. Conversely, in the case of type 2 diabetes, the advantages of simplicity and the capacity to easily make adjustments for both patients and healthcare providers may be more

Table 2. Insulin preparation and its properties

Type	Properties (action in hours)			
	Onset	Peak	Duration	[31,32].
Regular soluble (Crystalline)	0.5-0.7	1.5-4	5-8	[31,32].
Lispro <i>Intermediate</i>	0.25	0.5-1.5	2-5	[31,32].
NPH (isophane)	1-2	6-12	18-24	[31,32].
Lente Slow	1-2	6-12	18-24	[31,32].
Ultralente	4-6	16-18	20-36	[31,32].
Protamine zinc	4-6	14-20	24-36	[31,32].
Glargine	2-5	5-24	18-24	[31,32].

important than the drawback of not being able to precisely alter the doses. Rapid-acting insulin analogues are progressively replacing normal insulin because of superior postprandial glucose management reduced hypoglycemia and more convenience for patients at the time of insulin injections [34].

4. COMPLICATIONS OF INSULIN THERAPY

The two most frequent side effects of insulin are hypoglycemia and weight gain. Inappropriate dose mismatches between insulin peak delivery and food intake or the addition of factors that increase insulin sensitivity (adrenal insufficiency, pituitary insufficiency) or that increase glucose uptake without insulin (exercise) can all result in hyperglycemia [35,36]. Episodic hypoglycemia episodes are more common the more vigorously euglycemia is attempted. In one clinical research (DCCT) the group receiving intensive insulin therapy experienced a three-fold increase in hypoglycemia responses compared to the group receiving conventional therapy. Hypoglycemia can occur less frequently and carries a lower risk when physiological insulin regimens are used in conjunction with education. Increased muscle fat and truncal mass result in weight gain which is an inevitable side effect of initiating insulin therapy for uncontrolled diabetes [37]. This is also because there are fewer energy losses from glycosuria. Instance physiological insulin regimes can limit weight gain by decreasing the demand for snacks in both adults and children due to improper insulinemia and hypoglycemia between meals. When insulin is initiated to treat type 2 diabetes metformin can help reduce weight gain [38-40], insulin allergy and resistance. Using human insulin or extremely pure forms of the hormone has resulted in a significant reduction in the prevalence of insulin resistance and allergic responses. Allergy responses were particularly common with bovine insulin [41]. The small amounts of aggregated or denatured insulin that

are present in all preparations, the tiny amounts of contaminants, or the sensitivity to any of the components that go into making insulin (protamine, Zn²⁺, phenol etc.) are still the causes of these reactions [42,43]. Allergy reactions mediated by IgE were the most common although at present they are very uncommon. Although there are benefits to skin testing, many people respond well to intradermal insulin without suffering any negative side effects. Desensitization is a recommended treatment if the allergy continues and it works for 50% of people [44,45].

5. SOME FRUITS FOR A DIABETES-FRIENDLY DIET

5.1 Berries from the Refreshing Treat and Disease-fighting Antioxidant

Feel free to indulge in your love of berries whether they're blueberries, strawberries or another kind. The American Diabetes Association claims that berries are a superfood for diabetics for their high fiber and antioxidant content. The U.S. Department of Agriculture (USDA) reports that a cup of fresh blueberries contains 84 calories and 21 grams (g) of carbs. Try layering berries with plain nonfat yogurt in a parfait if you can resist the impulse to just pop them straight into your mouth. It provides a terrific dessert or brunch for diabetes as shown in Fig. 1 [46,47].

5.2 Tart Cherries Help Fight Inflammation

According to the USDA, a cup of cherries contains 52 calories and 12.5 g of carbohydrates. Cherries may have anti-inflammatory properties. According to a review that was published in *Nutrients* in March 2018 tart cherries are also a great source of antioxidants which may help prevent heart disease, cancer and other illnesses. Fresh, canned, frozen or dried fruits

are available for purchase. To avoid a blood sugar surge, make sure to read the labels of

canned and dried fruits because many of them include added sugar [46,47].

1 Berries for a Refreshing Treat and Disease-Fighting Antioxidants



2 Tart Cherries Help Fight Inflammation



3 Sweet, Juicy Peaches for Metabolism-Boosting Potassium



4 Apricots for a Scrumptious, Fiber-Rich Bite



5 Apples for a Quick Fibrous and Vitamin C-Rich Snack



6 Oranges for a Juicy, Refreshing Source of Vitamin C



7 Pears for Easy Snacking, Plus Vitamin K and Fiber



8 Zesty Green Kiwi for Potassium, Fiber, and Vitamin C



Fig. 1. Fruits for a diabetes-friendly diet

5.3 Sweet Juicy Peaches for Metabolism-Boosting Potassium

Incorporating fragrant and luscious peaches into your diabetes-friendly diet is a great way to apply them during the warmer months. The USDA reports that one medium peach has 14 g of carbs and 59 calories. Furthermore, it contains 285 mg of potassium (six percent of the daily intake) and 10 mg of vitamin C or eleven percent of the DV for that nutrient. The fruit tastes great either way on its own or added to iced tea for a refreshing twist. Blend peach slices with low-fat buttermilk, crushed ice and a dash of ginger or cinnamon to make a simple smoothie that's perfect for those with diabetes [46,47].

5.4 Apricots for a Scrumptious Fibre-Rich Bite

A great way to incorporate apricots into your diabetic meal plan is to add them as a staple sweet summer fruit. According to the USDA one apricot contains just 17.0 calories and 4.0 g of carbs. 15.0% of your daily value (DV) of vitamin A or 134 micrograms (mcg) can be obtained from four fresh apricots. These juicy gems are a fantastic source of fiber as well. The dietary fiber content of four apricots is 3.0 g or 10.0% of the DV. Consider adding some chopped fresh apricots to a salad or incorporating them into hot or cold cereal [46,47].

5.5 Apples for Quick Fibrous and Vitamin C-Rich Snack

A daily apple could indeed fend off illnesses. As a terrific fruit option, medium-sized apples have only 95 calories and 25 grams of carbohydrates, so pack one in your purse or tote bag if you're on the run. In addition, apples have a good amount of fiber (4.0 g per medium fruit or 16 percent of your DV) and vitamin C (8.73 mg, or roughly 9.0 percent of the DV) in a midsize apple [46,47]. Apple skins are nutrient-dense with additional fiber and heart-protective antioxidants; nevertheless, avoid peeling them, advises the Harvard T.H. Chan School of Public Health [46,47].

5.6 Orange for a Juicy Refreshing Source of Vitamin C

One medium orange has 70.0 milligrams of vitamin C, which equates to 78.0 percent of your daily requirement. According to the USDA, this light option has just 62.0 calories and 15.0 g of

carbs. Additionally, one medium orange has 237.0 mg of potassium (5.0 % of the DV) and 40.0 mcg of folate (10.0 percent of the DV), both of which may help stabilize blood pressure [46,47]. While you're savoring this delicious delight, keep in mind that grapefruit along with other citrus fruits is also an excellent option [46,47].

5.7 Pears for Easy Snacking Plus Vitamin K

Pears are a smart addition to your diabetic meal plan since they are a great source of fiber (one medium fruit has around 5.5 g or 20.0 percent of the DV) according to the USDA. Moreover, their texture and flavour enhance after picking, in contrast to most other fruits [46,47]. According to USA Pears keep your pears at room temperature until they are mature and ready to eat (at which point you can store them in the refrigerator). Here's a delightful treat, cut a pear into slices and add them to the next green salad [46,47].

5.8 Zesty GREEN Kiwi for Potassium

A kiwi's zesty bright green fruit is hidden beneath its fuzzy brown peel which you may not be aware of if you haven't had one. One tasty, nutritious kiwi contains 5.0 percent of the daily value (DV) of potassium (215.0 mg), 71.0 percent of the daily value (64.0 mg) of vitamin C and 8.0 percent of the daily value (g) of fiber [46,47]. A single kiwi provides approximately 42.0 calories and 10.0 g of carbs, making it a wise inclusion in your diet for those with diabetes. Zespri Kiwifruit claims that kiwis are accessible all year round and can be stored in the fridge for as long as seven days [46,47].

6. DIABETIC MEAL PLAN

Anyway, it is anticipated that a balanced diet with lots of fruits and vegetables and less processed food will lower the chance of developing several chronic degenerative diseases. Furthermore, eating a diet high in whole foods facilitates better hunger control and typically keeps people from becoming overweight or obese. Conversely, a fast-food diet raises the risk of type II diabetes and obesity [48]. Meal plans for diabetics balance the calories from foods (carbs, proteins, and lipids) with each person's insulin dosage and amount of bodily activity. Foods including milk, fruits, vegetables, legumes, bread and grain, meat and fat are frequently consumed. Proteins, lipids and carbs are all present in these foods

[49]. A meal plan should be designed so that 50–60% of daily calories come from carbohydrates, 15.0% through proteins, and the remaining portion is derived from lipids. Maintaining a regular daily schedule is crucial if insulin treatment is taken. The design of the plan will balance the peaks of insulin and carbohydrates. The World Health Organization (WHO) suggests consuming at least five to ten servings of fruits and vegetables each day [49].

6.1 Role of Fruits in Diabetes

An essential component of any person's basic diet should be fruits. Fruits supply the body with fiber, vitamins, and minerals that are necessary for a full, balanced diet, regardless of whether a person has type 2 diabetes or not. People with type 2 diabetes can still benefit from fruits but only if they avoid ones with high sugar and carbohydrate content which might raise blood glucose levels [50]. To manage diabetes 2, they ought to have fruits with a low Glycemic Index. Cherries, prunes, grapefruit, dried apricots, raisins, peaches, apples, pears, strawberries, plums, guava, orange, grapes, papaya, banana, kiwi, pineapple, figs and mango are among the fruits that have a low glycemic index. Despite watermelon having a high Glycemic Index, the glycemic load per food serving (120.0 g) is minimal, meaning it will not significantly impact blood glucose levels. Consequently, a standard portion, due to its higher water content, has few carbohydrates, resulting in a low glycemic load [50]. It is advisable to consume it in limited quantities. The consumption of these items is limited to their raw or cooked form or as juice,

without any additional sugar. When purchasing fruits, consider acquiring them in smaller portions. Avoid fruit juices and choose whole fruits instead as they have more fiber content and provide a greater feeling of satiety. When purchasing canned fruit juices, it is essential to examine the label for the terms 'unsweetened extra-light or no sugar added, if necessary. Avoid consuming fruits that have been preserved by freezing or canned in a dense syrup [51].

6.2 General Dietary Guideline

The contemporary approach to managing diabetes mostly entails making adjustments to the type and amount of food consumed by individuals with diabetes. Any form of diabetes regardless of weight status, age, gender or career is covered by the following rules as shown in Table 3 [52-61].

Weight loss in overweight diabetes patients through reduction in portion sizes and calorie intake leads to improved insulin sensitivity hence optimizing the effectiveness of medication therapy. The energy balance is the basic idea underlying maintaining body weight [62,63]. This demographic should be motivated to sustain their present weight by adhering to consistent portion sizes, consuming a similar quantity of food daily eating at consistent intervals throughout the day adhering to a consistent medication schedule and engaging in regular exercise at a consistent interval. These patients should strive to select their daily meals from sources of carbs, vegetables, fruits and protein while restricting the use of fats [64-66].

Table 3. General dietary guidelines for foods

SI No.	Guideline	
1	Most of the carbohydrates consumed should be in the form of starch (polysaccharides) such as maize, rice, beans, bread, potatoes etc.	[52, 61].
2	All refined sugars such as glucose, sucrose and their products (soft drinks, sweets, toffees, etc.) and honey should be avoided, except during severe illness or episodes of hypoglycemia. These foods contain simple sugar, which is easily absorbed causing a rapid rise in blood sugar.	[53, 62].
3	Non-nutritive sweeteners, e.g., Canderel, saccharine, NutraSweet and aspartame are suitable sugar substitutes for diabetic subjects.	[54, 59].
4	Animal fats such as butter, lard, egg yolk and other foods high in saturated fatty acids and cholesterol should be reduced to a minimum and replaced with vegetable oils, particularly polyunsaturated fats.	[55, 63].
5	Salt should be reduced whether hypertensive or not.	[56, 60].
6	Protein (fish, meat, beans, crab, crayfish, soybean, chicken, etc.) and salt are restricted for those with diabetic nephropathy.	[57].
7	Cigarette smoking should be avoided by diabetic patients. Alcohol should be taken only in moderation.	[58, 64].

SI No.	Guideline	
8	The items allowed for free consumption include Water, green leafy vegetables, tomatoes, onions, cucumber, aubergine, peppers and vegetable salad without cream. Any brand of tea, coffee or drinks that contain very low or no calories.	[59].
9	For patients too ill to eat solid food a fluid or semi-solid diet should be substituted (papaya, soya bean, custard, etc.).	[52, 61].
10	Patients treated with insulin or certain oral hypoglycemic agents, e.g., sulfonylureas must be advised to eat regularly and often to prevent hypoglycemia 3 meals a day plus suitable snacks in between e.g., fresh fruits.	[60, 65].
11	Small meals spaced over the day rather than 1 or 2 big meals help avoid post-pyramidal peaks in blood sugar.	[61, 66].

6.3 Dietary Approach to Diabetes

A healthy weight and normal growth and development particularly during pregnancy and lactation are all supported by dietary treatment for diabetes which aims to attain ideal blood sugar levels and cholesterol concentrations as well as adequate calories. By addressing diabetes-related problems and preventing delays a balanced diet can enhance overall health in Table 4 [67-71]. Additionally, the patients have to make an effort to choose meals that are high in fruits, vegetables, carbohydrates and proteins and low in fats.

Food has a significant role in both preventing and treating diabetes. However, as our understanding of the illness has grown, so too have nutritional strategies [67-68]. The conventional treatment of diabetes centres on reducing refined sugar intake and meals which release sugars upon digestion such as starches, bread, fruits and other foods that have high fat and protein content and low carbohydrate content [69]. Thus, diabetic specialists have been careful to restrict protein for individuals with compromised kidney function as well as lipids, particularly saturated fats that might increase cholesterol [70]. The revised strategy emphasizes fat more. Diabetes sufferers have issues with fat. Insulin's ability to transport

glucose into cells will be hampered by a diet high in fat. On the other hand, cutting down on fat consumption and body fat improves insulin's function significantly. More recent treatment plans significantly cut back on meats, oils and high-fat dairy goods [71]. They also boost the number of fruits, vegetables, grains and legumes. Consume a diet rich in fruits and vegetables daily. Fruit juices are never as good as the actual fruit. Water is a better option than typical sodas and fruit drinks; low-calorie juices, tea and coffee are also excellent options. Consume lower GI meals, which will benefit your diabetes by preventing blood sugar spikes and maintaining control over your blood sugar levels [72]. Diabetes can be effectively prevented and reversed with food. Still, as our understanding of the illness has grown so too have nutritional strategies. Reducing refined sugar intake and meals like baked goods, fruits and carbohydrates that release sugar after digestion is the mainstay of conventional diabetic treatment. Reduced carbs could mean an unhealthy proportion of fat and protein in the diet. Thus, diabetic specialists have been careful to restrict protein for individuals with compromised kidney function as well as lipids, particularly saturated fats which might raise cholesterol [73-74]. The revised strategy emphasizes fat more and the sufferer's issues with fat. Insulin has a tougher job

Table 4. Main food categories: include unlimited amounts of grains, legumes, fruits and vegetable

Food categories	Sources	
Grains	Rice, high-fiber cereals, corn, oatmeal, wheat, millet, barley, rye.	[67-71].
Legumes	Beans, peas, lentils, nonfat soy products.	[67-71].
Fruits	All, except avocados, olives, pineapple and watermelon, bananas, apples, grapes, pears, peaches, oranges, melon, grapefruits, kiwi and barriers among others all a good choice.	[67-71].
Vegetables	All, except white potatoes. They include tomatoes, cucumbers, carrots, broccoli, cauliflowers, spinach, kale, cabbage, green, beans, sweet potatoes and artichokes.	[67-71].

delivering glucose into cells with the higher fat content of a diet. On the other hand, cutting down on fat consumption and body fat improves insulin's function significantly. More recent treatment plans significantly cut back on meats, oils, and high-fat dairy goods. Grain, legume, fruit, and vegetable intake all rise together. According to the study, patients who were taking insulin or oral drugs were able to stop taking them after a few days if they followed an almost vegan dietary and physical activity regimen [75]. Diet and lifestyle guidelines evidence-based ayurvedic perspective for diabetes remedy [76,77,78], vegetable oil [79] and banana [80] also introduced. The majority of diabetic patients treated using this regimen maintained their improvements after two and three-year follow-ups. These small but effective dietary adjustments are effective [81].

7. CONCLUSION

Successful lifestyle changes such as weight loss counselling, adopting a nutritious diet just like the Mediterranean diet and increasing physical activity are the mainstays in the fight against diabetes. So, especially for those who are at high risk, the focus needs to be placed on encouraging a better lifestyle and figuring out ways to improve adherence as well as compliance with lifestyle changes and diet. The Mediterranean food pattern has a preventive effect against diabetes, according to findings from clinical trials and epidemiological research assessing the pattern's impact on the onset and control of management of the disease. An essential part of caring for a diabetic patient is diet. Both the patient and the diabetic healthcare professional should be aware of the patient's fundamental dietary requirements. Even if the blood contains an abundance of insulin in this form, the cells are not receptive to it. Because it is difficult for glucose to enter cells it builds up in the blood vessel. Short-term symptoms of uncontrolled diabetes include thirst, weariness, frequent urination and impaired eyesight. Over an extended period, people may experience heart illness, kidney issues, visual impairments, nerve damage and further challenges. This review indicates that there is a notable variation in the connections between specific fruits and the likelihood of developing type 2 diabetes. Increased intake of some whole fruits specifically blueberries, grapes, and apples, was strongly linked to a reduced risk of developing type 2 diabetes. Conversely, higher consumption of fruit juice was connected with an elevated risk.

ACKNOWLEDGEMENT

We apologize to other authors who have worked on several aspects of Diabetes Mellitus but whom we have unintentionally overlooked. We have cited as many references as permitted and apologize to the authors of those publications that we have not cited due to limitation of references. Words are inadequate to express my heartiest obligation and gratitude to *Dr. Md. Abdus Sattar*, Professor, Department of Applied Chemistry & Chemical Engineering (ACCE), Kushtia-7003 for his keen interest, proficient suggestions, sympathetic guidance and persistent encouragement. We feel great honour and pleasure in expressing thanks to the Chairman, Department of Applied Chemistry & Chemical Engineering (ACCE), Kushtia-7003 for his guidance and kind behaviour during my Project.

COMPETING INTERESTS

The authors have declared that no competing interests exist.

REFERENCES

1. American Diabetes Association. Diabetes e-newsletter er; Sat. 2 August 2003.
2. American Diabetes Association. Screening for diabetes. *Diabetes Care* 2002;25:21-4.
3. Karjalainen J, Martin JM, Knip M, Ilonen J, Robinson BH, Savilahti E, et Asif.: Role of food additives in diabetes on Tuesday, [December 8, 2020, IP: 118.179.37.161] *International Journal of Nutrition, Pharmacology, Neurological Diseases*. January 2011;1(1):35.
A bovine albumin peptide is a possible trigger of insulin-dependent diabetes mellitus. *N Engl J Med* 1992;327:302-7.
Available:<http://www.ijnpnd.com>
4. Peterson KF, Dufour S, Befroy D, Garcia R, Shulman GI. Impaired mitochondrial activity in the insulin-resistant offspring of patients with type 2 diabetes. *N Engl J Med*. 2004;350:554-671.
5. Barnard RJ, Massey MR, Cherny S, O'Brien LT, Pritikin N. Longterm use of a high complex carbohydrate, high-fiber, low-fat diet and exercise in the treatment of NIDDM patients. *Diabetes Care*. 1983;6: 268-73.
6. Mann JI. Diabetes mellitus: Nutritional aspects of etiology and management. *Recent Adv Clin Nutr* .1986;2:255-70.

7. Lepore M, Pampanelli S, Fanelli CG, et al. Pharmacokinetics and pharmacodynamics of the long-acting human insulin analogue glargine, NPH insulin, and ultralente human insulin injected subcutaneously, and continuous subcutaneous infusion of insulin lispro. *Diabetes* 2000;49:2142-2148.
8. Riddle MC. New tactics for type 2 diabetes: Regimens based on intermediate-acting insulin taken at bedtime. *Lancet*. 1985;1: 192-195.
9. Available: <https://www.daily-bangladesh.com/english/Best-Foods-to-Control-Diabetes/5634>
Health & Lifestyle Desk Daily-Bangladesh Published: 04:07 PM, 7 May 2018.
10. Davis SN, Thompson CJ, Brown MD, et al. A comparison of the pharmacokinetics and metabolic effects of human regular and NPH insulin mixtures. *Diabetes Res Clin Pract*. 1991;13:107-117.
11. Colagiuri S, Villalobos S. Assessing effect of mixed insulins by glucose-clamp technique in subjects with diabetes mellitus. *Diabetes Care*. 1986;9:579-586.
12. Owens DR, Zinman B, Bolli GB. Insulins today and beyond. *Lancet*. 2001;358:739-746.
13. Owens DR, Barnett A, Designer insulins: have they revolutionized insulin therapy? In: Betteridge DJ, ed. *Diabetes: Current Perspectives*. London: Dunitz. 1999;223-267.
14. Bolli GB. Hypoglycemia unawareness. *Diabetes Metab*. 1997;23(suppl 3):29-35.
15. Bolli GB. How to ameliorate the problem of hypoglycemia in intensive as well as non-intensive treatment of type 1 diabetes mellitus. *Diabetes Care*. 1999;22 (suppl 2):B43-B52.
- 15a. Islam MR, Ahmed S, Sadia SI, Sarkar AK, Alam MA. Comprehensive Review of Phytochemical Content and Applications from *Cestrum nocturnum*: A Comparative Analysis of Physicochemical Aspects. *Asian Journal of Research in Biochemistry*. 2023;13(4):43-58.
- 15b. Haque NN, Alam MA, Roy CK, Zenat M, Akther E, Munshi JL. Cyanobacteria Mediated CO₂ Segregation: A Promising Alternative Method for Sustainable Bioremediation and Biomass Production. *Asian Journal of Research in Biochemistry*. 2023;13(3):28-43.
- 15c. Bhardwaj K, Najda A, Sharma R, Nurzyńska-Wierdak R, Dhanjal DS, et al. Fruit and vegetable peel-enriched functional foods: potential avenues and health perspectives. *Evidence-Based Complementary and Alternative Medicine*; 2022.
- 15d. Sharma R, Kakodkar P, Kabra A, Prajapati PK. Golden ager Chyawanprash with meager evidential base from human clinical trials. *Evidence-Based Complementary and Alternative Medicine*. 2022;1-6.
- 15e. Sharma R, Martins N, Chaudhary A, Garg N, Sharma V, Kuca K, et al. Adjunct use of honey in diabetes mellitus: A consensus or conundrum?. *Trends in Food Science & Technology*. 2020;106:254-274.
16. Diabetes Control and Complication Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Eng J Med*. 1993;329:977-986.
17. Ciofeta M, Lalli C, Del Sindaco P, et al. Contribution of postprandial versus antepandial blood glucose to HbA_{1c} in type I diabetes on physiologic intensive therapy with lispro insulin at mealtime. *Diabetes Care*. 1999;22:795-800.
18. Binder C, Lauritzen T, Faber O, Pramming S. Insulin pharmacokinetics. *Diabetes Care*. 1984;7:188-199.
19. Bethel MA, Feinglos MN. Insulin analogues: New therapies for type 2 diabetes mellitus (review). *Curr Diab Rep*. 2002;2:403-408.
20. Dimitriadis GD, Gerich JE. Importance of timing of pre-prandial subcutaneous insulin administration in the management of diabetes mellitus. *Diabetes Care*. 1983;6:374-377.
21. Bolli GB, Marchi RD, Park GD, et al. Insulin analogues and their potential in the management of diabetes mellitus. *Diabetologia*. 1999;42:1151-1167.
22. Davis SN, Granner DK. Insulin, oral hypoglycaemic agents, and the pharmacology of the endocrine pancreas. In: *The Pharmacological Basis of Therapeutics* by Goodman & Gilman. 10th Ed., Chapter 61, Edition: McGraw Hills Medical Publishing Division, USA. 2001; 1679-1714.
23. Dreyer M, Pein M, Schmidt C, et al. Comparison of the pharmacokinetics/dynamics of the Gly (A₂)-Arg (B₃₁, B₃₂)-human insulin (HOE71GT) with NPHinsulin following

- subcutaneous injection by using euglycaemic clamp technique. *Diabetologia*. 1994;37(Suppl):A78.
- 23a. Alam MA, Bishwas RK, Mostofa S, Jahan SA. Low-temperature synthesis and crystal growth behavior of nanocrystal anatase-TiO₂. *Materials Letters*. 2024;354:135396.
- 23b. Tabassum M, Alam MA, Mostofa S, Bishwas RK, Sarkar D, Jahan SA. Synthesis and crystallinity integration of copper nanoparticles by reaction medium. *Journal of Crystal Growth*. 2024;626:127486.
- 23c. Alam MA, Tabassum M, Mostofa S, Bishwas RK, Sarkar D, Jahan SA. The effect of precursor concentration on the crystallinity synchronization of synthesized copper nanoparticles. *Journal of Crystal Growth*. 2023;621:127386.
24. Pieber TR, Eugine-Jolchine I, Derobert E. The European Study Group of HOE901 in type 1 diabetes: Efficacy and safety of HOE901 versus NPH insulin in patients with type 1 diabetes. *Diabetes*. 2000;23:157-162.
25. Mohn A, Strang S, Wernicke-Panteau J, et al. Nocturnal glucose control and free insulin levels in children with type 1 diabetes by use of the long-acting HOE901 as part of a three-injection regimen. *Diabetes Care*. 2000;23:557-559.
26. Henry RR, Gumbiner BN, Ditzler T, et al. Intensive conventional insulin therapy for type II Diabetes. Metabolic effects during 6-month outpatient trial. *Diabetes Care*. 1993;16:21-31.
27. Kudlacek S, Schernthaner G. The effect of insulin treatment on HbA1c, body weight and lipids in type 2 diabetic patients with secondary failure to sulfonylureas. A five-year follow-up study. *Horm Metab Res*. 1992;24:478-483.
28. Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Eng J Med*. 1993;329:977-986.
29. Pampanelli S, Porcellati F, Fanelli C, et al. Glycaemic control, awareness of and counter-regulation to hypoglycemia, in the second decade of intensive treatment of T1DM (abstract). *Diabetes*. 2002; 51(Suppl2):A492.
30. Bott S, Bott U, Berger M, Mühlhauser I. Intensified insulin therapy and the risk of hypoglycemia. *Diabetologia*. 1997;40:926-932.
31. Lalli C, Ciofette M, Del Sindaco, et al. Long-term intensive treatment of type 1 diabetes with the shortacting insulin analogue lispro in variable combination with NPH insulin at mealtime. *Diabetes Care*. 1999;22:468-472.
32. Cryer PE. Iatrogenic hypoglycemia as a cause of hypoglycemia-associated autonomic failure in IDDM. *Diabetes*. 2002;41:255-260.
33. Diabetes Control and Complications Trial Research Group. The effects of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Eng J Med*. 1993;329:977-986.
34. Henry RR, Gumbiner BN, Ditzler T, et al. Intensive conventional insulin therapy for type II Diabetes. Metabolic effects during 6-month outpatient trial. *Diabetes Care*. 1993;16:21-36.
35. Kudlacek S, Schernthaner G. The effect of insulin treatment on HbA1c, body weight and lipids in type 2 diabetic patients with secondary failure to sulfonylureas. A five-year follow-up study. *Horm Metab Res*. 1992;24:478-483.
36. Diabetes Control and Complications Trial Research Group. The effect of intensive treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Eng J Med*. 1993;329:977-986.
37. Pampanelli S, Porcellati F, Fanelli C, et al. Glycaemic control, awareness of and counter-regulation to hypoglycaemia, in the second decade of intensive treatment of T1DM (abstract). *Diabetes*. 2002; 51(Suppl2):A492.
38. Bott S, Bott U, Berger M, Mühlhauser I. Intensified insulin therapy and the risk of hypoglycaemia. *Diabetologia*. 1997;40: 926-932.
39. Lalli C, Ciofette M, Del Sindaco, et al. Long-term intensive treatment of type 1 diabetes with the shortacting insulin analogue lispro in variable combination with NPH insulin at mealtime. *Diabetes Care*. 1999;22:468-472.
40. Cryer PE. Iatrogenic hypoglycaemia as a cause of hypoglycaemia-associated autonomic failure in IDDM. *Diabetes*. 2002;41:255-260.
41. Diabetes Control and Complications Trial Research Group. The effects of intensive

- treatment of diabetes on the development and progression of long-term complications in insulin-dependent diabetes mellitus. *N Engl J Med.* 1993;329:977-986.
42. Yki-Jarvinen H, Ryysy L, Nikkilä K, Tulokas T, Vanamo R, Heikkilä M. Comparison of bedtime insulin regimens in patients with type 2 diabetes mellitus: a randomized trial. *Ann Intern Med.* 1999;130:289-396.
 43. Kahn CR, Rosenthal AS. Immunology reaction to insulin: insulin allergy, insulin resistance, and autoimmune insulin syndrome. *Diabetes Care.* 1979;2:283-295.
 44. Yki-Jarvinen H, Ryysy L, Nikkilä K, Tulokas T, Vanamo R, Heikkilä M. Comparison of bedtime insulin regimens in patients with type 2 diabetes mellitus: a randomized trial. *Ann Intern Med.* 1999;130:289-396.
 45. Kahn CR, Rosenthal AS. Immunology reaction to insulin: Insulin allergy, insulin resistance, and autoimmune insulin syndrome. *Diabetes Care.* 1979;2:283-295.
 46. Maria Masters, Contributing Health Writer, Bachelor of Arts (B.A.) from the University of Pittsburgh. Maria Masters is a contributing editor and writer for *Everyday Health*. In addition to *Everyday Health*, Masters is a contributing editor and writer for *What to Expect* and has held positions at *Men's Health* and *Family Circle*. Her work has appeared in *Health*, *Prevention.com*, *Men'sJournal.com* and *HGTV* magazine, among numerous other print and digital publications.
 47. Kelly Kennedy, RD; *Everyday Health* Staff Kelly is a registered dietitian who has both a bachelor's degree in nutrition from the University of Massachusetts in Amherst and a master's degree in nutrition from the State University of New York College at Oneonta. She is also a licensed dietitian and nutritionist in the state of Massachusetts. Kelly manages and oversees nutrition content, meal planning, and diet and nutrition coaching at *Everyday Health*. In her spare time, Kelly enjoys anything that takes her outside — from gardening and playing in the yard with her kids to going for hikes and even feeding her pet chickens.
 48. Nitithan S, Komindr S, Nichachotsalid A. Phytate and fiber content in Thai fruits commonly consumed by diabetic patients. *J Med Assoc Thai.* 2004;87:1444-6.
 49. Fadupin GT, Keshinro OO. Factors influencing dietary compliance and glycaemic control in adult diabetic patients in Nigeria. *Diabetes Int.* 2001;11:59-61.
 50. Fadupin GT, Keshinro OO, Sule ON. Dietary recommendations: Example of advice given to diabetic patients in Nigeria. *Diabetes Int.* 2000;10:68-70.
 51. Prevention and management of the global epidemic of obesity. Report of the WHO consultation on obesity. Geneva: WHO; 1998.
 52. Hutch M. Best fruit and vegetables for type-2 diabetes; April 23, 2010.
 53. Barnard ND, Cohen, J, Jenkins DJ, Turner-McGrievy G, Gloede L, Jaster B, et al. A low-fat, vegan diet improves glycemic control and cardiovascular risk factors in a randomized clinical trial in individuals with type 2 diabetes. *Diabetes Care.* 2006;29:1777-83.
 54. Puwastien P, Valaipatchara U, Kongkachuichai R. Dietary fiber content in common Thai foods. *J Nutr Assoc Thailand.* 1990;24:43-53.
 55. The International Expert Committee. International Expert Committee report on the role of the A1C assay in the diagnosis of diabetes. *Diabetes Care.* 2009;32:1327–34. [PMC free article] [PubMed] [Google Scholar]
 56. Sinitskaya N, Gourmelen S, Schuster-Klein C, Guardiola-Lemaitre B, Pevet P, Challet E. Increasing fat-to-carbohydrate ratio in a high-fat diet prevents the development of obesity but not a prediabetic state in rats. *Clin Sci.* 2007;113:417–25. [PubMed] [Google Scholar]
 57. Sjostrom CD, Peltonen M, Wedel H, Sjostrom L. Differentiated long-term effects of intentional weight loss on diabetes and hypertension. *Hypertension.* 2000;36:20–5. [PubMed] [Google Scholar]
 58. Schroder H. Protective mechanisms of the Mediterranean diet in obesity and type 2 diabetes. *J Nutr Biochem.* 2007;18:149–60. [PubMed] [Google Scholar]
 59. Selvin E, Crainiceanu CM, Brancati FL, Coresh J. Short-term variability in measures of glycemia and implications for the classification of diabetes. *Arch Intern Med.* 2007;167:1545–51. [PubMed] [Google Scholar]
 60. Shai I, Schwarzfuchs D, Henkin Y, Shahar DR, Witkow S, Greenberg I, et al. Weight loss with a low-carbohydrate, Mediterranean, or low-fat diet. *N Engl J*

- Med. 2008;359:229–41. [PubMed] [Google Scholar]
61. Lindeberg S, Jönsson T, Granfeldt Y, Borgstrand E, Soffman J, Sjöström K, et al. A Palaeolithic diet improves glucose tolerance more than a Mediterranean-like diet in individuals with ischaemic heart disease. *Diabetologia*. 2007;50:1795–807. [PubMed] [Google Scholar]
 62. Lovejoy JC, Windhauser MM, Rood JC, de la Bretonne JA. Effect of a controlled high-fat versus low-fat diet on insulin sensitivity and leptin levels in African-American and Caucasian women. *Metabolism*. 1998;47:1520–4. [PubMed] [Google Scholar]
 63. American Diabetes Association. Clinical practice recommendations-2007. *Diabetes Management in Correctional Institutions: Diabetes Care*. 2007;30(Suppl 1):S77–84. [PubMed] [Google Scholar]
 64. American Diabetes Association. Diagnosis and classification of diabetes mellitus. *Diabetes Care*. 2010;33(Suppl 1):S62–9. [PMC free article] [PubMed] [Google Scholar]
 65. Bouchard C, Tremblay A, LeBlanc C, Lortie G, Savard R, Theriault G. A method to assess energy expenditure in children and adults. *Am J Clin Nutr*. 1983;37:461–7. [PubMed] [Google Scholar]
 66. Drewnowski A. Energy density, palatability, and satiety: Implications for weight control. *Nutr Rev*. 1998; 56:347–53. [PubMed] [Google Scholar]
 67. Duncan KH, Bacon JA, Weinsier RL. The effects of high and low energy density diets on satiety, energy intake, and eating time of obese and nonobese subjects. *Am J Clin Nutr*. 1983;37:763–7. [PubMed] [Google Scholar]
 68. Eaton SB, Eaton SB, Konner MJ, Shostak M. An evolutionary perspective enhances understanding of human nutritional requirements. *J Nutr*. 1996;126:1732–40. [PubMed] [Google Scholar]
 69. Esposito K, Maiorino MI, Di Palo C, Giugliano D. Adherence to a Mediterranean diet and glycaemic control in Type 2 diabetes mellitus. *Diabet Med*. 2009;26:900–7. [PubMed] [G46. Franz MJ, Splett PL, Monk A, Barry B, McClain K, Weaver T, et al. Cost-effectiveness of medical nutrition therapy provided by dietitians for persons with non-insulin-dependent diabetes mellitus. *J Am Diet Assoc*. 1995;95:1018–24. [PubMed] [Google Scholar]
 70. Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma without the use of the preparative ultracentrifuge. *Clin Chem*. 1972;18:499–502. [PubMed] [Google Scholar]
 71. Garg A, Bantle JP, Henry RR, Coulston AM, Griver KA, Raatz SK, et al. Effects of varying carbohydrate content of the diet in patients with non-insulin-dependent diabetes mellitus. *JAMA*. 1994;271:1421–8. [PubMed] [Google Scholar]
 72. Garg A. High-monosaturated fat diets for patients with diabetes mellitus: A meta-analysis. *Am J Clin Nutr*. 1998;67:577S–82. [PubMed] [Google Scholar]
 73. Howarth NC, Saltzman E, Roberts SB. Dietary fiber and weight regulation (Review) *Nutr Rev*. 2001;59:129–39. [PubMed] [Google Scholar]
 74. Pérez-Jiménez F, López-Miranda J, Pinillos MD, Gómez P, Paz-Rojas E, Montilla P, et al. A Mediterranean and a high-carbohydrate diet improves glucose metabolism in healthy young persons. *Diabetologia*. 2001;44:2038–43. [PubMed] [Google Scholar]
 75. Schakel SF, Sievert YA, Buzzard IM. Sources of data for developing and maintaining a nutrient database. *J Am Diet Assoc*. 1988;88:1268–71. [PubMed] [Google Scholar]
 76. Sharma R, Amin H, Ruknuddin G, Prajapati PK. Efficacy of Ayurvedic remedies in type 2 diabetes: A review through works done at Gujarat Ayurved University, Jamnagar. *Journal of Medical Nutrition and Nutraceuticals*. 2015;4(2): 63.
 77. Sharma R, Prajapati PK. Diet and lifestyle guidelines for diabetes: Evidence based ayurvedic perspective. *Romanian Journal of Diabetes Nutrition and Metabolic Diseases*. 2014;21(4):335-346.
 78. Sharma R, Bolleddu R, Maji JK, Ruknuddin G, Prajapati PK. *In-Vitro* α -amylase, α -glucosidase inhibitory activities and *in-vivo* anti-hyperglycemic potential of different dosage forms of guduchi (*Tinospora cordifolia* [willd.] miers) prepared with ayurvedic bhavana process. *Frontiers in Pharmacology*. 2021;12: 642300.
 79. Hasan M, Jahan R, Alam MA, Khatun MK, Al-Reza SM. Study on physicochemical properties of edible oils available in Bangladeshi local market. *Archives of*

- Current Research International. 2016;6(1): 1-6.
80. Shruthi D. Medicinal uses of banana (*Musa paradisiaca*). Drug Invention Today. 2019; 12(1).
81. Simopoulos AP. The Mediterranean diet: What is so special about the diet of Greece? The scientific evidence. J Nutr. 2001;131:3065S–73.

© 2024 Ali et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/111953>