



Pharmacological and Therapeutic Features of COVID - 19 Infection in Hodeidah, Yemen

**Mohammed A mood AL-Kamarany ^{a*},
Hussien Omer Alkadi ^b, Mohammed Yahia Almadani ^b,
Zainulbedeen Ali Alsharma ^c,
Rashad Mohammed Al-Salehy ^c
and Nabil Ahmed Albaser ^d**

^a *Pharmacy Practice Department, College of Clinical Pharmacy, Center of Tropical Medicine and Epidemiology Studies - Hodeidah University (CTMES HU), Yemen.*

^b *Department of Pharmacology and Therapeutics, Faculty of Medicine and Health Sciences, Sana'a University, Sana'a, Yemen.*

^c *Center of Tropical Medicine and Infectious Diseases (CTMID), Al-Thawra Public Hospital Authority, Hodeidah, Yemen.*

^d *Department of Pharmacology, Faculty of Medical Sciences, AL Razi University, Sana'a, Yemen.*

Authors' contributions

This work was carried out in collaboration among all authors. Author MAA contributed in analyzing of data, writing, revising and submitting of article. HOA supervised and revised the article. Author MYA collected the clinical data, authors ZAA and RMA contributed in clinical practice and Author NAA wrote the article. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMPS/2024/v26i5683

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/111318>

Original Research Article

Received: 07/11/2023

Accepted: 13/01/2024

Published: 23/03/2024

*Corresponding author: E-mail: alkamarany@gmail.com;

ABSTRACT

Background: The symptoms and signs of coronavirus diseases 2019 (COVID-19) present at illness onset vary, but over the course of the disease, most persons with COVID-19 will experience the following: fever, cough, fatigue, anorexia, shortness of breath, hypoxia, sputum production, and myalgia. During the first year (2020), there was no effective cure for COVID-19 infection and the most common treatment for patients was supportive therapy. Ministry of Public Health and Population in Yemen presented national guideline that is extracted from other countries and World Health Organization (WHO).

Objective: Therefore, in this research, we discuss the pharmacological and therapeutic approaches for management of COVID-19 during its early period of rampage, using availability of medicines in Yemeni local pharmaceutical marketing. On the other hand, the efficacy of therapeutic was monitored.

Methodology: 505 patients with COVID-19 from Hodeidah showed that respiratory infection illness can range from mild to critical. Mild to moderate was 386 case (78.93 %), severe cases was 73 cases (15.13 %) and critical cases was 28 cases (5.93 %). The major criteria for cases admission namely clinical examination, case definition, clinical investigation, radiological finding, hematological and biochemical finding and molecular biology assay namely real – time - polymerase chain reaction (RT- PCR). The pharmacological and therapeutic properties of supportive therapy of COVID-19 included" antipyretic agent, antiviral (oseltamivir) analgesic agent, glucocorticoid (dexamethasone), anticoagulant agent (enoxaparin), anti-cough (acetylcysteine), antibiotics broad spectrum to treat the secondary infection namely azithromycin, ceftriaxone, ceftazidime, piperacillin and tazobactam, meropenam, and moxifloxacin. In addition, oxygen therapy is the major treatment that used in treatment of severe and critical cases. Also, vitamins (C and D) and mineral namely zinc were prescribed and proton pump inhibitor (PPI) (pantaprazol), anti-emetic (Ondansetron). Fluid therapy namely ringer lactate or normal saline were used in management of COVID -19.

Results: The results showed that the age of the patients included in this study between 3- 92 years with 4.0 to 1.0 males: females. 49 patients were admitted in COVID-19 isolation department of Hodeidah, Yemen. 25/49 cases (54.34%) were recovered with national guideline, the average of RR pre - treatment was 30 ± 5 breath / minute and reduced to 23 ± 2 breath / minute post – treatment with significantly statistically different ($p < 0.05$). On the other hand, the clinical symptoms namely difficult in breathing, cough, fever, and headache reduced to absent absolutely (100 %). In addition, the results showed that the average of WBC pre - treatment was 20 ± 4 ($\times 10^9/L$) and reduced to 7 ± 2 ($\times 10^9/L$) post – treatment with significantly statistically different ($p < 0.05$). Also, lymphocytes decreased with COVID -19 infection to 2 % with average ($10 \pm 5\%$) and improved post - treatment to 20 %. In addition, neutrophil increased to 95 % with average ($80 \pm 15\%$) and improved to 70 % post-treatment. the average of WBC pre - treatment was 20 ± 4 ($\times 10^9/L$) and reduced to 7 ± 2 ($\times 10^9/L$) post-treatment with significantly statistically different ($p < 0.05$). The outcome finding of management that 19 cases (3.88 %) of severe and 6 cases (1.22 %) of critical were recovered. On the other hand, 1 case of severe and 21 cases of critical COVID-19 were died.

Conclusion: In spite of lack of specific antiviral drugs, the approach of pharmacology and therapeutics had good impact at level of mild to moderate cases, and severe cases while the critical cases needs to develop at level critical cases in intensive care unit (ICU) with mechanical ventilation (MV) procedure.

Keywords: COVID-19; pharmacology; therapeutic; Hodeidah; Yemen.

1. INTRODUCTION

“Since December 2019, the world faced a new pandemic caused by a virus in the Coronaviridae family, namely SARS-CoV-2, disrupting global public health and world economies. COVID-19, the disease caused by SARS-CoV-2, is a viral

infection that affects the respiratory system, resulting in respiratory syndrome” [1]. “COVID-19 is considered the most serious of pandemics since 1918 when the “Spanish flu” (H1N1) emerged” [2]. “Before 2021, there were no drugs or vaccines proven to treat or prevent infection caused by SARS-CoV-2” [3]. “However,

numerous studies were being conducted” [3,4]. “At the end of December 2019, the dramatic story of COVID-19 began and new species of the virus from the Coronaviridae family emerged in the city of Wuhan, China” [2,5]. “As of April 10, 2020, there were cases confirmed infection in Yemen and to spread in different area of Yemen. The signs and symptoms of COVID-19 present at illness onset vary, but over the course of the disease, most persons with COVID-19 will experience the following: Fever or chills, cough, shortness of breath or difficulty breathing, fatigue, muscle or body aches, headache, new loss of taste or smell, sore throat, congestion or runny nose, nausea or vomiting and diarrhea”. [6-8]. Symptoms differ with severity of disease such as fever, cough and shortness of breath are more commonly reported among hospitalized patients than non-hospitalized. This study reviews the therapeutic strategies to deal with the disease before effective vaccines and antiviral drugs against the causative virus became available, and only symptomatic, supportive and prevention were at our disposal.

2. METHODOLOGY

2.1 Study Design

This was a case series study based on medical file about monitoring the efficacy and safety of national guideline based on using laboratory and clinical outcome in COVID – 19 outbreak in Hodeidah, Yemen. The study was conducted during the period 1st June 2020 to 30th September 2020.

2.2 Study Area

This study was carried out in COVID -19 Isolation Department, Center of Tropical Medicine and Infectious Diseases (CTMES) , Authority of Public AL- Thawara Hospital , Hodeidah . The study included all patients that were admitted and treated in isolation department and patients were treated at home. Note : The area is vector – borne diseases endemic , therefore the malaria and dengue were diagnosed to manage the coinfection [18-25] .

2.3 Study Population

The study included 505 patients 386 cases (76.43 %) of mild and moderate cases) ; 70 cases (13.86 %) of severe were treated at home. On the other hand , the pharmacological

profiles were applied on 21 patients of severe illness (4.16 %) and 28 patients (5.54%) of critical illness were treated at isolation department. 49 patients (9.7 %) needed admission in an intensive care unit (ICU). Note : The biomedical parameters were monitored carefully in severe and critical cases only (49 cases).

2.4 Diagnosis of COVID-19

The major criteria for cases admission namely clinical examination, case definition, clinical investigation, radiological finding, hematological finding. On the other hand, the cases were confirmed using molecular biology assay namely real – time - polymerase chain reaction (RT-PCR) [26-32].

2.5 National Guideline

The national guideline treatment of COVID -19 for management of mild, moderate, severe, and critical cases were described in Table 3. [33-34].

2.6 Pharmacological and Therapeutics Approach

The pharmacological and therapeutic properties of supportive therapy of COVID - 19 included " antipyretic agent to reduce the fever, analgesic agent to reduce the pain, glucocorticoid (dexamethasone) as anti-inflammatory agent, anticoagulant agent (enoxaparin) to prevent the small embolism formation in lung, bronchodilator agent (salbutamol), anti-cough to reduce productive and non-productive cough (acetylcysteine), antibiotics broad spectrum to treat the secondary infection namely azithromycin, ceftriaxone, ceftazidime piperacillin and tazobactam, meropenam, and moxifloxacin. In addition, oxygen therapy is the major treatment that used in treatment of severe and critical cases. Also , vitamins (C and D) and mineral namely zinc are used to support the immune system. Finally , proton pump inhibitor (PPI) to prevent the gastric stress (pantoprazol) , anti-emetic (ondansetron). Fluid therapy namely ringer lactate or normal saline were used in management of COVID – 19.

2.7. Statistical Analysis

The differences between the females and males groups were analysed by using Excel 2010 and Statistical Process Social Sciences (SPSS)

Table 1. Pharmacological properties of support therapy used in treatment of COVID-19 patients, Hodeidah, Yemen

Medicines Used	MOA	Side Effects	Monitoring	Antidote
Paracetamol	Paracetamol has a central analgesic effect that is mediated through activation of descending serotonergic pathways.	Nausea, vomiting, constipation	Temperature Pain and fever relief , Caution must be observed due to liver toxicity at high doses of paracetamol.	N-acetylcysteine (NAC)
Azithromycin	Prevents bacteria from growing by interfering with their protein synthesis. It binds to the 50S subunit of the bacterial ribosome, thus inhibiting translation of mRNA	Anorexia, nausea, vomiting, and diarrhea are common	Concomitant use of hydroxychloroquine with azithromycin as the association can lead to a higher risk of QT interval prolongation and cardiac arrhythmias ⁽¹⁰⁾ .	No specific antidote
Ondansetron	5-HT ₃ antagonists	Headache, dizziness, and constipation	ECG (if applicable in high-risk or elderly patients); potassium, magnesium. Monitor for signs of serotonin syndrome; monitor for decreased bowel activity.	No specific antidote
Enoxaparin	low-molecular-weight heparin inhibit activated factor X	Hypochromic anemia, thrombocytopenia, hemorrhage, bleeding complications.	Measuring factor Xa inhibition (anti-factor Xa activity).	Limited experience suggests that 1 mg of protamine sulfate may be used to partially neutralize 1 mg of enoxaparin.
Pantoprazole	Proton pump inhibitors (PPIs)	Diarrhea, headache, and abdominal pain Reduction in oral cyanocobalamin absorption		
Dexamethasone	Dexamethasone bind to the DNA of glucocorticoid properties to modify transcription proteins inhibiting leukocyte infiltration to the site of inflammation, interfering with the effect of inflammatory mediators, suppress humoral responses and reduce edema and/or scarring of tissues.	Metabolic Effects Adrenal Suppression, Peptic ulcer, Nausea, dizziness, and weight loss in some patients. Hypomania or acute psychosis. Increased intraocular pressure and glaucoma. Benign intracranial Hypertension. Growth-suppressing. Sodium and fluid retention and loss of potassium. Potential adverse events:	A thorough history and physical examination should be performed to assess for risk factors.	No specific antidote

Medicines Used	MOA	Side Effects	Monitoring	Antidote
		Increased risk for infection Hyperglycemia. Peripheral edema . Increased appetite Insomnia, irritability, delirium		
Oseltamivir	Neuraminidase inhibitor	Vomiting, diarrhea, headache, and trouble sleeping. Other side effects may include psychiatric symptoms and seizures	A thorough history and physical examination	No specific antidote
Ceftriaxone	Inhibition of cell wall synthesis	Rash, pruritus, fever, eosinophilia, urticarial, anaphylaxis, colitis, diarrhea, nausea and vomiting, pseudomembranous colitis.	Therapeutic: Culture and sensitivities, serum levels, signs and symptoms of infection, white blood cell count Toxic: Urinalysis, BUN, SCr, AST and ALT, skin rash, Neutropenia and leukopenia, Prothrombin time in patients with renal or hepatic impairment or poor nutritional state, as well as patients receiving a protracted course of antimicrobial therapy, and patients previously stabilized on anticoagulant therapy.	No specific antidote.
Ceftazidime	Inhibits bacterial cell wall synthesis	Pain, swelling, burning, or irritation around the IV needle; nausea, vomiting, diarrhea, stomach pain; or vaginal itching or discharge.	As ceftriaxone	No specific antidote
Piperacillin and Tazobactam	Piperacillin, a broad spectrum, semisynthetic penicillin exerts bactericidal activity by inhibition of both septum and cell wall synthesis. Tazobactam, a beta-lactam structurally related to penicillins, is an inhibitor of many beta-lactamases, which commonly cause resistance to penicillins and cephalosporins. Tazobactam extends the antibiotic spectrum of piperacillin to include	The same for the parent compound	As ceftriaxone	No specific antidote

Medicines Used	MOA	Side Effects	Monitoring	Antidote
	many beta-lactamase-producing bacteria that have acquired resistance to piperacillin alone.			
Acetylcysteine	Acetylcysteine exerts its mucolytic action through its free sulfhydryl group, which opens the disulfide bonds and lowers mucus viscosity	Bronchospasm , Disagreeable odor. Drowsiness. Fever. Coughing up blood. Increased volume of bronchial secretions.Irritation of tracheal or bronchial tract.	Asses patient for nausea, vomiting and skin rash. Reassess LFTs for possible hepatotoxicity every 4 to 6hours,	
Moxifloxacin	It functions by inhibiting DNA gyrase, a type II topoisomerase, and topoisomerase IV, enzymes necessary to separate bacterial DNA, thereby inhibiting cell replication Targeting the bacterial DNA gyrase (type II topoisomerase) and topoisomerase IV thus inhibiting bacterial DNA synthesis and leading to cleavage of bacterial DNA and rapid bacterial death.	Diarrhea , dizziness, headache nausea , stomach cramps	Therapeutic: Culture and sensitivities, signs and symptoms of infection Toxic: Urinalysis, BUN, SCr, AST and ALT, Physical examination: encephalopathic changes Monitor for QT prolongation	
Meropenam	It inhibits bacterial cell wall synthesis like other β -lactam antibiotics.	A skin rash, constipation, diarrhea, headache, nausea and vomiting	Therapeutic: Resolution of clinical signs of infection (fever, decreased white blood cell count), Culture and sensitivity, CBC w/differential, urinalysis, temperature Toxicity: Hepatic and renal function tests during therapy	
Vitamin D3 cholecalciferol	Is a fat-soluble vitamin helps body absorbs calcium, and phosphorus	Hypersensitivity reactions such as angioedema or laryngeal edema, nusea and vomiting	A blood test is recommended to monitor blood levels of 25(OH)D three months after beginning treatment.	Supportive treatment
Vitamin C (ascorbic acid)	A water-soluble vitamin Antioxidant and free radical scavenger that has anti-inflammatory properties	Individuals who have a history of kidney stone formation and those who experience iron overload should exercise caution before using supplemental vitamin C.		
Zinc	It is a naturally occurring mineral	nausea, vomiting, diarrhea		

Table 2. Pharmacokinetics properties of support therapy used in treatment of COVID-19 patients, Hodeidah, Yemen

Medicines Used	Absorption	Distribution	Metabolism	Excretion
Paracetamol	Absorbed from the GIT peak plasma concentrations occurring about 30 minutes to 2 hours after ingestion	Distributes rapidly and evenly throughout most tissues and fluids and has a volume of distribution of approximately 0.9L/kg. 10 to 20% of the drug is bound to red blood cells	Metabolized in the liver (90-95%) and excreted in the urine mainly as the glucuronide and sulphate conjugates	Excreted in the urine mainly as the glucuronide and sulphate conjugates. Less than 5% is excreted as unchanged
Azithromycin	Rapidly absorbed and well tolerated orally. It should be administered 1 hour before or 2 hours after meals.	Penetrates into most tissues except(CSF) and phagocytic cells extremely well, with tissue concentrations exceeding serum concentrations by 10- to 100-fold.	Not inactivate cytochrome P450 enzymes	
Ondansetron	Completely and rapidly absorbed from GIT after oral administration	Widely distributed (volume of distribution approximately 160L) and binds moderately (70 to 76%) to plasma proteins;	Extensive hepatic metabolism	Eliminated by renal and hepatic excretion.
Enoxaparin	Full-dose enoxaparin therapy is 1 mg/kg subcutaneously every 12 hours. Bioavailability is 92%.	Volume of distribution of anti-factor Xa activity is about 6 L.	NA	Elimination half-life based on anti-factor Xa activity is about 4.5 hours after S.C. administration.
Pantoprazole	Well absorbed, undergoes little first-pass metabolism resulting in an absolute bioavailability of approximately 77%.	The apparent volume of distribution (V) is approximately 11.0 to 23.6 L, and the serum protein binding is ~98%	Extensively metabolized in the liver through the cytochrome P-450 system, predominantly by CYP2C19 demethylation with subsequent sulfation	The serum elimination half-life of about 1.1 hours
Dexamethasone	Following 20 mg dose of dexamethasone median time to peak concentrations (Tmax) (range: 0.5 to 4 hours).	Dexamethasone is about 77% bound to human plasma proteins in vitro.	Dexamethasone is metabolized by CYP3A4.	Renal excretion of dexamethasone is less than 10% of total body clearance. Less than 10% of dexamethasone is excreted in the urine.
Oseltamivir	Oral bioavailability is over 80% and is extensively metabolised to its active form upon first-pass through the liver.	It has a volume of distribution of 23–26 litres. Its half-life is about 1–3 hours .	Rapidly metabolized to active form , oseltamivir carboxylate.	More than 90% of the oral dose is eliminated in the urine as the active metabolite.
Ceftriaxone	Parenteral route of administration Rapidly and completely absorbed	C _{max} : 123-151mcg/L Volume of distribution: 10.7L		33- 67 % is excreted in the urine as unchanged drug, and the

	following intramuscular administration.			remainder is secreted in the bile and ultimately is found in the feces as microbiologically inactive compounds.
Ceftazidime	Administered I.V. and I.M.	Distributed widely into most body tissues and fluids, including the gallbladder, liver, kidneys, bone, sputum, bile, and pleural and synovial fluids; unlike most other cephalosporins, ceftazidime has good CSF penetration; it crosses the placental barrier. Ceftazidime is 5% to 24% protein-bound.	Not metabolized.	Excreted mainly in urine by glomerular filtration; small amounts of drug appear in breast milk. Elimination half-life is about 1 1/2 to 2 hours in patients with normal renal function; up to 35 hours in patients with severe renal disease. Hemodialysis or peritoneal dialysis removes ceftazidime.
Piperacillin and Tazobactam	Piperacillin/Tazobactam 4 g/0.5 g Powder for Solution for Infusion. The peak piperacillin and tazobactam concentrations after 4 g / 0.5 g administered over 30 minutes by intravenous infusion are 298 µg/ml and 34 µg/ml respectively.	Both piperacillin and tazobactam are approximately 30% bound to plasma proteins. widely distributed in tissue and body fluids including intestinal mucosa, gallbladder, lung, bile and bone.	Piperacillin is metabolised to a minor microbiologically active desethyl metabolite. Tazobactam is metabolised to a single metabolite, which has been found to be micro-biologically inactive.	Piperacillin and tazobactam are eliminated by the kidney via glomerular filtration and tubular secretion.
Acetylcysteine	Most inhaled acetylcysteine acts directly on mucus in the lungs; the remainder is absorbed by pulmonary epithelium. After oral administration, drug is absorbed from the GI tract.	Unknown	Hepatic. Acetylcysteine undergoes rapid deacetylation in vivo to yield cysteine or oxidation to yield diacetylcystine.	Unknown
Moxifloxacin	Well absorbed from the gastrointestinal tract. Absolute oral bioavailability is approximately 90%. Food has little effect on absorption.	Volume of distribution is 1.7 to 2.7 L/kg% 50% bound to serum proteins, independent of drug concentration.	Approximately 52% of oral or intravenous dose is metabolized via glucuronide and sulphate conjugation. The cytochrome P450 system is not involved in metabolism. The sulphate conjugate accounts for 38% of the dose, and the glucuronide conjugate accounts for 14% of the dose.	Approximately 45% of an oral or intravenous dose of moxifloxacin is excreted as unchanged drug (~20% in urine and ~25% in feces).

Meropenam	Administered I.V	Distributed into most body fluids and tissues, including CSF. It's only about 2% bound to plasma protein.	Thought to undergo minimal metabolism. One inactive metabolite has been identified.	Excreted unchanged primarily in urine. Elimination half-life of drug in adults with normal renal function and children age 2 and older is about 1 hour and 1.5 hours in children age 3 months to 2 years.
Vitamin D3 cholecalciferol			The three main steps in vitamin D metabolism, 25-hydroxylation, 1 α -hydroxylation, and 24-hydroxylation are all performed by cytochrome P450 mixed-function oxidases (CYPs). These enzymes are located either in the endoplasmic reticulum (ER) (e.g., CYP2R1) or in the mitochondria (e.g., CYP27A1, CYP27B1, and CYP24A1).	
Vitamin C (ascorbic acid)	Vitamin C absorption occurs through small intestine (distal intestine) by active transport mechanism.	Vitamin C is widely distributed in all the body tissues.	Vitamin C metabolites (oxalate salts) and unmetabolized vitamin C are excreted by kidneys. Few percentage of vitamin C is excreted through feces.	The urinary excretion of vitamin C is dose dependent.
Zinc	Zinc is absorbed throughout the whole small intestine	distributed to the peripheral tissues, including skeletal muscle (60%), bone (30%), skin (5%), and other tissues (5%)		Zinc is lost from the body through the kidneys, skin, and intestine.

[9-17]

Table 3. National guideline of COVID-19 Patients, Hodeidah, Yemen

Medicines Used	Mild cases	Moderate	Severe	Critical
Paracetamol 500 mg Tablet	three times a day (7 days)	three times a day (7 days)	NA	NA
Paracetamol 1000 mg IV	NA	NA	three times a day (10-14 days)	three times a day (10-14 days)
Azithromycin 500 mg Tablet	once a day (6 days)	once a day (6 days)	once a day (6 days)	once a day (6 days)
Ondansetron 8 mg IV	NA	NA	If necessary	If necessary
Enoxaparin 60 mg Subq.	NA	NA	once a day (14 days)	once a day (14 days)
Pantoprazole 40 mg IV	NA	NA	once a day (14 days)	once a day (14 days)
Acetylcysteine IV 300 mg	NA	NA	three times a day (14 days)	three times a day (14 days)
Oseltamivir 75 mg Tablet	NA	NA	once dose (14 days)	once dose (14 days)
Dexamethasone 10 – 20 mg IV	NA	NA	twice a day (10 days)	twice a day (10 days)
Ceftriaxone 1 g IV	NA	NA	twice a day (10-14 days)	NA
Ceftazidime 1 g IV	NA	NA	twice a day (10-14 days)	NA
Piperacillin and Tazobactam 4.5 g IV	NA	NA	NA	three times a day (10-14 days)
Moxifloxacin 400 mg Tablet	NA	If necessary	NA	NA
Moxifloxacin 400 mg IV	NA	NA	once a day (10 -14 days)	once a day (10 -14 days)
Meropenam 1 g IV	NA	NA	NA	three times a day (10-14 days)
Vitamin D 50000 IU Tablet	once weekly	once weekly	once weekly (two weeks)	once weekly (two weeks)
Viatmine C 1g Tablet	once a day (14 days)	once dose (14 days)	NA	NA
Vitamine C 1g IV	NA	NA	once a day (14 days)	once a day (14 days)
Zink 300 mg Tablet	once a day (14 days)	once dose (14 days)	once dose (14 days)	once dose (14 days)

Note :

- 1) Dose of dexamethasone was prescribed 20 mg in first five days and reduced into 10 mg in other five (first wave of COVID-19)
- 2) The doses of antibiotics were adjusted according to creatinine clearance
- 3) NA : Not Applicable
- 4) *Mild cases (symptoms of respiratory infection namely fever , cough , sore throat , headache , ...etc) , moderate cases (clinical signs of non severe pneumonia) , severe cases (clinical signs of severe pneumonia namely dyspnea, hypoxia, or lung involvement on imaging) and critical cases (acute respiratory distress syndrome "ARDS: , respiratory failure, shock, or multi-organ system dysfunction) [33-34].*

version 15 to calculate the descriptive analysis and paired t- test at $\alpha = 0.05$ that were used to explore the effectiveness of national guideline treatment of COVID -19 pre and post-treatment in Hodeidah city, Yemen.

3. RESULTS

3.1. Clinical Response

The results showed that the age of the patients included in this study between 3- 92 years with 1.5 : 2.0 male: female. 49 patients were admitted in COVID – 19 isolation department of CTMID. 25/46 cases (54.34%) were recovered with national guideline, the average of RR pre - treatment was 30 ± 5 breath / minute and reduced to 23 ± 2 breath / minute post – treatment with significantly statistically different ($p < 0.05$) . On the other hand , the clinical symptoms namely difficult in breath, cough , feve , and headachic absent abso[utely]ly (100 %).

3.2. Laboratory Response

3.2.1 Hematological parametrs

The results showed that the average of WBC pre - treatment was 20 ± 4 ($\times 10^9/L$) and reduced to 7 ± 2 ($\times 10^9/L$) post – treatment with significantly statistically different ($p < 0.05$) (Table 5). On the other hand , the WBCs count, which had reached a peak of 34 ($\times 10^9/L$) as maximum value . Also , lymphocytes decreased with COVID -19 infection to 2 % with aveage ($10\pm 5\%$) and improved post - treatment to 20 % . In addition, neutrophil increased to 95 % with aveage ($80\pm 15\%$) and improved to 70 % post

treatment . the average of WBC pre - treatment was 20 ± 4 ($\times 10^9/L$) and reduced to 7 ± 2 ($\times 10^9/L$) post – treatment with significantly statistically different ($p < 0.05$).

3.2.2 Biochemical parameters

The results showed that the average of blood glucose pre - treatment was 370 ± 200 mg/dl and reduced to 140 ± 40 mg/dl post - treatment based on insuline therapy with significantly statistically different ($p < 0.05$) (Table 6). On the other hand , the blood glucose, which had reached a peak of 550 mg/dl as maximum value Also , albumin decreased with COVID -19 infection to 1.5 g/dl with average (2.2 ± 0.5 g/dl) and improved post - treatment to 3 ± 0.5 g/dl ($p < 0.05$) (Table 7). In addition, CRP increased to 98 mg/L % with aveage (70 ± 28 mg/L) and very slow[e]ly decreased [kinetic] post – treatment (23 ± 8 mg/L) ($p > 0.05$). The liver enzyme (Table 7), serum creatinine (Table 9) and serum electrolytes (Table 10) were not correlated to COVID-19 infection , therefore , these parameters were not changed post – treatment (Tables 7 , 9 and 10).

3.3 Outcome

The outcome finding of management that 49 severe and critical cases were admitted recovered at center, 19 cases (3.88 %) of severe and 6 cases (1.22 %) of critical were recovered at isolation center . On the other hand ,1 case (0.19%) of severe and 21 cases (4.15 %) of crtical COVID-19 were died. On the other hand , 386 cases (76.43 %) of mild and 54 cases (11.04 %) of severe were admitted at home that recovered.




Fig. 1. Kinatic of lymphocytes pre – within and post – treatment of mild and moderate cases

Table 4. Monitoring the level of oxygen saturation and respiratory rate (RR) pre and post - treatment (X ± SD ; n:49)

Parameters	Pre - treatment	Post - treatment	p-value
Oxygen saturation (%)	75 ± 10	92 ± 2	$p < 0.05^*$
Respiratory Rate (perth/minute)	30 ± 2	25 ± 1	$p < 0.05^*$
Blood Pressure (mmHg)			
• Systolic	120 ± 20	122 ± 10	$p > 0.05$
• Diastolic	80 ± 10	82 ± 10	

* : Significant p – value less than 0.05

Table 5. Monitoring the level of WBC and differential count pre- and post-treatment (X ±SD: n:49)

Parameters	Pre - treatment	Post - treatment	p-value
WBC (×10 ⁹ /L)	20 ± 4	7 ± 2	$p < 0.05^*$
Neutrophils (%)	80 ± 15	50 ± 15	$p < 0.05^*$
Eosinophils (%)	Normal	Normal	$p > 0.05$
Basophils (%)	Normal	Normal	$p > 0.05$
Lymphocytes (%)	10 ± 5	25 ± 5	$p < 0.05^*$
Monocytes (%)	Normal	Normal	$p > 0.05$

* : Significant p – value less than 0.05




Fig. 2. Kinatic of neutrophils pre – within and post – treatment of mild and moderate cases




Fig. 3. Kinatic of lymphocytes pre , within and post – traetment of severe cases




Fig. 4. Kinatic of neutrophils pre , within and post – traetment of severe case

Table 6. Monitoring the level of blood glucose pre and post - treatment (X±SD; n:49)

Parameters	Pre - treatment	Post - treatment	p-value
Blood glucose (mg/dl)	370 ± 200	140 ± 40	p < 0.05*

* : Significant p – value less than 0.05

Table 7. Liver function test pre and post - treatment (X±SD , n:49)

Parameters	Pre - treatment	Post - treatment	p-value
GOT	45.42±26.31	44.31±20.25	p > 0.05
GPT	45.38 ± 29.31	43.40 ± 29.31	p > 0.05
Albumin	2.2 ± 0.5	3.100 ± 0.5	p < 0.05*

Note : The liver enzymes need long time to change (cohort) while the albumin level was improved : : Significant p – value less than 0.05

Table 8. Immunological marker (CRP) response pre and post - treatment (X±SD ; n:49)

Parameters	Pre - treatment	Post - treatment	p-value
CRP mg/dl	70 ± 28	65 ± 20	p > 0.05

Note : The CRP needs long time to change

Table 9. Renal function test pre and post – treatment (X±SD; n:49)

Parameters	Pre – treatment	Post – treatment	p-value
Creatinine mg /dl	0.5 – 1.5	0.4 – 1.6	p > 0.05

Table 10. Electrolytes assay pre and post treatment (X±SD ; n:49)

Parameters	Pre - treatment	Post - treatment	p-value
K ⁺	5.19±4.67	5.19±4.67	p > 0.05
Na ⁺	135.35±5.49	134.44±4.55	p > 0.05
Cl ⁻	100.28±6.31	99.12±5.33	p > 0.05

4. DISCUSSION

“Clinicians namely Intensive Care Unit (ICU) , tropical medicine, and clinical pharmacologist are frequently asked to monitor the effects of

national guideline treatment with the objective of ensuring safe and effective therapy. Already in monitoring of supportive therapy care includes therapeutic response bilogically and clinically” [35]. Our study monitored the efficacy of

national guideline treatment based hematological parameters and clinical response to treatment was used namely .

“Findings of the present study showed that almost all the WBCs of COVID-19 patients sampled from Hodeidah Yemen (collected systemically from COVID-19 patients) were found to be more than normal values . These results were estimated according to direct hematological analyzer that was validated using a classical approach for the assay of WBC and related parameters. This approach gives enough guarantees for the results that will be generated by this method during blood analysis” [36-37].

“Monitoring efficacy of supportive therapy care and response to treatment of COVID-19 infection can be assessed using biological and clinical parameters of improvement include symptoms and signs (eg, a decrease in fever, tachycardia, cough , respiratory rate), (eg, decreasing WBC count and neutrophil % within normal range ,and increase lymphocytes to normal range. Although hematological criteria namely WBC , neutrophil , and lymphocytes are commonly used as economical indicators in assessing response to infectious disease therapy in Hodeidah outbreak management, WBCs improvement can frequently lag behind clinical improvement. In our study of hematological and clinical follow-up of patients with COVID-19, clinical cure was observed in 128/149 cases (85.90%) after 14 days of follow-up of patients. During the study period, there were no drugs or other therapeutics presently approved by the U.S. Food and Drug Administration (FDA) to prevent or treat COVID-19. Current clinical management includes infection prevention and control measures and supportive care, including supplemental oxygen and mechanical ventilatory support when indicated” [38]. In Yemen, the pharmacological and therapeutic approaches for management of COVID-19 were using available [of] medicines in local pharmaceutical marketing.

“Although systemic corticosteroids for the treatment of viral pneumonia or ARDS were not recommended, dexamethasone treatment is recommended by the National Health Service in the UK and the National Institutes of Health (NIH). In the US for patients with COVID-19 who [are] require supplemental oxygen but are not mechanically ventilated. Dexamethasone is not recommended in patients with COVID-19 who do not require supplemental oxygen or hospitalization” [39-41].

“The Infectious Diseases Society of America (IDSA) guideline panel suggests the use of glucocorticoids for patients with severe COVID-19 where severe is defined as patients with SpO₂ ≤94% on room air, and those who require supplemental oxygen, mechanical ventilation, or extracorporeal membrane oxygenation (ECMO). The IDSA recommends against the use of glucocorticoids for those with COVID-19 without hypoxemia requiring supplemental oxygen. The World Health Organization (WHO) recommends systemic corticosteroids rather than no systemic corticosteroids for the treatment of people with severe and critical COVID-19 (strong recommendation, based on moderate certainty evidence)” [42-43] . “The WHO suggests not to use corticosteroids in the treatment of people with non-severe COVID-19 (conditional recommendation, based on low certainty evidence). Dexamethasone reduced deaths by approximately one third in patients requiring ventilation and by one fifth in those requiring oxygen” [32] . “Among patients hospitalized with mild-to-moderate COVID-19, the use of azithromycin, did not improve clinical status at 15 days as compared with standard care” [44].

“For pain management during COVID-19, the paracetamol used safely to alleviate symptoms of COVID-19 such as fever, headache, and acute or chronic pain. However, caution must be observed due to liver toxicity at high doses of paracetamol” ⁽¹⁰⁾ . “Of importance, in current pain practice, opioids are generally used as only part of a treatment plan, which includes physical therapy, pain psychology, interventional procedures, and other ancillary therapies. Because COVID-19 patients have a higher incidence of venous thromboembolism and anticoagulant therapy is associated with reduced ICU mortality, it is suggested that patients should receive thromboprophylaxis. Moreover, in the case of known thrombophilia or thrombosis, full therapeutic-intensity anticoagulation (e.g., enoxaparin 1 mg/kg twice daily) is indicated” [45].

“Different countries recommended the use of chloroquine in management of COVID-19 but WHO stopped its using” [46]. Our study did use Chloroquine in CTMID, Authority of Public AL Thawarah Hospital , Hodeidah, Yemen. Chloroquine (500 mg every 12 hours), and hydroxychloroquine (200 mg every 12 hours) were proposed by Gautret et al [47]. “The study showed that hydroxychloroquine was significantly associated with viral load reduction until viral disappearance and this effect was

enhanced by the macrolides azithromycin. In vitro and in vivo studies, indeed, have shown that macrolides may mitigate inflammation and modulate the immune system. In particular, these drugs may induce the downregulation of the adhesion molecules of the cell surface, reducing the production of proinflammatory cytokines, stimulating phagocytosis by alveolar macrophages, and inhibiting the activation and mobilization of neutrophils [11]. However, further studies are needed for recommending the use of azithromycin, alone or associated with other drugs such as hydroxychloroquine, outside of any bacterial overlaps. Again, attention must be paid with the concomitant use of hydroxychloroquine with azithromycin as the association can lead to a higher risk of QT interval prolongation and cardiac arrhythmias and chloroquine can also induce QT prolongation” [11].

“In patients with confirmed COVID-19 pneumonia, community-onset bacterial co-infection is uncommon, so the empirical treatment of antibiotics for patients 18 years and older with suspected community-acquired pneumonia was azithromycin as oral for moderate or severe pneumonia, in case of unsuitable we used moxifloxacin. However, intravenous antibiotics for moderate or severe pneumonia, we used ceftriaxone. A recent in silico study demonstrated that the fluoroquinolones, ciprofloxacin and moxifloxacin, exert strong capacity for binding to SARS-CoV-2 main protease (Mpro), indicating that fluoroquinolones may inhibit SARSCoV-2 replication” [48]. The intravenous antibiotics for severe pneumonia (for example, symptoms or signs of sepsis or ventilator-associated pneumonia) or when there is a higher risk of resistance, the options included piperacillin with tazobactam: 4.5 g three times a day, increased to 4.5 g four times a day if infection is severe, or Ceftazidime: 2 g three times a day if the above options are unsuitable for moxifloxacin.

“The emergence of the COVID-19 pandemic has affected individuals and society on multiple levels. The overall focus has been on preventing the spread of the disease and discovering treatment options. However, the psychological impact of the illness should not be overlooked. Patients face multiple stressors due to the COVID-19 crisis including but not limited to fear of becoming infected and of infecting others, especially family members, inadequate access to testing, disrupted regular medical care, financial

losses, distress related to social distancing and quarantine, and uncertainty of the duration of the pandemic” [49]. “Some reports raise concerns of a long-lasting mental health impact as a consequence of the pandemic. It is anticipated that patients infected with COVID-19 may suffer from mood dysregulation, anxiety, anger, and worsening of any preexisting mental illness” [50]. With long hospitalization and the potential need to intubation and mechanical ventilation, light sedation is required to reduce risk of post – intensive care unit syndrome and adverse outcomes.

Our study focused on the pharmacological and therapeutic properties of medications available in Yemen. On the other hand, in response to COVID-19, FDA quickly created, in March 2020, the Coronavirus Treatment Acceleration Program (CTAP), which was designed to help facilitate the development of drugs and biological products (other than vaccines) for COVID-19 therapeutics. The development of COVID-19 therapeutics that are effective against current and future variants remains a critical priority. CTAP is leveraging the agency’s scientific experience and knowledge of COVID-19 to support current and future clinical trials testing new treatments for COVID-19. From these efforts, we continue to gain valuable knowledge about the safety and effectiveness of potential therapies for COVID-19. These medications were used in other countries 1) antiviral agents (Remdesivir, Favipiravir, Molnupiravir, PF-07321332, Paxlovid), 2) convalescent plasma and immunoglobulin, 3) monoclonal antibodies usable in mono therapy (Sotrovimab), 4) monoclonal antibodies used in combination (Banlavitimab +Etesevimab, Casirivimab, +Imdevinab, AZD7442 namely Cilgavimab (AZD1061) + Tixagevimab (AZD8895) Evusheld, Combination of neutralizing monoclonal antibodies BR11-196 and BR11-198. BR11-196 and BR11-198 are noncompetitive anti-SARS-CoV-2 monoclonal antibodies, 5) Immunomodulatory drugs (Tocilizumab, Sarilumab, Anakinra, Canakinumab, Baricitinib, Tofacitinib, Ruxolitinib, Adalimumab, Certolizumab, Infliximab, Etanercept, Golimumab, Itolizumab, Ravulizumab, Lemilumab, 6) Renal Replacement Therapy Solutions. On the other hand, some medications available in Yemen but not prescribed by medical doctors in our center namely Ivermectin and Colchicine, and Lopinavir/Ritonavir [51,52]

5. CONCLUSION

In spite of lack of specific antiviral drugs, the approach of pharmacology and therapeutics had good impact at level of mild to moderate cases, and severe cases while the critical cases needs to develop at level critical cases in ICU with mechanical ventilation (MV) procedure.

CONSENT AND ETHICAL APPROVAL

The study was integrated within the clinical practice. Patients received simple explanation for the aim of the study. If agreed to participate verbally, biological and clinical data were collected and interview was conducted. Confidentiality of the collected data was achieved by keeping data record in a locked room with limited access to the research team only. according to Center of Tropical Medicine and Epidemiology Studies – Hoediad University (CTMES – HU , Yemen declaration.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization (WHO), Director-General's opening remarks at the media briefing on COVID-19;2020.
2. Al Hasan SM, Saulam J, Kanda K & Hirao T. The novel coronavirus disease (COVID-19) outbreak trends in mainland China: a joinpoint regression analysis of the outbreak data from January 10 to February 11, 2020. [Preprint]. Bull World Health Organ. E-pub: 17;2020.
3. Centers for Disease Control and Prevention (CDC) . Coronavirus disease 2019 (COVID-19) clinical care;2020.
4. National Institute of Health (NIH) . Adaptive COVID-19 Treatment Trial (ACTT)—ClinicalTrials;2020.
5. World Health Organization (WHO). Novel Coronavirus (2019-nCoV) Situation report– 1 , Geneva , Switerland;2020
6. Guan WJ, Ni ZY, Hu Y, et al. Clinical Characteristics of Coronavirus Disease 2019 in China. N Engl J Med 2020;382: 1708-20.
7. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: A descriptive study. Lancet 2020;395:507-13.
8. Pan L, Mu M, Yang P, et al. Clinical Characteristics of COVID-19 Patients with digestive symptoms in Hubei, China: A Descriptive, cross-sectional, multicenter study. Am J Gastroenterol; 2020.
9. Katzung BG, ed. Basic and clinical pharmacology. 13th ed. New York: McGraw-Hill Education; 2015.
10. Pergolizzi J, Varrassi G, Magnusson P, LeQuang JA, Paladini A, Taylor R, Wollmuth C, Breve F, Christo P. COVID-19 and NSAIDS: A narrative review of knowns and unknowns. Pain Ther;2020. Available:https:// doi.org/10.1007/s40122-020-00173-5.
11. Mercurio NJ, Yen CF, Shim DJ, Maher TR, McCoy CM, Zimetbaum PJ, Gold HS. Risk of QT interval prolongation associated with use of hydroxychloroquine with or without concomitant azithromycin among hospitalized patients testing positive for coronavirus disease 2019 (COVID-19). JAMA Cardiol. 2020 May 01; [PMC free article: PMC7195692] [PubMed: 32356863].
12. Liu D, Ahmet A, Ward L, Krishnamoorthy P, Mandelcorn ED, Leigh R, Brown JP, Cohen A, Kim H: A practical guide to the monitoring and management of the complications of systemic corticosteroid therapy. Allergy Asthma Clin Immunol 2013; 9:30Crossref, Medline, Google Scholar.
13. Schug SA, Palmer GM, Scott DA, Halliwell R, Trinca J. Acute pain management: Scientific evidence fourth edition,2015. Med. J Aust. 2016;204(8):315–17.
14. Chou R, Gordon DB, de Leon-Casasola OA, et al. Guidelines on the management of postoperative pain management of postoperative pain: A clinical practice guideline from the American Pain Society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists. J Pain. 2016;17(2):131–57.
15. Vallejo R, Barkin RL, Wang VC. Pharmacology of opioids in the treatment of chronic pain syndromes. Pain Phys. 2011;14(4):E343–E360360.
16. Hooper DC. Mode of action of fluoroquinolones. Drugs. 1999;58(Suppl 2):6e10.
17. Davies B. E. Pharmacokinetics of oseltamivir: an oral antiviral for the

- treatment and prophylaxis of influenza in diverse populations. *The Journal of antimicrobial chemotherapy*. 2010; 65 Suppl 2(Suppl 2), ii5–ii10. <https://doi.org/10.1093/jac/dkq015>
18. Alahdal M, Al-Shabi J, Ogaili M, Abdullah QY, Alghalibi S, Jumaan AO, AL-Kamarany MA. Detection of dengue fever virus serotype – 4 by using One-Step Real-Time RT-PCR in Hodeidah, Yemen. *Microbiology Research Journal International*. 2016;14(6):1–7. DOI:<https://doi.org/10.9734/BMRJ/2016/24380>
 19. Al-Areeqi A, Alghalibi S, Yusuf Q, Al-Masrafi I, Al-Kamarany MA. Epidemiological characteristic of malaria coinfecting with dengue fever in Hodeidah, Yemen. *International Journal of TROPICAL DISEASE & Health*, 2020; 40(3):1–10. DOI:<https://doi.org/10.9734/ijtdh/2019/v40i330230>
 20. Yusuf QA, Ogaili M, Alahdal M, Amood Al Kamarany M. Dengue fever infection in Hodeidah, Yemen risk factors and socioeconomic indicators. *British Biomedical Bulletin*. 2015;3(1):58–65.
 21. Yusuf Q, Al-Masrafi I, Al-Mahbashi A, Al-Areeqi A, Al-Kamarany MA, & Khan AS. First evidence of West Nile Virus in Hodeidah, Yemen: Clinical and epidemiological characteristics. *International Journal of TROPICAL DISEASE & Health*. 2019;38(4):1-9. DOI:<https://doi.org/10.9734/ijtdh/2019/v38430190>
Available:<https://journalijtdh.com/index.php/IJTDH/article/view/30190>
 22. Al Kamarany MA, Majam A, Suhail K, Zuhairy A, Alabsi E. Coronavirus Disease 2019 – Dengue Fever Coinfection: A case report. *International Journal of Pathogen Research*. 2023;12(4):27–32. Available:<https://doi.org/10.9734/ijpr/2023/v12i4234>
 23. Fuad Ahmed Balkam. Hematological changes among patients with dengue fever. *Abhath Journal of Basic and Applied Sciences*. 2022;1(1):72–82. Available:<https://doi.org/10.59846/ajbas.v1i1.362>
 24. Malik MR, Mnzava A, Mohareb E, Zayed A, Al Kohlani A, Thabet AA, El Bushra H. Chikungunya outbreak in Al-Hudaydah, Yemen, 2011: Epidemiological characterization and key lessons learned for early detection and control. *J Epidemiol Glob Health*. 2014;4(3):203-11. DOI: 10.1016/j.jegh.2014.01.004. Epub 2014 Mar 15. PMID: 25107656; PMCID: PMC7333817.
 25. Rezza G, El-Sawaf G, Faggioni G, Vescio F, Al Ameri R, De Santis R, Helaly G, Pomponi A, Metwally D, Fantini M, Qadi H, Ciccozzi M, Lista F. Co-circulation of Dengue and Chikungunya Viruses, Al Hudaydah, Yemen, 2012. *Emerg Infect Dis*. 2014;20(8):1351-4. DOI: 10.3201/eid2008.131615 PMID: 25061762; PMCID: PMC4111199.
 26. AL Kamarany MA, Al-Masrafi I, Ghouth AB, Suhail K, Majam A, Zuhairy A, Alabsi E. Outbreak investigation: First Ten COVID – 19 Infection related deaths in Hodeidah, Yemen. *Asian Journal of Research in Infectious Diseases*. 2023; 13(3):19–27. Available:<https://doi.org/10.9734/ajrid/2023/v13i3268>
 27. AL-Kamarany MA, Suhail KA, Majam AS, Abdulabari Alabsi E, Hamoud Dowbalah M, Mohammed Zuhairy A. Epidemiological and clinical features of COVID-19 in Hodeidah, Yemen. *International Journal of TROPICAL DISEASE & Health*.2021; 42(21):28–40. Available:<https://doi.org/10.9734/ijtdh/2021/v42i2130550>
 28. Suhail FA, Al Kamarany MA. Radiological features of COVID-19 patients in Hodeidah, Yemen. *Asian Journal of Research in Infectious Diseases*. 2021; 8(4):117–127. Available:<https://doi.org/10.9734/ajrid/2021/v8i430256>
 29. AL Kamarany MA, Abdulkarim T, & Nasser M. Biomolecular Features of COVID-19 in Hodeidah, Yemen. *Asian Journal of Biochemistry, Genetics and Molecular Biology*, 2023;14(3):29–38. Available:<https://doi.org/10.9734/ajbgmb/2023/v14i3317>
 30. Al-Kamarany MA, Abdulkarim T. Immunological Features of COVID-19 in Hodeidah, Yemen. *Asian Journal of Immunology*. 2023;6(1):76–84. Available:<https://journalaji.com/index.php/AJI/article/view/97>
 31. Balkam FA, Al-Kamarany MA. Hematological and coagulation features of COVID-19 patients in Hodeidah, Yemen.

- International Journal of Pathogen Research. 2023;12(6):46–54.
Available:<https://doi.org/10.9734/ijpr/2023/v12i6252>
32. Balkam FA, Al-Kamarany MA. Hematological and coagulation features of COVID-19 patients in Hodeidah, Yemen. International Journal of Pathogen Research. 2023;12(6): 46–54.
Available:<https://doi.org/10.9734/ijpr/2023/v12i6252>
 33. World Health Organization (WHO). Operation consideration for case management of COVID-19 Health Facility;2020.
 34. Ministry of Public Health and Population (MOPHP) of , Syrian National Guideline for Control of Coronavirus Disease 2019 (COVID-19);2020.
 35. Meador CL. Monitoring response to drug treatment , best tests , know the abnormality that you are going during treatment . Pick something you can measures , Lyons IARC Press;1999 .
 36. Vis JY, Huisman A. Verification and quality control of routine hematology analyzers. International Journal of Laboratory Hematology. 2016;38(Suppl. 1):100–109
 37. Saeed AL-Shaibany, Salma Taha, Adel Dhfash, Fouad Balkam, and Mohammed Amod AL-Kamarany. Validation hematological analyzer for assay of erythrogram in Hodeidah City, Yemen. Asian Hematology Research Journal . 2018;1(1):1
 38. Centers for Disease Control and Prevention (CDC) . Information for Clinicians on Investigational Therapeutics for Patients with COVID-19;2020.
 39. RECOVERY Collaborative Group; Horby P, Lim WS, Emberson JR, Mafham M, Bell JL, Linsell L, Staplin N, Brightling C, Ustianowski A, Elmahi E, Prudon B, Green C, Felton T, Chadwick D, Rege K, Fegan C, Chappell LC, Faust SN, Jaki T, Jeffery K, Montgomery A, Rowan K, Juszczak E, Baillie JK, Haynes R, Landray MJ. Dexamethasone in Hospitalized Patients with Covid-19. N Engl J Med. 2021;25;384(8):693-704.
DOI: 10.1056/NEJMoa2021436. Epub 2020 Jul 17.
PMID: 32678530; PMCID: PMC7383595.
 40. Ledford H. Coronavirus breakthrough: Dexamethasone is first drug shown to save lives. Nature. 2020;582(7813):469.
DOI: 10.1038/d41586-020-01824-5.
PMID: 32546811.
 41. RECOVERY trial news release: Dexamethasone. Oxford University News Release. EMBARGOED UNTIL;2020
 42. Villar J, Ferrando C, Martinez D, et al. Dexamethasone treatment for the acute respiratory distress syndrome: A multicentre, randomised controlled trial. Lancet Respir Med 2020;8(3): 267-76.
 43. Horby P, Lim WS, Emberson J, et al. Effect of Dexamethasone in Hospitalized Patients with COVID-19: Preliminary Report. medRxiv 2020:2020.06.22.20137273.
 44. Cavalcanti AB, Zampieri FG, Rosa RG, et al, Coalition Covid-19 Brazil I Investigators. Hydroxychloroquine with or without azithromycin in mild-to-moderate Covid-19. N Engl J Med2020.
DOI:10.1056/NEJMoa2019014
PMID:32706953Abstract/FREE Full TextGoogle Scholar.
 45. Kollias A, Kyriakoulis KG, Dimakakos E, Poulakou G, Stergiou GS, Syrigos K. Thromboembolic risk and anticoagulant therapy in COVID-19 patients: Emerging evidence and call for action. Br. J. Haematol. 2020;189(5):846-847.
 46. World Health Orgnization (WHO), WHO discontinues hydroxychloroquine and lopinavir/ritonavir treatment arms for COVID-1. WHO discontinues hydroxychloroquine and lopinavir/ritonavir treatment arms for COVID-19;2020.
 47. Gautret P, Lagier JC, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. Int J Antimicrob Agents 2020 March 20 (Epub ahead of print).
 48. Marciniak K, Beberok A, Boryczka S, et al. Ciprofloxacin and Moxifloxacin Could Interact with SARS-CoV-2 Protease: Preliminary in Silico Analysis.
Available at SSRN:
<https://ssrn.com/abstract53562475>. 2020
 49. Brooks SK, Webster RK, Smith LE, et al. The psychological impact of quarantine and how to reduce it: Rapid review of the evidence. Lancet 2020;395(10227):912-920.
DOI: 10.1016/S0140-6736(20)30460-8
 50. Wang C, Pan R, Wan X, et al. Immediate psychological responses and associated factors during the initial stage of the 2019 Coronavirus Disease (COVID-19) Epidemic among the general population in

- China. Int J Environ Res Public Health. 2020;17(5):E1729.
DOI: 10.3390/ijerph17051729
51. Food and Drug Administration – United States (FDA 0- US). Coronavirus Treatment Acceleration Program (CTAP) , 2023.
52. García-Lledó A, Gómez-Pavón J, González Del Castillo J, Hernández-Sampelayo T, Martín-Delgado MC, Martín Sánchez FJ, Martínez-Sellés M, Molero García JM, Moreno Guillén S, Rodríguez-Artalejo FJ, Ruiz-Galiana J, Cantón R, De Lucas Ramos P, García-Botella A, Bouza, E. Pharmacological treatment of COVID-19: an opinion paper. Revista española de quimioterapia: publicación oficial de la Sociedad Española de Quimioterapia. 2022;35(2):115–130.
Available:<https://doi.org/10.37201/req/158.2021>

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/111318>