



An Assessment of the Nutritional Status and its Predictors amongst Women in a Rural Community of Dharwad Taluk, Karnataka, India

Vinutha U. Muktamath ^{a++*}, Sannapapamma K.J ^{a#}, Ravi Y ^{a†},
Geetha Channal ^{a‡}, Rajeshaari Desai ^{a†}
and Sumangala Badami ^{a^}

^a All India Coordinated Research Project-WIA, University of Agricultural Sciences, Dharwad, India.

Authors' contributions

This work was carried out in collaboration among all authors. Authors VUM and RY contributed to data analysis. Author VUM wrote the manuscript. Author SKJ provided all the facilities to conduct the research. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/EJNFS/2024/v16i41405

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113501>

Original Research Article

Received: 09/01/2024

Accepted: 13/03/2024

Published: 27/03/2024

ABSTRACT

The connection between food insecurity and health is becoming more widely acknowledged, as it plays a major role in the onset and advancement of non-communicable illnesses. The family's overall well-being and health are directly impacted by the health status of the female. Rural women play a dominant role in the country's economy and their nutritional status has direct bearing on the

⁺⁺ Assistant Professor & Scientist;

[#] Professor & Head;

[†] Assistant Professor & Senior Scientist;

[‡] Associate Professor & Senior Scientist;

[^] Young Professional-1;

*Corresponding author: Email: vinumuktamth@gmail.com;

whole family. Hence the present study was conducted with the objective to assess nutritional status and to identify the predictors of poor nutrition among rural women of Dharwad taluk, Karnataka. The sample of the study included 500 rural women in the age group of 16-80 years. A self structured questionnaire, SES Agarwal 2005, Anthropometric measurements were used to assess BMI, Hb level, knowledge level and morbidity status. More than one third of the rural women were selected were in the age group of 40-80 years, were illiterates and involved in agriculture. Majority belong to social tribes and staying in nuclear families. Two third of the selected rural women consume non-vegetarian food. More than half of the families belonged to upper middle socio economic status. Majority of them had health related problems and were in mild to moderate anemia category. Low SES, low nutritional knowledge, non working status, morbidity condition, middle age group, anemia status and poor protein diet were found to be the risk factors of poor nutritional status among rural women. Low SES was found to be 2.6 times riskier; anemia condition was 4.68 times risk when compared to normal Hemoglobin level.

Keywords: *Nutritional status; SES; anemia; morbidity; knowledge.*

1. INTRODUCTION

India's economy greatly depends on agriculture, and women make up the majority of those working in the field from planting to harvesting. Based on data from the Agriculture Census (2010–11), women made up 30.3% of the estimated 118.7 million cultivators and 42.6% of the 144.3 million agricultural laborers. Of all female primary laborers, 24% were cultivators and 55% were agricultural laborers. Increased "casualization" of work, unprofitable crop production, and men's distress migration "for higher casual work in agriculture and non-agriculture sectors" have all contributed to the feminization of agriculture in India. Women are increasingly responsible for caring for the family land when men are not there. Job overload, feminization of poverty, job risks, and family burden are all consequences of feminization in agriculture, and these factors have led to severe physical and mental health issues (<https://agcensus.nic.in/>).

An important factor in poor health is inadequate nutrition, which can also significantly shorten life expectancy and raise the risk of non-communicable diseases (NCDs), obesity, and obesity-related morbidity. The connection between food insecurity and health is becoming more widely acknowledged, as it plays a major role in the onset and advancement of non-communicable illnesses. The family's overall well-being and health are directly impacted by the health status of the female. The entire family is impacted by maternal mortality and morbidity, not just the mother. Most of the time, women manage the home and take care of the children in addition to actively participating in financially rewarding work outside the home or perhaps

making a substantial contribution to the family business, whether it be in industry or agriculture. They are the main players in the rural economy. It suggests that their nutritional and physical well-being is critical to the growth of the country as well as to their own families. It health limits both their economic potential and their ability to care for family members [1,2].

Some of the elements that operate against rural females are the patriarchal social structure, gender discrimination from birth, malnourishment, illiteracy, unskilled labor, hard physical work, early marriage, frequent and multiple pregnancies, and poverty. Female morbidity rates are higher even after they have an advantage over males biologically. Women are triple jeopardized by some of the elements that operate against rural females like the patriarchal social structure, gender discrimination from birth, malnourishment, illiteracy, unskilled labor, hard physical work, early marriage, frequent and multiple pregnancies, and poverty. One of the main reasons for the current state of health is low literacy and ignorance about nutrition and health. According to the 2001 census, only 39.3% of Indian women are literate, making up 52.21 percent of the country's overall literate population. In particular, 30.6 percent of the females in rural areas are literates when compared to the overall 44.70 percent literates [3,4].

Anaemia is one of the most widely spread deficiency disease among women. Anaemia occurs due to deficiency of iron and folic acid. Iron is needed for the formation of haemoglobin that is the carrier of oxygen in our blood. According to the fifth round of the National Family Health Survey (NFHS-5), 57 percent of

women are anaemic as per NFHS-5 in comparison to 53 percent reported by NFHS-4 in all the round-II states and union territories in India [5,6]. In Karnataka 20 per cent of rural women have poor status and 37 per cent are obese. And the anemic rate among women in Karnataka is 59 per cent. And in Dharwad district 16.70 and 18.40 per cent of rural women had low BMI and obese respectively. And the anemic rate is 50.20 Per cent. (NFHS-5 Report). Hence under nutri smart village programme, five villages were adopted to improve the nutritional status of women. In view of this, the following study was conducted with an objective to study the nutritional status of the rural women in Dharwad taluk and its predicting factors.

2. METHODOLOGY

2.1 Population and Sample

The population of the sample was rural women from Dharwad taluk in the age group of 20-80 years. Among the 118 villages in Dharwad taluk five villages namely Belur, Dubbanamardi, Kalkeri, Kumbarkoppa and Kogelgeri were selected purposively as they were adopted under "Nutri Smart Villages" programme by ICAR- India Co-ordinated research Project-Women in Agriculture, UAS, Dharwad. The sample consisted of 500 women (100 from each village) in the age group of 20-80 years who were selected randomly.

2.2 Tools Used for the Study

An in-depth detailed self-structured questionnaire was used to collect the demographic information, personal profile, clinical symptoms and morbid condition of rural women. Anthropometric measurements like height and weight of women was used to calculate Body Mass Index(BMI) to assess nutritional status of women. The BMI (Body Mass Index) of mothers was calculated using the formula weight in kg/ (height in meter)² and was classified according to WHO BMI classifications [7]. To assess anemia, the women's hemoglobin (Hb) level was tested by drawing blood sample and testing with help of medical practitioners of local government hospital. Socio-economic status was measured using socio-economic status scale developed by Aggarwal et al. [8]. The nutritional knowledge, attitude and practices of rural women were studied using the standard checklist developed by ICAR- AICRP-WIA in the year 2022.

2.3 Research Design

An exploratory design was used to assess the nutritional status, anemia levels and nutritional knowledge, attitude and practices, clinical symptoms and morbid condition of rural women. A correlational design was used to study the factors influencing the nutritional status of rural women.

3. RESULTS AND DISCUSSION

It is clear from the table (Table 1) that, 23.60 per cent were in the age group of 20-40 years, while 34.20, 36.00 and 6.20 percent were in the age group of 41-60 years and 61-80 and 15-20 years respectively. Majority of the rural farm women (28.40%) were illiterate, followed by primary educated (26%), middle school educated (15.20%), high school educated (15%) and remaining 9 percent and 6 percent were having intermediate and graduation level of education respectively. Majority of the subjects were housewives (35%), followed by agriculture as a main occupation (34%). Wherein 16.6 percent, 11.4 percent were laborers and had services in Govt/private sectors. Only 3 percent women were involved in small sale business activities [9,10].

Table 2 Indicates family profile of rural women where majority were scheduled tribes (52.2%), followed by OBC (19.2%), scheduled caste (15.4%) and upper caste (13.2%) respectively. With respect to the marital status, 94.60 percent were married, 4 per cent were in widowhood stage, separated were 0.80 percent and 0.60 were single. About 59 per cent and 37.80 per cent rural women hailed from nuclear and joint family, while 3.20 per cent belonged to extended family type [11,12].

With respect to food pattern, majority were non vegetarian (61.20%) and 38.8 percent were vegetarian respectively. About 48.8 percent rural farm women reported that they were landless. Wherein 23.60 percent were small land holders, 13.80 percent were marginal land holders, 9.4 percent were semi-medium land holders, 4.20 percent were medium land holders and only 0.2 percent were large land holders respectively. Majority of the rural farm women belonged to lower middle socio economic status (51.80%), followed by upper middle SES (47.80%) and high SES (0.4%) respectively.

Table 1. Personal profile of the rural women

| Variables | Categories | Number | N=500 |
|------------------|------------------------|---------------|-------------------|
| | | | Percentage |
| Age (Years) | 20-40 | 149 | 29.80 |
| | 41-60 | 171 | 34.20 |
| | 61-80 | 180 | 36.00 |
| | Total | 500 | 500.0 |
| Education | Illiterate | 142 | 28.40 |
| | Primary | 130 | 26.00 |
| | Middle school | 76 | 15.20 |
| | High school | 75 | 15.00 |
| | Intermediate | 45 | 9.00 |
| | Graduation & above | 32 | 6.40 |
| Occupation | Agriculture | 170 | 34.00 |
| | Labour | 83 | 16.60 |
| | Service (Govt/Private) | 57 | 11.40 |
| | Small business | 86 | 17.20 |
| | House wife | 104 | 20.80 |

Table 2. Family Profile of Rural Women

| Variables | Categories | Number | N=500 |
|---------------------|-------------------|---------------|-------------------|
| | | | Percentage |
| Caste | Upper caste | 66 | 13.20 |
| | OBC | 96 | 19.20 |
| | SC | 77 | 15.40 |
| | ST | 261 | 52.20 |
| Marital status | Single | 3 | 0.60 |
| | Married | 473 | 94.60 |
| | Widow | 20 | 4.00 |
| | Divorcee | - | - |
| | Separated | 04 | 0.80 |
| Family type | Nuclear | 295 | 59.00 |
| | Joint | 189 | 37.80 |
| | Extended | 16 | 3.20 |
| Veg /Non vegetarian | Vegetarian | 194 | 38.80 |
| | Non vegetarian | 306 | 61.20 |
| SES | High | 02 | 0.40 |
| | Upper middle | 259 | 51.80 |
| | Lower middle | 239 | 47.80 |

Table 3 describes on clinical symptoms observed among the farm women. With respect to the general appearance, majority were fair (52%), followed by good appearance (24.40%) and poor appearance (23.60%) respectively. About 80 percent informed about hair fall, wherein 30.80 percent found to have white hair and 14.40 percent with dry hair symptoms respectively. Symptoms associated with dark circle and burning eyes (12.20%), while blurred vision was observed among 5.60 percent rural women. Mouth ulcer was found among 26.80 percent rural women, about 29.20 percent had decayed tooth and 9.80 percent found to have decayed and broken tooth respectively. Muscular problems were found among 34.80 percent rural

women. Results are in collaboration with the study conducted by Nayak, J, et.al [13] who explained about the main hazards leading to Eye itching, irritation, breathing, water in eyes, blurred vision etc among farm women were Burning of cut-wood, forest wood, dung cake, agriculture waste and kerosene. Respondents faced back, shoulder, leg, chest pain, crushing finger, etc due to work load and type of work in farm activities. The shoulder pain was highest among respondents. Lack of safety measures in farm machine/ equipment were the source of hazard to farm women due to their loose clothes (saree). Occupational risk factors include static position, forward bending, heavy lifting and carrying, kneeling and vibration in agriculture. The lower

the individual's socioeconomic position, the higher was their risk of poor health status. Social factors like education, employment status, income level, gender, and ethnicities have significant influence on women health disparities [14]. The major health problems expressed by the respondents were back pain, joint pain, leg pain, head ache, dust allergy, swellings in joints, fever, cold and cough, general malaise, chest pain, pain in shoulders, eye problem, hearing problem, respiratory problem, throat infection, skin irritation, nervous problem, abdominal pain, constipation, urinal infection, mouth infection, cuts and wounds, musculo-skeletal disorders Meenakshi et al. [15].

Table 4 indicates nutritional knowledge, attitude and practice among rural women. Majority of the rural farm women found to have low nutritional knowledge (62%), followed by medium nutritional knowledge (30%) and high nutritional knowledge (7.4%) respectively. The results are in line with the study conducted by Jyoti S. et al. [16] which indicated the overall knowledge, attitude and practices regarding utilization of underutilized green leafy vegetables in selected rural women that is 43.12 percent, 46.12 percent and 49 percent respectively. Another study by Hafiz et al. [17] showed that malnutrition affects mother and their infant's growth as both have a positive relationship among them and several factors, such as lack of knowledge of proper nutrition,

lack of awareness of malnutrition, low income, cultural and socio-economic factors. Lack of education, awareness and resources are the most common causes of malnutrition deficiency of adequate nutrients affects not only women but also their infants as it reduces their intelligence level. So there is need to create awareness regarding the nutritional status of women. The study by Geetha et al. [18] assessed Knowledge, Attitude, and Practice (KAP) of rural farm women on nutrition along with personal, socio economic profile. More than one third (34.5 %) of the rural farm women had high knowledge, followed by low (33.5%) and medium (32.0%). Hence, intervention strategies like promotion of Nutri-gardens in Schools, Households and the rural Community, promotion of Nutri-sensitive crops or biofortified crops to address the malnutrition.

It is clear from the Table 5 that more than forty per cent of the rural women were under normal BMI category followed by over weight (28.10 %) and underweight (20.40 %). Only 6.20 per cent of the rural women were under obese BMI category. The trend indicated that overweight and obese together were more than undernourished. Similar results were observed by Prakash [19] where 18.70 per cent of women were suffering with chronic energy deficiency, 42.30 per cent were normal, 27 per cent were overweight and 12 per cent were obese.

Table 3. Clinical symptoms observed among the farm women

(N=500)

| Clinical symptoms * | No. | % |
|----------------------------|-----|-------|
| General appearance | | |
| • Good | 122 | 24.40 |
| • Fair | 260 | 52.00 |
| • Poor | 118 | 23.60 |
| Hair* | | |
| • Hair fall | 402 | 80.40 |
| • Dry hair | 72 | 14.40 |
| • White hair | 154 | 30.80 |
| Eye* | | |
| • Dark circle | 61 | 12.20 |
| • Blurred vision | 28 | 5.60 |
| • Burning eyes | 61 | 12.20 |
| Tongue * | | |
| • Mouth ulcers | 134 | 26.80 |
| Teeth* | | |
| • Decayed tooth | 146 | 29.20 |
| • Decayed and broken tooth | 49 | 9.80 |
| Muscular problems * | 174 | 34.80 |

* Multiple responses

Table 4. Nutritional knowledge, attitude and practice among rural women

| Particulars | N | (%) | N=500 |
|---|-----------|--------|-------|
| I. Nutritional knowledge of farm women | | | |
| Low | 310 | 62.00 | |
| Medium | 153 | 30.60 | |
| High | 37 | 7.40 | |
| Total | 500 | 100.00 | |
| Mean ±SD | 26.2±4.18 | | |

Mesharam [20] also observed that the prevalence of chronic energy deficiency has declined from 52 per cent during 1975–79 to 34 per cent during 2011–12, while that of overweight/obesity has increased from 7 to 24 per cent during the same period.

Table 5. BMI of Rural women

| BMI in kg/ (height in meter)2 | N | % | N=500 |
|-------------------------------|-----|-------|-------|
| Underweight (<18.5) | 102 | 20.40 | |
| Normal (18.50 – 24.90) | 226 | 45.10 | |
| Over Weight (>24.90 -30) | 141 | 28.10 | |
| Obese (>30) | 31 | 6.20 | |

It is clear from the Table 6 that, morbidity status of farm women from last three months was found to be high in case of ailments viz., Fatigue/weakness(68%), Backache (59%), Body pain (46%), Acidity (29.8%), Fever (27.2%), Giddiness (19%), Eye strain (17.2%), Joint pain (16.6%), Cold & Cough (16.2%), Headache (13.8%), Pain in shoulder (8.4%) and Asthma (4.8%) respectively. Results are in line with the study conducted by Pratibha J. [21] Results revealed that prevalence of chronic energy deficiency (CED) was found to be higher among the farm women of Haryana and Uttar Pradesh. Morbidity pattern among women agricultural workers reveals that their ailments were more with poverty and occupation rather than lifestyle. Apart from poverty and deprivation, the insensitivity of the health-care system too adds to the ill health of agricultural women [22]. The study by Hemant Kumar [23] revealed that the number of obese people was more than indicated unhealthy food intake and lack of physical activity. Prevalence of hypertension and diabetes as found to be on lower side.

Fig. 1 indicates anemia status of rural farm women. Majority of the farm women had mild anemia (50.2%), followed by moderate anemia(38.40 %). While only 10.60 percent were found to be normal (12 and above) and 0.8 percent were found to be severely anemic. The study by Jethi R. [13] indicated In India nearly 70

percent of women are estimated to be iron deficient. Iron deficiency can exist without anemia. Iron deficiency Anemia (IDA) is very late manifestation of iron deficiency because iron deficiency is very well tolerated. Anemia does not develop till storage iron is exhausted. Saha [24] conducted cross-sectional study on non-iron deficiency anemia in rural Indian women. Results revealed that two out of every three women are anemic; one out of four were anemic with depleted iron storage. Importantly, two out of five women had anemia but iron storage was sufficient.

Table 6. Morbidity status of farm women (last three months)

| Ailments | No. | Per cent * | N=500 |
|-------------------|-----|------------|-------|
| Body pain | 230 | 46.00 | |
| Backache | 295 | 59.00 | |
| Headache | 69 | 13.80 | |
| Fatigue/ weakness | 340 | 68.00 | |
| Giddiness | 95 | 19.00 | |
| Eye strain | 86 | 17.20 | |
| Fever | 136 | 27.20 | |
| Cold & Cough | 81 | 16.20 | |
| Acidity | 149 | 29.80 | |
| Joint pain | 83 | 16.60 | |
| Pain in shoulder | 42 | 8.40 | |
| Asthma | 24 | 4.80 | |

*Multiple responses

The factors influencing the nutritional status of women depicts (Table 7) that the risk of poor nutritional status significantly increased with increase in age of rural women. When compared to the young women aged 15-20 years age group women, the risk of poor nutritional status was 2.86 times higher among 20-40 years old women, 1.59 times risk among 60-80 years women and 1.16 times risk for 40-60 year old women. Also, SES was found to significantly influence the BMI where the risk of poor nutritional status was high among women belonging to lower middle SES (2.68 times more) when compared to high and upper middle SES groups. The occupational status indicated that the risk of

poor nutritional status was more in home makers (1.1 times) when compared to working women. With respect to nutritional knowledge the women with low nutritional knowledge (6.11 times) and medium knowledge were at high risk of having poor nutritional (3.02 times) status. The Hb levels indicate a significant relationship where the rural women belonging to severe anemia category were 4.5 times high risk, women with mild anemia were 1.27 times and moderate anemia were 1.12 times risk of having poor status when compared to women having normal Hb level. The morbidity status as per the last three months prevalence among rural women was found to 4.27 times risk of poor BMI when compared to women who did not suffer from any morbid conditions. With respect to the eating pattern the

risk was very high for vegetarian women where it was 9 times more than non-vegetarians. Age, SES, nutritional knowledge, anemia status, morbid condition and eating pattern were significantly associated with BMI. Whereas, educational was not found to significantly influence BMI of rural women.

Similar results were observed by Bhandari et.al.,2016 and Geeta et.al.,2022 where they observed that the risk was more among 20 to 35 years women. Prakash [19] observed that Body mass index increased in women with increasing age with an increase waist-hip-ratio and foods low on protein, food frequency where the diets were cereal based with less of vegetables and fruits, low in diversity, and over-dependence

Table 7. Bivariate analysis of factor influencing the nutritional status of women based on BMI N=500

| Variables | Nutritional Status | | OR (95 % CI) | P-Value |
|--|--------------------|-------------|----------------------|---------------------|
| | Poor n (%) | Normal n(%) | | |
| I. Age (Years) | | | | |
| 15-20 | 11(35.50) | 20(64.50) | 1 | |
| 20-40 | 68 (57.60) | 50(42.40) | 2.857 (1.291-6.323) | 0.024* |
| 40-60 | 85(49.70) | 86 (50.30) | 1.155 (0.720-1.853) | |
| 60-80 | 110(61.10) | 70 (38.90) | 1.590 (1.041-2.429) | |
| II. Socio economic status (SES) | | | | |
| High | 01(50.0) | 01(50.0) | 1 | |
| Upper Middle | 143(59.80) | 96 (40.20%) | 1.008 (0.062-16.284) | 0.031* |
| Lower Middle | 130(50.20) | 129(49.80) | 2.677(0.476-12.320) | |
| III. Education | | | | |
| Post graduation | 45(54.20) | 38(45.80) | 1 | |
| Graduation | 42(57.50) | 31(42.50) | 0.528(0.741-1.373) | 0.443 ^{NS} |
| High School | 130(57.50) | 96(42.50) | 0.710(0.413-1.621} | |
| Primary | 52 (48.10) | 56 (51.90) | 1.275(0.719-2.263) | |
| IV. Occupation | | | | |
| Working | 219(55.30) | 177 (44.70) | 1 | 0.035* |
| Home maker | 55(52.90) | 49 (47.10) | 1.102(0.715-1.700) | |
| V. Nutritional Knowledge | | | | |
| High | 11(29.72) | 26(70.27) | 1 | 0.003** |
| Medium | 82(53.59) | 71(46.41) | 3.02(0.897-5.789) | |
| Low | 181(58.39) | 129(41.61) | 6.11(1.273-18.135) | |
| VI. Anemia | | | | |
| Normal | 31(58.50) | 22 (41.50) | 1 | |
| Mild | 120(62.50) | 72(37.50) | 1.266(0.916-2.899) | 0.001** |
| Moderate | 119 (47.40) | 132 (52.60) | 1.12 (0.990-2.906) | |
| Severe | 4 (100.00) | 0 | 4.5 (2.012-5.990) | |
| VII. Morbidity Status | | | | |
| No | 42 (26.42) | 117(73.58) | 1 | |
| Yes | 232 (68.04) | 109(31.96) | 4.27(0.476-13.320) | 0.011** |
| VIII. Vegetarian/Non-Vegetarian | | | | |
| Non vegetarian | 171(17.89) | 135 (44.12) | 1 | |
| Vegetarian | 103(53.09) | 91(46.90) | 9.01(1.273-18.135) | 0.000*** |

*Significant at 5 % level ** Significant at 1 % level *** Significant at 0.001 % level




Fig. 1. Anemia levels of rural women

on locally grown produce. Biswas et.al. 2016 also observed the poor nutritional status was significantly associated with age, marital status, religion, place of residence, educational status, SES and wealth quintile being significant factors affecting nutritional status. Food and dietary habits also impact nutritional status, with milk or curd, pulses or beans, dark green leafy vegetables, eggs, and fish being associated with lower rates of underweight. Upadhyay et.al.,2011 also observed that rural women who had poor nutritional knowledge, anemia, low SES, increased age were the risk factors of poor nutritional status. Ferdoushi et al. [25] and Hasnain et al. [26] also observed that the illness and health related problems were the risk factors of poor nutritional status.

4. CONCLUSION

The present study clearly indicates the rural women had poor nutritional knowledge which was a risk factor for the poor nutritional status irrespective of their educational status. Low SES, non-working status, health related problems, anemia and poor diet pattern with low protein intake were found to be the risk factors of poor nutritional status. Majority of the rural women had mild to moderate anemia despite the action taken by the Government. However, a trend towards overweight and obesity was rise in the rural area was prevalent which is alarming and calls for more strong actions to take up the initiatives to combat poor nutritional status among rural women.

FUTURE SCOPE

The article studies the present nutritional status and its predictor factors. There is further scope to

conduct intervention studies and study its efficacy.

ETHICAL APPROVAL

Ethical approval to conduct the study was obtained by the ethical committee of University of Agricultural Sciences, Dharwad.

ACKNOWLEDGEMENT

Authors greatly acknowledge Indian Council of Agricultural Research- Central Institute of Women in Agriculture (ICAR -CIWA), Bhubaneshwar for providing financial facility to conduct research

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Aiyar A, Rahman A, Pingali P. India's rural transformation and rising obesity burden. *World Dev.* 2021;138.
2. McKay F, John P, Sims A, Kaur G, Kaushal J. Documenting the food insecurity experiences and nutritional status of Women in India: Study Protocol. *Int. J. Environ. Res. Public Health.* 2020;17:3769
3. Annual report. Department of Agriculture, cooperation and farmers welfare, Ministry of Agriculture and farmers welfare, Government of India, Krishi Bhawan, New delhi; 2021.

4. Bhandari S, Sayami JT, Thapa P. Dietary intake patterns and nutritional status of women of reproductive age in Nepal: findings from a health survey. *Arch Public Health.* 2016;74:2 Available:<https://doi.org/10.1186/s13690-016-0114-3>
5. Biswas S, Pramanik KR, Sonowal CJ. Marginalised social groups differentials in nutritional status (BMI) among reproductive-aged women in West Bengal. *BMC Public Health.* 2023;23:842. Available:<https://doi.org/10.1186/s12889-023-15635-6>
6. Biswas S, Pramanik KR, Sonowal CJ. Marginalised social groups differentials in nutritional status (BMI) among reproductive-aged women in West Bengal. *BMC Public Health.* 2023;23:842 Available:<https://doi.org/10.1186/s12889-023-15635-6>
7. WHO, Training Course on child growth assessment, WHO child growth standards interpreting growth indicators, department of nutrition for health and development, published by WHO press, World Health Organization, Geneva, Switzerland. 2006; 20-45.
8. Aggarwal OP, Bhasin SK, Sharma AK, Chhabra P, Aggarwal K, Rajoura OP. A new instrument (scale) for measuring the socio-economic status of a family: Preliminary study. *Indian J. Comm. Medicine.* 2005;30(4):111-114.
9. Government of India, Ministry of Health and Family Welfare, Compendium of fact sheets in India and 14 states/UTIs (Phase-11), National Family Health Survey (NFHS-5) 2019-21, Available:https://main.mohfw.gov.in/sites/default/files/NFHS-5_Phase-II_0.pdf
10. Joshi, Pratibha & Sharma JP, Sharma Nishi, Singh BK, Ahmad, Nafees & Gills, Reshma. Health and nutrition status of farm women in rural India: Case of Indo Gangetic plains. *Ind. J. of Agril. Sci.*, 2018;88:978-984. DOI: 10.56093/ijas.v88i6.80657
11. Kulkarni VS, Kulkarni VS, Gaiha R. Double burden of malnutrition Reexamining the Coexistence of Undernutrition and Overweight Among Women in India. *Int. J. Health Serv.* 2017;47:108–133.
12. Upadhyay S, Kumar AR, Rita Singh Raghuvanshi & B.B. Singh. Nutritional Status and Knowledge of Hill Women on Anemia: Effect of Various Socio-demographic Factors, *Journal of Human Ecology.* 2011;33:1:29-34. DOI: 10.1080/09709274.2011.11906346
13. Nayak J, Singh SP, Moharana G. Occupational health hazard faced by farm women at their Workplaces. DRWA (ICAR), Bhubaneswar. 2013:1-32.
14. Tarar MA, Fatima T, Salik MH, Akhtar S, Khan YN, Sultan T, Ahmad F, Warraich IA, Yasmin S. Health problems faced by female farm workers in rural areas of tehsil Dera Ghazi Khan: A sociological investigation, *J Dow Uni Health Sci;* 2016;10(1):35-38.
15. Meenakshi JR, Panneer S. Occupational health of agricultural women workers in India. *Ind. J. Community Med.* 2020;45: 546-9.
16. Singh Jyoti, NeeruBala, Verma Anisha. Knowledge attitude and practices of rural women regarding underutilized green leafy vegetables in Allahabad district. *International Journal of Applied and Pure Science and Agriculture (IJAPSA),* 2015;01 (9): e-ISSN: 2394-5532.
17. Hafiz Ali Raza, Rana Muhammad Amir, Muhammad Qamar Bilal, Saima Afzal, Muhammad Iftikhar, Mortala Boye, Ayesha Qayyum, Adeela Manzor. Knowledge, attitude and practices regarding malnutrition among rural women in punjab, pakistan. *J Agric. Res.* 2020;58(2):125-131.
18. Geetha K, Geetha M, Yankanchi AB. Sunitha, Sneha Shigihalli. Knowledge, Attitude and Practice (KAP) on nutrition among rural farm women of Chickballapur District of Karnataka, India. *Current Journal of Applied Science and Technology.* 2022;41(35):30-39. ISSN: 2457-1024
19. Prakash, Jamuna. Nutritional status and dietary pattern of Indian rural women with reference to energy intake and expenditure, *Journal of Community nutrition and Health.* 2013;2:1-8.
20. Meshram I, Balakrishna N, Sreeramakrishna K, Rao K, Kumar R, Arlappa N, Laxmaiah A. Trends in nutritional status and nutrient intakes and correlates of overweight/obesity among rural adult women (≥ 18 –60 years) in India: National Nutrition Monitoring Bureau (NNMB) national surveys. *Public Health Nutrition.* 2016;19(5):767-776. DOI: 10.1017/S1368980015002268

21. Renu Jethi, Pratibha Joshi, Nirmal Chandra ML, Roy Kushagra Joshi. Investigation of Nutritional and Health Status of Indian Rural Hill Farm Women. Journal of Community Mobilization and Sustainable Development. 2018;13(1):169-176,
22. Thresia CU. Women workers in Agriculture: Gender discrimination, working conditions, and health status Kerala Research Programme on Local Level Development, Centre for Development Studies; 2004.
23. Hemant Kumar, Sajjan Madappady, Prajna Kini. Morbidity Profile of Women Aged 15-45 Years in Rural Mangaluru, A Cross Sectional Study. Indian J Prev Med. 2020;8(2):49-54.
24. Saha S, Puwar T, Shah K, Pandya A, Wanjari MB, Saxena D, Non-iron Deficiency Anemia in Rural Indian Women: A Cross-Sectional Study, Cureus. 2022; 14(8):e28565.
DOI:10.7759/cureus.28565.
PMID: 36185907; PMCID: PMC9520231.
25. Ferdoushi, Sarmin Akter , Md. Masud Rana , Fatema-Tuz-Zohora and A.N.M. Kawsar, Nutritional Status and Morbidity Pattern of Girls residing at selected female halls in Tangail District, International Journal of Immunological Nursing. 2015;1 (1).
26. Hasnain G, Akter M, Sharafat SI, Mahmuda A. Morbidity patterns, nutritional status, and healthcare-seeking behavior of female garment workers in Bangladesh. Electron Physician. 2014;6 (2):801-7.
DOI: 10.14661/2014.801-807
PMID: 25763149; PMCID: PMC4324275.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/113501>