



# Micro Irrigation in Rice: A Review

**Koyi Jyothsna <sup>a</sup>, Jayadeva, H. M. <sup>a</sup>, B. S. Yenagi <sup>b</sup>  
and Aakash <sup>c\*</sup>**

<sup>a</sup> Department of Agronomy, UAS, GKVK, Bengaluru- 560065, Karnataka, India.

<sup>b</sup> Department of Agronomy, UAS, Dharwad- 580005, Karnataka, India.

<sup>c</sup> Department of Agronomy, R.S.M. (P.G.) College, Dhampur-246 761 (Bijnor) (Affiliated to Mahatma Jyotiba Phule Rohilkhand University, Bareilly, Uttar Pradesh), India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/JSRR/2024/v30i51991

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115400>

**Review Article**

**Received: 01/02/2024**

**Accepted: 04/04/2024**

**Published: 09/04/2024**

## **ABSTRACT**

Rice (*Oryza sativa*) is the most important staple food crop over half of the world's population and it provides 21 per cent of global human per capita energy. In India, rice is widely grown as the second food crop occupying about 43 million hectares of area. Water scarcity is a major constraint in the development of rice farming in some parts of the world. The water required to produce unit kg of rice is as high as 4000-5000 liters under conventional methods of water application (Transplanting under submerged conditions). During the crop growth period of rice, the amount of water usually applied to field is often much more than the actual field requirement because of adopting traditional flood irrigation methods. In order to meet the growing demand for water it is necessary to use water efficiently and reduce the loss of water. One such practice to conserve water is adoption of micro irrigation to a water loving crop like rice to reduce the wastage of water and to increase the water use efficiency. Micro irrigation is defined as the application of small quantities of water directly above and below the soil surface, usually as discrete drops, continuous drops or tiny streams through emitters placed along a water delivery line.

\*Corresponding author: E-mail: [akash.agro10@bhu.ac.in](mailto:akash.agro10@bhu.ac.in);

*Keywords: Rice; drip; sprinkler; water use efficiency; grain yield.*

## 1. INTRODUCTION

“Rice is the staple food for more than half of the world’s population and plays an important role in food security of many countries. It is the staple food for over 65 per cent population in India”. [1]. “Globally rice is cultivated in an area of 165.04 million hectares with a production of 782 m t and with a productivity of 4,679 kg ha<sup>-1</sup> [2]. “In India the rice is being cultivated in 47.8 m ha, which is the largest area among all rice-growing countries with the production of 129.4 m t and the average productivity of 2.80 t ha<sup>-1</sup>. [3]. “90 per cent of the world rice comes from Asia of which 85 per cent of production is used for human consumption and it deserves a special status among cereals as world’s most important wetland crop” [4]. Rice is primarily composed of carbohydrate, which makes up almost 80 per cent of its total dry weight. Raw, long grain white rice is a relatively good source of energy, carbohydrates, calcium, iron, thiamin, pantothenic acid, folate and vitamin E, compared to maize, wheat and potatoes.

“Rice is a semi aquatic plant which requires 2-3 times higher amount of water than any other cereal crop when grown under transplanted conditions” [5]. “Among the major constraints affecting rice production across the globe, water scarcity appears to be a major challenge. More than 80 per cent of the fresh water resources in Asia are used for agriculture of which about half of the total irrigation water is used for rice production. Water is a very fundamental resource in agriculture and it is more or less the very essence of most if not all agricultural practices. However, this “liquid gold” is a resource that is gradually becoming scarce and expensive to access and as a result of the impending needs, combined water saving mitigation strategies in all sectors ought to be enforced as a means to minimize water wastage and maximize its output when used”. [6]. “The estimated water availability for agriculture which is 83.3 percent of total water used today will shrink to 71.6 percent in 2025 and to 64.6 percent in 2050” [7]. “The shrinking water resources and competition from other sectors, the share of water allocated to irrigation is likely to decrease by 10 to 15 percent in the next two decades. Hence, the expansion of irrigation may depend upon the adoption of new systems such as pressurized irrigation methods with the limited water resources. Water input to paddy fields is largely used for saturating fields,

performing puddling operation, maintenance of water film, and meeting for seepage, percolation, evaporation and transpiration losses. High water consumption is mainly attributed to the water lost by soil evaporation (E), by crop transpiration (T) and by deep percolation (D) below the rooting zone consequently to the hydrostatic load of a few saturated centimeters of shallow soil. On average, an ET loss from rice fields is about 4-5 mm per day during wet months and about 6-7 mm per day during dry months; this can be as high as 10-11 mm per day in subtropical provinces. It was projected that evaporation alone accounts for about 30-40 per cent of ET” [8]. “Seepage and percolation losses of water account for 1-5 mm per day in heavy clay soils and 25-30 mm per day in sandy and sandy loam soils. It is estimated that the combined losses through seepage and percolation maybe 25-50 per cent of total water loss in heavy soils with shallow groundwater table (20-50 cm depth); and 50-85 per cent of total water loss in coarse textured soils with deep groundwater table (1.5 m depth or more). Such an alarming rate of declining ground water is urging scientists and grower to choose new approaches for increasing the water productivity of rice. The amount of water lost by transpiration is the only loss of water that is involved in crop production because transpiration is a driven process for growth and yield components formation. The reduction in E and D components is possible through the increase of the irrigation efficiency, which can be achieved through the micro irrigation which could allow to grow rice in areas with limited water availability and in soils which are not suitable for basin irrigation. It is an alternative method for rice to save water, to simplify crop management, to reduce environmental impact and to make possible the traditional crop rotation for rice”. [9].

“Traditional method of irrigation not only consumes more water, but also causes severe wastage of water and nutrients thus leading to nutrient losses under anaerobic condition” [10]. “In arid countries with large population growth and reduction of fresh water, there is considerable pressure on the agricultural sector to reduce water consumption and access to fresh water for industrial and urban sectors” [11]. “The agricultural sector faces a serious challenge of producing more food with as little water as possible, which can be achieved by increasing crop water productivity” [12]. “Demand for rice is

growing every year and it is estimated that in 2025 AD the requirement would be 140 m t. Decreasing water availability for agriculture threatens the productivity of the irrigated rice ecosystem and ways must be sought to save water and increase the water productivity of rice" [13]. "Because of the increasing demand for food with the increasing population growth and scarcity of water, rice producers face three major challenges viz., i) To face water scarcity, ii) To increase productivity, iii) To produce more rice with less water. Hence, it is necessary to develop an alternative rice production system that requires less water and increases water productivity. The traditional flooding irrigation in rice cultivation system has the problems of high water consumption and low water use efficiency (WUE), so the adoption of an alternative water-saving rice production system is necessary to tackle the problems of water scarcity" [14]. Rice cultivation under micro irrigation is the most effective technology to improve the efficiency of irrigation water applied and to enhance the water use efficiency especially in arid areas. Micro irrigation (Drip and sprinkler) is one such tool that can reduce water use and at the same time increase yield. It is the time to save water by adopting micro irrigation systems.

The aim of this review is to highlight the importance of adoption of micro irrigation system for the cultivation of rice which is a staple food crop of the country and the importance of conservation of water by increasing water use efficiency due to precise application of water under the current scenario of growing water scarcity due to climatic aberrations and weather anomalies.

## 2. INFLUENCE OF MICRO IRRIGATION ON GROWTH PARAMETERS

"In an experiment conducted in Bangalore, Karnataka, pooled data of 3 years showed that significantly higher plant height, number of tillers per plant, leaf area per plant, leaf area index, total dry matter accumulation were recorded with drip irrigation scheduled at  $2.0 E_{Pan}$  throughout the growth stages in aerobic rice as compared to surface irrigated puddled rice which could be attributed to continuous supply of water there by maintaining optimum water and nutrient availability throughout crop growth in drip irrigation" [15]. "Drip irrigation at 1.4 IW:CPE ratio recorded significantly higher plant height, number of leaves per hill/plant, leaf area index and effective tillers per plant as compared to other

drip irrigation levels (0.6, 0.8, 1.0 and 1.2 IW:CPE) and sprinkler and flood irrigation in an experiment conducted for 2 years at Chattisgarh" [16]. "Sprinkler irrigation at 125 per cent pan evaporation recorded significantly higher plant height, number of tillers per hill in aerobic rice as compared to other sprinkler irrigation levels (75 % and 100 % pan evaporation) and life saving irrigation treatments" [17]. "Reduction in growth parameters with increase in severity of water stress might be due to anatomical changes in the plant like decrease in cell volume, cell division, cell elongation, intercellular space and thickening of cell wall as reported by" [18]. "Significantly higher plant height, tiller number  $m^{-2}$ , leaf area index and dry matter production per plant was recorded with drip fertigation at 150 per cent  $ET_c$  with 125 per cent RDF over surface irrigation with soil application of 100 per cent RDF in Coimbatore, Tamil Nadu" [19]. "Drip irrigation at 1.5 IW:CPE ratio recorded significantly higher plant height, number of tillers per plant, number of leaves per plant, leaf area, leaf area index and total dry matter production per plant over other levels (0.75, 1.0 and 1.25 IW:CPE ratio) in an experiment conducted in clay soil at Raichur, Karnataka which might be due to maintenance of adequate soil moisture by frequent irrigation which favoured faster cell division and cell elongation ultimately resulting in higher tiller production, more number of leaves and leaf area development and also higher dry matter production" [20]. "In a study conducted at Pantnagar, significantly higher plant height was recorded with Drip irrigation at 100 per cent CPE 2 days gap over the farmers practice (transplanted and flooded rice) by 14.7 per cent which could be due to even distribution of water with adequate availability and enhanced favorable rhizosphere for more nutrient uptake which in turn accelerate the growth, leading to the formation of higher growth attributes in the course contributing more photosynthates towards the sink" [21]. "A pooled data of 2 years study at Tamil Nadu in clay loam soil showed that significantly higher plant height was noticed with sprinkler Irrigation at 150% PE as compared to surface irrigation" [1]. "An experiment was conducted in sandy loam soils of Tanjavur" [22], "Tamil Nadu and showed that significantly higher plant height, leaf area index, dry matter production, crop growth rate and relative growth rate were recorded with drip fertigation at 200% PE + 125% RDF as compared to surface irrigation and other levels of drip fertigation. In an experiment conducted at Raichur, Karnataka, direct seeded rice with drip irrigation at 4 hours

daily recorded significantly higher plant height, leaf area index, number of tillers  $m^{-2}$  and total dry matter per plant as compared to other treatments" [5]. "Aerobic rice with drip fertigation at 150 per cent  $E_{pan}$  recorded better plant height, number of tillers and leaf area as compared to drip fertigation at 100 per cent  $E_{pan}$  and surface irrigation" [23]. "This variation in growth parameters under surface and drip irrigation system was mainly due to the differences in wetting patterns, water distributions in soil and relative water use by crop" [24]. "Drip fertigation @ 100% PE up to 30 DAS+150% PE up to 60 DAS+ 200% PE up to 90 DAS with 125% RDF (25% through NF+75% through WSF) resulted significantly superior growth characters viz., higher plant height, no. of tiller  $m^{-2}$ , total dry matter production  $m^{-2}$ , as compared to other treatments during summer 2022 and 2023" [25]. "In general, the plant ability to grow depends on the availability of moisture and nutrients in continuous manner. The higher plant growth might be due to higher cell elongation and cell division was caused by constant availability of moisture at the root zone and an increase in nutrient uptake by the plant. Higher number of tillers might be due to good crop, root growth and availability of nutrients throughout growth stages which ultimately increased tiller density" [26]. "Dry matter production increased with increased level of water and nutrients which might be due to water soluble fertilizer through fertigation resulted in continues supply of nutrients besides maintaining optimum water availability which lead to higher uptake of nutrients which in turn good soil aeration throughout crop growth period might have favored faster cell division and elongation focused ultimately increased plant height, tiller numbers, more number of leaves and leaf area development leading to maintain total dry matter accumulation" [26].

### 3. INFLUENCE OF MICRO IRRIGATION ON YIELD AND YIELD PARAMETERS

"In an experiment conducted at Jalgaon, Maharashtra, significantly higher number of panicles  $m^{-2}$ , panicle weight, number of grains per panicle, 1000 grain weight and grain yield were recorded with Drip Irrigation at 100% CPE 2 days gap as compared to other treatments which might be due to even distribution of water with adequate availability and enhanced favorable rhizosphere for more nutrients uptake which in turn accelerate the growth, leading to the formation of higher yield attributes in the course contribute more photosynthates towards the sink"

[21]. "Highest grain yield and straw yield was recorded with application of 100 % RDF through drip fertigation with water soluble fertilizers as compared to surface irrigation with soil application of RDF which can be mainly attributed to continuous availability of water and nutrients that resulted in higher uptake of nutrients in turn production of higher dry matter under drip fertigation" [27]. "In Cauvery new delta zone of Tamil Nadu, highest grain yield and straw yield were recorded with drip fertigation at 200 % PE with 125 % RDF as compared to surface irrigation with application of 100 % RDF" [22]. "Significantly highest number of panicles  $m^{-2}$ , spikelets per panicle, filled grains percentage, test weight and grain yield were recorded with lateral distance of 0.8 m, row spacing of 20 cm with dripper flow rate 1 lph + 30% more water on surface as compared to other treatments in Coimbatore, Tamil Nadu" [28]. "An experiment conducted in Sandy clay loam soils of Bangalore, Karnataka showed that significantly higher productive tillers per hill, panicle weight, filled grains per panicle, 1000 grain weight, grain yield and straw yield were recorded with Irrigation at 2.0  $E_{pan}$  throughout growth stages which was found to be on par with irrigation at 1.5  $E_{pan}$  upto tillering and 2.0  $E_{pan}$  from tillering to harvest as compared to other treatments" [15]. "Higher grain yield recorded in these treatments might be due to its superiority in producing higher productive tillers  $hill^{-1}$ , panicle length, panicle weight, thousand seed weight and total number of grains  $panicle^{-1}$ " [29]. "The increase in yield attributes and yield under drip irrigation might be due to efficient water utilization, higher absorption and accumulation of nutrients by crop and maintenance of excellent soil-water-air relationship with higher oxygen concentration in the root zone" [5]. "Significantly highest panicle length, sound grains per panicle, grain yield and straw yield were recorded with drip irrigation scheduled at 1.4 IW:CPE as compared to other drip irrigation treatments, sprinkler irrigation and flooding method in deep black clay soils of Raipur which might be due to sufficient availability of water whereas stress condition reduced grain yield" [16]. "An experiment was conducted in Sandy loam soils of Karnataka and results revealed that significantly highest number of productive tillers per hill, number of panicles  $m^{-2}$ , panicle length, number of grains per panicle, grain yield and straw yield were recorded with drip irrigation at 150 % CPE + 125 % RDF as compared to conventional flooding method of irrigation" [30]. "The increase in yield attributes under drip fertigation might be due to enhanced

availability and uptake of nutrients leading to enhanced photosynthesis, expansion of leaves and translocation of nutrients to reproductive parts compared to conventional method of soil application of nutrients. Higher grain yield and straw yield might be due to better uptake of nutrients by the crop, because of higher root ramification *i.e.* both horizontal and vertical spread resulting in better contact with the soil particles might have helped in higher nutrient content and uptake and might have also been to the fact that higher soil moisture status. Significantly higher number of effective tillers  $m^{-2}$ , number of grains per panicle, panicle length, panicle weight, grain yield, straw yield and harvest index were recorded with SRI method with drip irrigation emitters spaced at 20cm which was found to be on par with SRI method with drip irrigation emitters spaced at 30cm over conventional paddy cultivation under continuous flooding in heavy clay soils at Bhopal" [31]. "In a field study conducted at Pantnagar in Silty clay loam soils grain yield increased to the extent of 35.31 per cent under direct seeded rice with drip irrigation at 20% CPE on 1 day gap over the conventional transplanted rice which might be due to initial higher plant growth in terms of number of shoots, higher shoot heights, LAI, total dry matter yield and higher yield attributing characters" [32]. "Significantly higher number of panicles  $m^{-2}$ , panicle weight, total filled grains per panicle, test weight, grain yield, straw yield and harvest index were recorded with drip irrigation scheduled at 1.5 IW:CPE which was found to be on par with drip irrigation scheduled at 1.25 IW:CPE" [20]. "These improvements with 1.50 IW/CPE ratio might be due to the continuous and uninterrupted moisture supply throughout the crop growth period which resulted in increased moisture besides higher cell division and elongation. The soil water distribution under drip irrigation is vertical in the beginning but later there will be a lateral distribution whereas, distribution remained primarily vertical under surface irrigation" [33]. "This resulted in low soil suction, better water and nutrient uptake by plants under higher IW/CPE of drip irrigation. Higher number of leaves, leaf area and leaf area index were positively contributed to higher dry matter production" [34]. "Direct seeded rice with drip irrigation 4 hours daily recorded significantly higher productive tillers per hill, panicle length, panicle weight, 1000 grain weight and grains per panicle over the other treatments in medium deep black clayey soils of Karnataka" [5]. "Paddy grain yield was highest by 53.4 per cent with application of irrigation water through drip

scheduled at four hours per day over four hours once in every three days. Further, it was highest by 28 and 32.5 per cent over DSR farmer's practice and surface irrigation, respectively. Irrespective of the irrigation schedules drip irrigation treatments gave significantly higher grain yield over surface irrigation and farmers practice. The increase in grain yield was due to improved yield attributes. Intermittent irrigation and maintaining moist, mostly aerobic soils not only enhances tillering but also the root system's development and functioning" [35]. "Yield and yield attributes of rice were improved under drip irrigation than aerobic practice" [36]. "Soil water tension of 10 kPa was adequate to manage the sprinkler irrigation in rice, especially in the reproductive stage and when using cultivars developed for flooded environments at Brazil in lowland rice" [37]. "Significantly highest grain yield was recorded with 0.8 Epan under drip irrigation @ 4 l/h at 3 days interval over the other treatments at West Bengal" [6]. "Results of a research in Pakistan showed that yield of rice increased by 18 per cent by adopting sprinkler irrigation" [38]. "Drip fertigation @ 100% PE up to 30 DAS+150% PE up to 60 DAS+ 200% PE up to 90 DAS with 125% RDF (25% through NF+75% through WSF) resulted significantly higher no. of filled grains panicle<sup>-1</sup>, total no. of grains panicle<sup>-1</sup>, fertility percentage and lower no. of ill filled grains panicle<sup>-1</sup>, grain yield and straw yield as compared to other treatments, during summer 2022 and 2023" [25]. "A sound source in terms of plant height, number of tillers to support and the number of leaves were rationally able to increase the total dry matter and later lead to higher grain yield. Production and partitioning of dry matter are vital for determination of overall yield of the crop. The higher grain yield recorded might be due to higher conducive situation for efficient water and nutrients uptake which boost their growth and yield attributes through supply of more photosynthates towards the reproductive sink. The adoption of sub surface drip irrigation system (SDIS) to grow the rice cultivars Tarm Hashemi (TH) and Daillman Mazandarani (DM), with planting distances of 18, 20, and 22 cm, was compared with the flood irrigation system (FIS). The SDIS achieved water in abundance as reflected in the improvement of all the studied characteristics of crop yield, *i.e.*, plant vigor index, and increased grain yield and harvest index for both rice cultivars. It can refer to an increase in moisture content by increasing the volume of added water and the role of disintegration and non-compact soil, increasing its permeability and thus increasing its

ability to retain irrigation water at the root zone, making it less vulnerable to environmental conditions, such as sunlight and wind, as well as, an increase in water movement at the root zone and an increase in the volumetric distribution of soil pores, which in turn increases the soil ability to hold water and retain moisture” [39].

#### 4. INFLUENCE OF MICRO IRRIGATION ON WATER USE EFFICIENCY

Highest water productivity was noticed with drip fertigation (1.0+1.5 PE) with 75 % RDF through WSF as compared to surface irrigation with soil application of RDF and other drip fertigation treatments. Increase in water productivity was attributed to increased grain yield with decreased water consumption as compared to surface irrigation [29]. A pooled data of two years showed that drip irrigation scheduled at 1.5  $E_{Pan}$  recorded significantly highest water productivity as compared to continuous submergence (Transplanted) [40]. The water requirement was down by 50 per cent by reducing seepage, percolation and evaporation losses in view of ET based irrigation scheduling in aerobic rice [41]. Drip irrigation scheduled at 0.8  $E_{Pan}$  @4 lph at 3 days interval recorded highest water use efficiency as compared to other treatments at West Bengal [6]. Lower amount of water consumption and higher water productivity was recorded with sprinkler irrigation over the other irrigation methods in a 2 year field study at Meerut, Uttar Pradesh in Loamy soils [42]. Highest water use efficiency and 30.7 per cent saving of water over flood irrigation was achieved with drip irrigation in an on farm trial conducted in Haryana. The drip irrigation methods used less water due to restriction of water loss through evaporation from large amount of ground, conveyance losses resulted in maximum water use by crops [43]. In a field experiment conducted at Tamil Nadu, pooled data of 2 years showed that highest per cent water saving over surface irrigation was recorded with sprinkler Irrigation at 125% pan evaporation [1].

#### 5. ECONOMICS

Highest gross returns, net returns and B:C ratio was recorded with drip irrigation as compared to other irrigation methods (Sprinkler and flood irrigation) in a field study conducted at Raipur [44]. The finding of a study indicated that the higher net return was obtained under drip irrigation method followed by flood irrigation method of irrigation. The lowest net return was in

sprinkler irrigation method. It showed that drip irrigation produced 12 per cent more net income than flood irrigation method, and 40 per cent higher net income as compared to sprinkler irrigation system. Further, drip irrigation system proved 0.29 t higher incremental yield than any other method. Similarly, net increment income of drip irrigation system was Rs. 4511 over traditional flood irrigation method [45]. Optimal water use can enhance returns with enhanced labour productivity and far higher net income than traditional methods for the cultivation of rice [46].

#### 6. CONCLUSION

Micro irrigation practices can be adopted to reduce the wastage of water and supply required quantities of water to rice so as to decrease the water crisis because rice is the major consumer of huge quantities of water when cultivated under flooding method of irrigation. Water saving can be achieved, fertilizer requirement can be reduced as less quantities of fertilizer was required when applied through irrigation and energy can be saved. Water use efficiency and fertilizer use efficiency can be increased by adopting a suitable micro irrigation practice based on the soil type, slope of the land and availability of water.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Kumar GS, Ramesh K, Subrahmaniyan, Ravi V, Effect of sprinkler irrigation levels on the performance of rice genotypes under aerobic condition. *Int. J. Curr. Microbiol. App. Sci.* 2018; 7(3):1848-1852.
2. Anonymous, Area, production and average yield, directorate of economics and statistics, department of agriculture and cooperation report, New Delhi, available on the website; 2022. Available:[http:// www.agricoop.nic.in](http://www.agricoop.nic.in)
3. Anonymous, Area, production and average yield, food and agricultural organization; 2023. Available:[http:// www.indiastats.com](http://www.indiastats.com)
4. Rajeshkumar A, Ramdass S, Thirumeninathan S, Influence of drip fertigation in aerobic rice production. *Int. J. Agri. Sci.* 2019;11(4):8824-8827.
5. Prashant RN, Umesh MR, Basavanneppa MA, Ramesh YM, Manjunatha B, Effect of

- irrigation scheduling through surface drip on growth, yield and water saving in direct seeded rice (*Oryza sativa* L.). J. Farm. Sci. 2019;32(1):27-30.
6. Nabanita S, Ghosh U, Biswas KR, Effect of drip irrigation on yield and water use efficiency of summer rice cultivation in pots. J. Pharma. Phytochem. 2018;7(1):37-40.
  7. Yadav JSP, Conservation and managing water resource for sustainable agriculture. J. Water Manage. 2002;10(1-2):1-10.
  8. Tuong TP, Bouman BAM, In water productivity in agriculture: Limits and opportunities for improvement, CABI Publishing, UK. 2003;53-67.
  9. Antonino S, Antonio M, Francesca B. Water use and crop coefficients in sprinkler irrigated rice. Ital. J. Agron. Riv. Agron. 2009;2:47-58.
  10. Balaji ND, Murthy KR, Pushpa K. Yield and yield components of aerobic rice as influenced by drip fertigation. Int. J. Sci. Nat. 2015;6(3):362-365.
  11. Abdelraouf RE, Abuarab ME. Effect of irrigation frequency under hand move lateral and solid jet sprinkler irrigation on water use efficiency and yield of wheat. J. Appl. Sci. Res. 2012;8(11):5445-5458.
  12. Abdelraouf RE, Habbasha S, Taha MH, Refaie KM, Effect of irrigation water requirements and fertigation levels on growth, yield and water use efficiency in wheat. Middle-East. J. Scientific Res. 2013;16:441-450.
  13. Guerra IC, Bhuiyan SI, Tuong TP, Barker R. Producing more rice with less water from irrigated systems. SWIM Paper 5, IWMI/IRRI, Columbo, Sri Lanka. 1998;24.
  14. Monaco F, Sali G. How water amounts and management options drive irrigation water productivity of rice. A multivariate analysis based on field experiment data. Agric. Water Management. 2018;195:47-57
  15. Anusha S, Nagaraju Mallikarjuna GB, Gururaj K. Influence of drip irrigation scheduling on growth and yield of direct seeded aerobic rice (*Oryza sativa* L.). Int. J. Env. Sci. 2015;9(1&2): 329-332.
  16. Anamika S, Rathore AL, Hemlatha Deepti J, Suneel KR, Nandeha KL, Effect of pressurized irrigation systems on productivity, water and energy use efficiency of summer rice. The Ecoscan. 2015;9(1&2):249-254.
  17. Shahanila P, Prameela P, Visalakshi KP, John PS, Aerobic rice in uplands with micro sprinkler irrigation. Int. J. Tropic. Agril. 2016;54(1):79-83.
  18. Adriano S, Bartolomeo D, Cristos X, Andrea M. Antioxidant defenses in olive trees during drought stress: Changes in activity of some antioxidant enzymes. Funct. Plant Biol. 2005;32: 45-53.
  19. Ramdass S Ramnathan SP, Influence of drip fertigation levels on physiological parameters of aerobic rice in western zones of Tamil Nadu, India. Int. J. Curr. Microbiol. App. Sci. 2017;6(4): 2609-2613.
  20. Jagadish, Chittapur BM, Umesh UR, Halepyati AS, Mastana RBG, Vishwanath J. Srinivasa R. GV, Effect of irrigation scheduling and fertigation on performance of drip irrigated direct seeded rice (*Oryza sativa* L.). Int. J. Chem. Stu. 20186(4): 2216-2222.
  21. Bhardwaj AK, Pandiaraj T, Chaturvedi S, Singh CT, Soman P, Bhardwaj RK, Labh B. Growth, production potential and input use efficiency of rice under different planting methods in drip irrigation. Curr. J. App. Sci. Tech. 2018;26(6):1-9.
  22. Karthika N, Ramanathan SP. Effect of drip fertigation on growth, physiological parameters and grain yield of rice grown in Cauvery new delta zone of Tamil Nadu. Int. J. Chem. Stu. 2019; 7(3):2758-2761.
  23. Kumar PV. Optimization of water and nutrient requirement for yield maximization in hybrid rice (*Oryza sativa* L.) under drip fertigation system. M. Sc Thesis submitted to Agricultural college and research institute, Tamil Nadu Agricultural University, Madurai; 2009.
  24. Sharma S, Sanmay KRP, Gokul BR, Soumen B. Effect of drip irrigation and nitrogen fertigation on yield and water productivity of guava. The Bioscan. 2013;8 (3):783-786.
  25. Umil Singh N, Vaiyapuri K, Thavaprakash N, Selvakumar S, Vanitha K. Impact of Irrigation and fertigation levels on growth, yield components and yield of aerobic rice (*Oryza sativa* L.) under Drip System. Indian J. Agric. Res. 2023;57(6):774-779.
  26. Duary S, Pramanik K. Response of aerobic rice to irrigation and nitrogen management in red and lateritic soil of West Bengal. J. Crop. Weed. 2019;15(1):108-113.
  27. Rekha B, Jayadeva HM, Kambali G, Ngaraju GB, Mallikarjuna, Geethakumari A, Growth and yield of aerobic rice grown under drip fertigation. The Ecoscan. 2015 ;9(1&2):435-437.

28. Parthasarathi T, Mohandass S, Senthilvel S, Eli vered, Effect of drip irrigation systems on yield of aerobic rice. *Environ. Ecol.* 2013;31(4A):1826-1829.
29. Gururaj K, Optimization of water and nutrient requirement through drip fertigation in aerobic rice. Ph. D. Thesis. Univ. Agric. Sci., Bengaluru, Karnataka (India); 2013.
30. Balaji ND, Murthy KR, Venkatesh RJS, Water use efficiency and nutrient use efficiency of aerobic rice as influenced by levels of irrigation and drip fertigation. *Adv. Life. Sci.* 2016;5(2): 529-533.
31. Rao KVR, Gangwar S, Keshri R, Chourasia L, Bajpai A, Soni K, Effects of drip irrigation system for enhancing rice (*Oryza sativa* L.) yield under system of rice intensification management. *App. Ecol. Environ. Research.* 2017;15(4):487-495.
32. Singh TC, Prajapati B, Bhardwaj AK, Effect of drip irrigation on growth and yield of direct seeded rice (*Oryza sativa* L.). *Int. J. Chem. stud.* 2018;6(1):161-164.
33. Araujo F, Larry E, Mark A, A comparison study of young thompson seedless grapevines under drip and furrow irrigation. *Scientia Horticulturae.* 1995;60:235-249.
34. Zheng J, Xianjun Lu, Xinlu Jiang, Yonglu Tang. The system of rice ntensification (SRI) for super high yields of rice in Sichuan Basin. In: Proceedings of an International Conference, Sanya, China. 2004;1-4.
35. Yang C, Yang L, Yang Y and Ouyang Z, Rice root growth and nutrient uptake as influenced by organic manure in continuously and alternatively flooded paddy land. *Agricultural Water Management.* 2004;70(1):67- 81.
36. Kato Y, Okami M, Katsura K, Yield potential and water use efficiency of aerobic rice (*Oryza sativa* L.) in Japan. *Field Crops Res.* 2009;113:328-33.
37. Pinto MAB, Parfitt JMB, Timm LC, Faria LC, Concenco G, Stumpf L, Norenberg BG, Sprinkler irrigation in lowland rice: Crop yield and its components as a function of water availability in different phenological phases. *Field Crops Res.* 2020;248:107714,1-6.
38. Gilani A, Absalan S, Jalali S, Behbahani L. The effect of sprinkler irrigation on grain yield, yield components and water use efficiency of rice cultivars under drill-seed cultivation in Khuzestan. *Irri. Sci. Engi.* 2019;42(2):63-73.
39. Merza NAR, Atab HA, Al- Fatlawi, Alsharifi SKA, Effect of irrigation systems on rice productivity. *J. Breed. Genet.* 2023;55 (2): 587-597.
40. Ramulu V, Rao PV, Uma devi M, Kumar AK, Radhika K. Evaluation of drip irrigation and fertigation levels in aerobic rice for higher water productivity. 2<sup>nd</sup> World forum (WIF2): 6-8, November, Chiang Mai, Thailand; 2016.
41. Medley JC, Wilson LT, The use of subsurface drip irrigation in rice. Paper presented in Tenth Annual National Conservation Systems Cotton and Rice Conference Proceedings Book. 2008;33-34.
42. Rajiv K, Rana NS, Sharawat YS, Mishra A, Kumar V, Mahesh K, Gathala Robin K. Dwivedi A, Improvement in water productivity without yield penalty of direct seeded rice under micro irrigation systems and tillage options in Indo - Gangetic plains of India. *Int. J. Pure. App. Biosci.* 2017;5 (1):147-155.
43. Bansal R, Sharma N, Soman P, Singh S, Bhardwaj RK, On-farm drip irrigation in rice for higher productivity and profitability in Haryana, India. *Int. J. Curr. Microbiol. App. Sci.* 2018;7(2):506-512.
44. Hemlata, Effect of drip irrigation and establishment methods on productivity, water use and economics of summer rice. Ph. D. Thesis. Indira Gandhi Krishi Vishwavidyalaya, Chattisgarh (India); 2015.
45. Soman P, Singh S, Bhardwaj AK, Pandiaraj T, Bhardwaj RK. On-farm drip irrigation in rice for higher productivity and profitability in Haryana, India. *Int. J. Curr. Microbiol. App. Sci.* 2018;7(2): 1-7.
46. Richakhanna. Effect of precision nutrient and water management with different sources and levels of fertilizers on rice production. M. Sc. (Agri.) Thesis, Univ. Agric. Sci., GKVK, Bangalore; 2013.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/115400>