



Assessment of Macro and Micro Mineral Compositions at Different Depths of Forest Nursery Soils in Federal University of Agriculture Abeokuta, Nigeria

Ojekunle O.O. ^{a*}, Oyelowo O.J. ^b, Yisau J.A. ^{a,b}
and Majolagbe M.O ^b

^a Department of Forestry and Wildlife Management, Federal University of Agriculture, Abeokuta, Ogun State, Nigeria.

^b Forestry Research Institute of Nigeria, Ibadan, Oyo State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJEE/2024/v23i6556

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/116046>

Original Research Article

Received: 26/02/2024

Accepted: 30/04/2024

Published: 02/05/2024

ABSTRACT

Forest soil is characteristically fundamental in the establishment of healthy seedlings for eventual planting out on the field. It is composed of mineral particles, organic matter, water, air, and numerous microorganisms. Two forest nursery soils at different depths were investigated at the Federal University of Agriculture Abeokuta (FUNAAB) for their mineral compositions. With the aid of a soil auger, twenty soil samples were randomly collected from different points in two sites at different depths (0-15 cm and 15-30 cm, air dried and taken to the laboratory for further analysis.

*Corresponding author: E-mail: ojekunleoo@funaab.edu.ng;

Site 1 and site 2 (frequently cultivated portion for agroforestry bare-root seedlings and intermittently used portion). The obtained data were subjected to descriptive statistics. The composition of mineral elements varied with sites and depths. The level of Potassium was higher at depths 0 – 15 cm and 15 – 30 cm in site 2 than other macro-minerals (Sodium, Calcium and Magnesium) when compared between sites. The order of concentration includes $K > Mg > Na > Ca$. Micro-minerals were significantly higher at soil depth 0 – 15 cm in site 2 than site 1 ($Fe > Zn > Mn > Cu$). At soil depth 15 – 30 cm, levels of copper, zinc and manganese were significantly higher ($p < 0.05$) in site 1 with the following trend $Cu > Fe > Zn > Mn$. The study showed that forest nursery soils varied in their mineral composition along with depth in the studied sites. Site 2 had better physicochemical properties and higher level of some essential minerals needed for good performance of seedling growth at nursery stage.

Keywords: Forest nursery soil; macro-minerals; micro-minerals; soil depth.

1. INTRODUCTION

“A nursery is an area where plants are raised for subsequent planting. It consists of transplant beds, paths, irrigation channels and so on. Forests are significant to life on earth. They purify the air we breathe, filter the water we drink, prevent erosion, and act as an important buffer against climate change. Cultural and spiritual values, biodiversity conservation, nutrient recycling and carbon dioxide sequestration are among other services functions of the forest [1]. Despite the key roles forest play in the world’s environmental and economic health, forests continue to be lost, along with the endangered animals that live in them” [2,3].

“A forest soil is a natural or only slightly distributed materials that took centuries to develop under permanent forest cover. Forest soils are influenced by forest vegetation. A succession of genetic soils layers is present, ranging from the very important surface organic layers down to the mineral parent materials. The persistent depositing of tree litter upon the ground for many years has developed the features surface layers of organic matter found in forest areas. When forests are removed and the land is used for agriculture, the soil structure generally deteriorates and this is evident by reduced pore space, increased bulk density, increased compaction, reduced content of water stable aggregates, and reduced rates of infiltration, has marked effects on surface water runoff, stream flow and sedimentation. Forest soils are important globally for many reasons which includes the relatively large amount of carbon stored in forest soil organic matter” [4].

“The structure of the forest soil directly influences the growth of the forest biodiversity. Organic matter in the forest soil helps improve and

maintain soil structure. In addition to microorganisms of a forest soil has a relatively high population of macro-organisms that favours soil structure and produces many large burrows and channels. Soil structure is defined as the manner in which the primary soil particles (sand, silt and clay) are combined and arranged with other solid soil components to form clumps or aggregates. It is also the arrangement and organization of the particles and substances that constitute soil. The quality of soil structure is measured by the capacity of the soil particles to clump together or aggregate” [5].

“Most forest soils have acidic soil reaction, with pH values lying within 3.0-7.0 though some forest soils have pH values greater than 7.0. The forest soils are generally more acidic (have less pH) than grassland and agricultural soils due to the continuous production of organic acids and CO_2 from decomposing litters. Chemical properties of soils include Cation exchange capacity which is the maximum quantity of total cation that a soil is capable of holding. Soil pH, soil nutrients, soil organic carbon, soil salinity among others” [6]. The aim of this study is to assess the structural characteristic and chemical properties of forest nursery soil in Federal University of Agriculture Abeokuta (FUNAAB) for optimum productivity at the nursery stage.

2. MATERIALS AND METHODS

2.1 Description of the Study Area

The study was carried out at the nursery site of the Department of Forestry and Wildlife Management, Federal University of Agriculture Abeokuta (FUNAAB), Ogun State. This area falls within Latitudes $7^{\circ}N$ and $7^{\circ}58'$ and Longitudes ranges from $3^{\circ}20'E$ to $3^{\circ}37'E$. The area has a tropical climate with a bimodal distribution of

rainfall; it lies within the humid lowland tropical rainfall with two distinct seasons (the wet season from March to October and the dry season from November to February). The mean annual rainfall is about 1113 mm which peaks in July and September. The relative humidity of the area is 82.4% and the average monthly temperature of 35.8°C [7]. The forest nursery is divided into two main parts on the basis of planting stocks (bare-root & container nursery) as well as seedling size (transplant & seedling nursery). The bare root plant section is the study area for this research. This site is divided into two; site 1 (plants grow directly in the nursery soil and the roots are separated from the soil at the time of lifting such as tomatoes. The lifted planting stock is further handled and planted without soil surrounding the roots. Also, it has only seedling beds in which seedlings only are raised, no transplanting is done. We use this bed for some agroforestry practical for growing crops e.g Solanum spp., amaranthus, celosia etc) and site 2 (planting activities are not carried out here; the land is untilled).

2.2 Soil Sample Collection and Laboratory Analysis

Soil samples were collected randomly from ten points at two different depths 0-15 cm and 15-30 cm in each site (site 1 and site 2) at the FUNAAB forest Nursery with soil auger. Site 1 and site 2 represent frequently cultivated for agroforestry bare-root seedling and intermittently used parts respectively. The samples were poured into individual sample bags, appropriately labelled and taken to the laboratory for analysis.

The collected samples were air dried, crushed and sieved through a 0.5 mm mesh size sieve and analyzed. Soil particle size was determined the pipet method [8], soil pH was determined in 0.01M CaCl₂ by using a soil collection solution ratio for 1:2.5 by means of a Phillip analogue pH meter. The soil was determined using the pH meter [9]. The organic carbon content of the soil was determined by the wet oxidation method of Walkley-Black as described by Allison [10]. The total nitrogen content of the soil was determined by Micro Kjeldahl procedure Bremner [11]. C:N was computed as ratio of N:C. Available phosphorus (P) was extracted by the Bray 1 method. The P concentration in the extract was determined colorimetrically by using the Spectronic 20 and absorption was read-off as described by Bray and Kurts [12] and modified by Murphy and Riley [13]. Exchangeable K, Ca and

Mg were extracted using ammonium acetate, K was determined on flame photometer and Ca and Mg by Atomic Absorption Spectrophotometer (AAS). The trace nutrients are copper (Cu), iron (Fe), manganese (Mn) and Zinc (Zn).

The Soil textural class were identified following the ISSS soil texture classification system. The proportion of clay, silt and sand were used to characterize particles size distribution while the soil physiochemical properties such as bulk density, porosity, soil organic matter, pH, CEC, moisture content, and soil organic carbon (SOC) which were considered as indicators of soil quality were determined.

2.3 Statistical Analysis

The data obtained were analysed using analysis of variance (ANOVA) with the Statistical Analysis System [14] computer package at 5% level of significance to determine difference in the treatments effect, while, the means of differences among the treatments were separated using Fisher's Least Significant Difference (F-LSD; $P \leq 0.05$).

3. RESULTS AND DISCUSSION

3.1 Macro-mineral Composition at Different Depths and Sites

Mineral composition of some macro minerals in site 1 and site 2 at different depths. Varied. At 0 – 15 cm and 15 – 30 cm soil depths, level of Potassium was higher in site 2 than other essential minerals (Sodium, Calcium and Magnesium) when compared between sites. Their order of concentration include $K > Mg > Na > Ca$ (Fig. 1).

3.2 Composition of Trace Minerals at Different Depths and Sites

The levels of some trace minerals recorded in the soils of the two sites are shown in Table 1. At 0 – 15 cm soil depth, results showed significantly higher ($p < 0.05$) levels of copper, iron and zinc in the site 2 than the site 1. On the other hand, manganese level was significantly higher ($p < 0.05$) in site 1 than site 2 (Table 1). At soil depth 15 – 30 cm, levels of copper, zinc and manganese were significantly higher ($p < 0.05$) in site 1 than site 2. However, level of iron was significantly higher ($p < 0.05$) in the site 1 than site 2 (Table 1).




Fig. 1. Mineral composition of the two studied depths and sites

Table 1. Trace minerals at different depths and sites

Micro-minerals	Site 1	0 – 15cm Site 2	Site 1	15 – 30cm Site 2
Cu	0.10±0.00 ^b	0.10±0.00 ^b	0.11±0.01 ^a	0.09±0.00 ^c
Fe	6.21±0.01 ^b	6.02±0.03 ^d	6.48±0.09 ^a	6.09±0.02 ^c
Zn	0.21±0.01 ^b	0.20±0.01 ^c	0.22±0.01 ^a	0.19±0.00 ^d
Mn	1.25±0.01 ^b	1.35±0.31 ^a	1.20±0.01 ^c	1.15±0.02 ^d

Means in the same row having similar superscripts are not significantly different ($p < 0.05$)

There were variations in the physicochemical parameters of the forest nursery soils assessed at the two different sites and depths. The particle size distribution (proportion of clay, silt and sand) also varied among soil depths. Generally, the soils in both sites and depths have a good balance of particle sizes which will have the best stability of pore space size and thus the best soil properties for maximum plant growth and productivity. Muoghalu and Awokunle [15] reported a similar trend in the particle size distribution in the Nigerian rainforest region, Omo biosphere reserve, Nigeria. While the texture is a basic soil property, vegetation disparity could have contributed to the variations in particle size distribution. The pH of the soils was slightly acidic. Consequently, availability of nutrients such as Nitrogen and Phosphorus needed by plants will be supported within this soils. This is corroborated by Alloway and Aryes [16] who stated that high value of pH is predictable as most soils in the tropics range from acidic to slightly neutral. The value may also be as a result of buffering effect of soil organic matter

against pH change in addition to the release of basic cations during organic matter decomposition. The percentage of organic matter in the two sites varied but was higher at 0-15 cm depth in site 1. The substantial difference is probably due to the accumulation and decay of leaf litter and root in the soil. Thus, improves soil physical properties by providing organic binding agents, increasing water holding capacity of soils, provides better aeration, lowering soil bulk density, and improving the elasticity and resilience of the whole soil. The decrease in organic matter with depth could be due to the decrease in the abundance of the fine roots with depth; at greater depth, larger diameter roots predominate, and this agrees with the work of Oyedele et al. [17] who stated that organic matter plays a significant role in adsorption reaction in the soil. It is widely known that 90% of soil nitrogen is in organic combination. High total nitrogen value observed in site 1 could be ascribed to the dominance of nitrogen fixing tree species composition such as *Leucaena leucocephala*, *Albizia lebbek* and *Gliricidia*

sepium around the forest nursery and their rate of decomposition. This agrees with the report of Vanlauwe *et al.* [18] and Deng, *et al.* [19] that species with high nitrogen content decompose more rapidly. Available phosphorus levels were higher at 15-30 cm depth in both sites. Higher phosphorus content in the soil may be an indication of excessive use of inorganic fertilizer or routine application of compost manure high in phosphorus in the area. According to Etabo *et al.* [20] and Fageria *et al.* [21] phosphorus availability or unavailability in the soil to plants critically affects the uptake of nitrogen and other nutrients.

In addition, variations in essential mineral composition between and within sites were observed. In site 1 and 2, the trend of micronutrients includes $K>Mg>Na>Ca$ at both 0 – 15 cm and 15-30 cm soil depths. Level of Potassium was higher than other essential elements Sodium, Calcium and Magnesium in site 2 at both 0 – 15 cm and 15 – 30 cm soil depths when compared between sites. The result of Oladoye [22] in a related study agrees with this study, where Calcium and Magnesium decreased significantly with soil depth in all the study site with 0 - 15cm depth varying significantly from 15- 30cm depth. Higher level of trace mineral was recorded in site 2 at 0 – 15 and the trend is as follows; $Fe>Zn>Mn>Cu$. At soil depth 15 – 30 cm, the order of concentration includes $Cu>Fe>Zn>Mn$. As reported by Kirchmamm and Eriksson [23]; and Combatt *et al.* [24], trace minerals which are required for the normal growth of plants vary in their composition in the soil. All the macronutrients assessed in this study varied with depth in each site.

4. CONCLUSION

The study on the assessment of physicochemical properties and mineral compositions of forest nursery soils indicated variations in these parameters at different depths and sites within the same locations. Physicochemical properties such as Bulk Density, Porosity, and Soil Organic Carbon are considered as indicators of soil quality which ensures optimum productivity at forest nursery stage. The proportion of Clay, Silt and sand were used to characterize particle size distribution also varied. There were disparities in mineral composition both micro and macro minerals between the two sites. Site 2 had better physicochemical properties and higher level of some essential minerals needed for good performance of seedling growth at nursery stage.

Future studies should investigate other likely factors which can affect the productivity of soils at nursery stage.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ajewole OI, Popoola L. Monetization of forest services functions for sustainable management. Journal of Environmental Extension, U.I. 2001;1(1):117-125.
2. Oliveira R, Lacerola LD. Produ e compsiq. Ca da serapilheira na Floresta da Tijula (Rz) Revit Brasil Bot. 1993;16:93-99.
3. Epstein HE, Lauenroth WK, Burke IC, Coffin DP. Regional productivities of plant species in the Great Plain of the United States. Plant Ecology. 1998;134:173-195.
4. Lowe KM, Shulmeister J, Feinberg JM, Manne T, Wallis LA, Welsh K. Using soil magnetic properties to determine the onset of Pleistocene human settlement at Gledswood Shelter 1, northern Australia. Geoarchaeology. 2016;31(3):211-228.
5. Lutz HJ, Chandler RF. Forest Soils. Science. 1947;105(2730):453.
6. Wilson MF, Sellers AH. A global archive of land cover and soils data for use in general circulation models. Journal of Climatology. 1985;2:119-143.
7. Aiboni VU. Characteristics and classification of soils of a representative topographic location in the University of Agriculture, Abeokuta (UNAAB), Nigeria. ASSEST Series A. 2001;1(1):35-50.
8. Gee GW, Bauder JW. Particle-Size Analysis. In: Klute A, Ed., Methods of Soil Analysis, Part 1. Physical and Mineralogical Methods, Agronomy Monograph No. 9, 2nd Edition, American Society of Agronomy/Soil Science Society of America, Madison, WI. 1986;383-411.
9. Black CA. Methods of Soil Analysis II. Chemical and Microbiological properties. Madiso, Wisconsin, American Society of Agronomy. 1965;9(2):1572.
10. Allison LE. Organic Carbon, In: Black CA (ed.) methods of soil analysis part 2 chemical and microbiological properties, American Society Agronomy, Madison, Wisconsin. 1965;1376-1378.

11. Bremner JM. Inorganic Forms of Nitrogen. In: Black CA, et al. Eds., Methods of Soil Analysis, Part 2, and Agronomy Monograph No. 9, ASA and SSSA, Madison. 1965;1179-1237.
12. Bray RH, Kurtz LT. Determination of total organic and available form of P in soil. Soil Science. 1945;59:39-45.
13. Murphy J, Riley JP. A modified single solution method for determination of phosphate in natural waters. Analytica Chimica Acta. 1962;27:31-36.
14. SAS. SAS Institute Inc. SAS/STAT® 14.1 User's Guide. Cary, NC: SAS Institute Inc; 2015.
15. Muoghalu JI, Awokunle HO. Spatial patterns of soil properties under tree canopy in Nigerian rainforest region. Tropical Ecology. 1994;35(2):219-228.
16. Alloway BJ, Ayres DC. Chemical Principles of Environmental pollution, Blackie Academic and Professional. 1997; 53-359.
17. Oyedele DJ, Gasu MB, Awotoye OO. Changes in soil properties and plant uptake of heavy metals on selected municipal solid waste dump sites in Ile-Ife, Nigeria. African Journal of Environmental Science and Technology. 2008;3(5):107-115.
18. Vanlauwe B, Diels J, Songinga N, Merckx R. Residue quality and decomposition; an unsteady relationship? In: Cadisch G, Giller KE. (Eds.) Driven by native plant litter quality and Decomposition. 1997;157-166.
19. Deng L, Huang C, Kim DG, Shangguan Z, Wang K, Song X, Peng C. Soil GHG fluxes are altered by N deposition: New data indicate lower N stimulation of the N₂O flux and greater stimulation of the calculated C pools. Global Change Biology. 2020;26(4): 2613-2629.
20. Etabo EM, Wekha WN, Korir NK, Gweyi-Onyango JP. Effect of phosphorus levels on soil properties and plant tissues of two Nerica varieties. Asian Soil Research Journal. 2018;(3):1-9. DOI: 10.9734/asrj/2018/v1i3682
21. Fageria NK, Baligar C, Clark RB. Micronutrients in crop production, Advances in Agronomy. 2002;77:186 – 268.
22. Oladoye AO. Physiochemical properties of soil under two different depths in a tropical forest of international institute of tropical agriculture, Abeokuta, Ibadan. Journal of Research in Forestry, Wildlife and Environmental. 2021;7(1):41 – 52.
23. Kirchmann H, Eriksson J. Trace elements in crops: Effects of soil physical and chemical properties. In: Glinski J, Horabik J, Lipiec J. (Eds) Encyclopedia of Agrophysics. Encyclopedia of Earth sciences Series. Springer, Dordrecht; 2011. Available:https://doi.org/10.1007/978-90-481-3585-1_175
24. Combatt CE, Jarma OA, Hernandez B, Jaraba NJ, Rodriguez PL. Macroelements and Microelements in soil and their Relationship with the Content of Steviol Glucosides in Stevia rebaudiana Bert from Five Regions of Colombia. Horticulture. 2021;7:547. Available:<https://doi.org/10.3390/horticultur ae7120547>

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/116046>