



# Primary Axillary Tail Malignancy with Nodal Metastatic Disease: A Case Report

**Girish D. Bakhshi <sup>a</sup>, Sumit Boricha <sup>a\*</sup>, Amit Thombare <sup>a</sup>, Chandrakant Sabale <sup>a</sup>, Urvashi Jain <sup>a</sup>, Soumi Das <sup>a</sup> and Ramkishore <sup>a</sup>**

<sup>a</sup> Department of General Surgery, Grant Government Medical College, and Sir JJ Group of Hospitals, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118584>

**Case Report**

**Received: 20/04/2024**

**Accepted: 24/06/2024**

**Published: 04/07/2024**

## **ABSTRACT**

The axillary tail of Spence, usually variable in size is a narrow part of breast tissue near the anterior axillary fold which extends into the axilla via the opening in the clavipectoral fascia. The incidence of malignancy from this tissue is rarely reported in literature. Being in close proximity to axilla, lymph nodal involvement is seen early in the disease process. This may result in an axillary lump presentation with no lump in the breast. A high index of clinical suspicion in the absence of a breast lump creates a diagnostic dilemma. The aim of present case study is to describe the presentation and management of primary malignancy of the axillary tail of the breast.

\*Corresponding author: E-mail: borichasumit5@gmail.com;

**Cite as:** Bakhshi, Girish D., Sumit Boricha, Amit Thombare, Chandrakant Sabale, Urvashi Jain, Soumi Das, and Ramkishore. 2024. "Primary Axillary Tail Malignancy With Nodal Metastatic Disease: A Case Report". *Asian Journal of Case Reports in Surgery* 7 (2):347-51. <https://journalajcrs.com/index.php/AJCRS/article/view/548>.

**Keywords:** Primary axillary; axillary tail; breast lump.

## 1. INTRODUCTION


The incidence of primary malignancy of axillary tail of Spence is 0.1% [1] among all the types of malignancy of the breast. Carcinoma of the axillary tail of Spence is a rare type of breast cancer that develops at a specific anatomical position in the breast [2]. The clinical presentation in such cases may not reveal any lump in the breast. Primary axillary lymphadenopathy in the absence of a breast lump is attributed either to tuberculosis or lymphoma, especially in the absence of inflammation. We present a case of a 39-year-old Indian female who presented with an axillary lump without any symptoms & signs of inflammation. There was no associated lump in main breast tissue. Investigations revealed an axillary lymph nodal mass and a trucut biopsy showed invasive ductal carcinoma. A PET scan revealed Primary in the axillary tail of Spence with secondary axillary lymph adenopathy.

A brief case report with a review of literature is presented.

## 2. CASE PRESENTATION

A 39-years-old female homemaker presented with a lump in the right axilla for 3 months. There was no history of malignancy in 1<sup>st</sup> or 2<sup>nd</sup> degree

relatives. On clinical examination, a 3x3cm firm, the non-tender axillary lymph node was palpable with no lump or swelling palpable in either of the main breast tissues. Examination of contralateral axilla was unremarkable. There were no other significant clinical findings Fig. 1a & b). The patient underwent a sono-mammography which was suggestive of multiple enlarged lymph nodes largest measuring 4.9X2.2X2.8 cms showing eccentric cortical hypertrophy, and preserved fatty hilum hypoechoic with no abnormal calcifications or cystic lesions. The largest necrotic node measured 14X16mm, BIRADS 4. with no lesion in the main Breast tissue. In the left breast, there were incidentally detected two BIRADS 2 lesions one at 1 o'clock position 2 cm from the nipple areola complex and the other were two discrete simple cysts measuring 4X4mm and 4X3mm at 9 o'clock and 10 o'clock position. A USG-guided core tru-cut biopsy was done, histopathological examination revealed invasive ductal carcinoma with Estrogen receptor and Progesterone receptor positive (ER PR +) and Herceptin receptor positive (Her2 positive). Positron emission Tomography (PET) CT revealed primarily in axillary tail of Spence with a few small enlarged FDG avid right axillary nodes. The largest node measures 4.8X2.5cm with SUV max of 4.2. (Fig. 2) No focal FDG avid lesion seen in either breast. Left breast lesions were benign. There were no distant metastases.



**Fig. 1a.** Picture depicting a ovoid lump in right axilla. The curvilinear dotted line indicating the incision line, which is a vertical shaped incision; towards the right breast parallel to the anterior axillary fold. Red arrows depicting the orientation of the breast tissue and axilla




Fig. 1b. Schematic representation of the right axillary lump (red mass) and the incision line (yellow dotted line)




Fig. 2. PET CT of the patient, red arrow depicting Lymph nodal mass in right axilla and axillary tail of spence depicted by blue arrow

A multidisciplinary approach involving medical oncologists, radiologists and surgeons was adopted. The patient was given a choice of Modified radical mastectomy and Breast Conservation Surgery (BCS) explaining pros & cons of both treatments. The patient preferred BCS as a mode of treatment. Hence patient underwent Axillary Lymph node dissection with excision of the axillary tail of Spence which confirmed the diagnosis of Invasive ductal carcinoma with free margins. The patient was then given chemoradiotherapy and was regularly followed up. Follow-up of 1 year has shown her to be disease & symptom-free.

### 3. DISCUSSION

Literature has documented the mean age of occurrence of carcinoma of the axillary tail of Spence to be 55.2 years, with a range of 44-79 years [1]. Notably, a tendency towards stage II or III disease and estrogen and progesterone receptor-negative neoplasms in individuals older than 45 years has been observed [1,3,4]. The present case was a 39-year-old female who was diagnosed with carcinoma of the axillary tail of the breast, presenting with a lump in the right axilla and no lesions in the main breast tissue. This case posed a diagnostic challenge, particularly given the high prevalence of extrapulmonary tuberculosis in the Indian subcontinent, where the axilla is a common site for extrapulmonary tuberculous lymphadenitis after the cervical region [5,6].

The early onset of the disease in this patient underscores the importance of considering carcinoma of the axillary tail of the breast as a differential diagnosis in patients presenting with a lump in the axilla, even in the absence of lesions in the main breast tissue. This emphasizes the need for a multidisciplinary approach in such cases.

Additionally, tailoring the appropriate management steps is crucial, as patient preferences must be considered in the planning process. Patient choice is a fundamental pillar of evidence-based medicine [7]. In the present case, the patient was given the option of breast conservation surgery, aligning with her preferences.

Furthermore, breast carcinoma confined to the axillary tail can be effectively managed using the principles of breast conservation surgery, which aims to achieve optimal patient outcomes. These patients have to be kept on regular follow up.

The limitation of this study is only 1 year of follow up which is less period to predict the long-term outcomes for patients managed with this approach. However, more studies focussed on this approach will help to establish guidelines for this group of patients.

### 4. CONCLUSION

Primary malignancy of axillary tail of the breast poses a clinical dilemma in diagnosis. Histopathology helps in establishing diagnosis. A localised disease on a PET scan can be managed by BCS if the patient opts for that approach. Regular follow-up is the key for success of this approach. Multicentric studies using this approach will help in establishing a standard management guideline for these patients.

### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc have been used during writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology.

#### Details of the AI usage are given below:

1. ChatGPT version 3.0 was used in editing the discussion part of paper only for better rephrasing of the paragraph. We have not used AI anywhere else

### ETHICAL APPROVAL

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

### CONSENT

As per international standards or university standards, patient(s) written consent has been collected and preserved by the author(s).

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Ampil F, Caldito G, Henderson B, Li B, Kim RH, Burton G, Chu Q. Carcinoma of the

- axillary tail of Spence: A case series. *Anticancer Res.* 2012;32:4057-9. PMID: 22993360
2. Kalita P, Gupta A, Sengupta P, Bhattacharyya S, Mishra S, Dasgupta S. Carcinoma of the axillary tail of Spence: A rare case report. *International Journal of Surgery Case Reports.* 2024 Jan 1;114:109151.
  3. Okubo M, Tada K, Niwa T, Nishioka K, Tsuji E, Ogawa T, Seto Y. A case of breast cancer in the axillary tail of Spence - enhanced magnetic resonance imaging and positron emission tomography for diagnostic differentiation and preoperative treatment decision. *World J Surg Oncol.* 2013;11:217. DOI: 10.1186/1477-7819-11-217. PMID: 24004816; PMCID: PMC3844328
  4. Evans DM, Guyton DP. Carcinoma of the axillary breast. *J Surg Oncol.* 1995;59: 190-5. DOI: 10.1002/jso.2930590311. PMID: 7609527
  5. Jerbi M, Hidar S, El Mouedded S, Jenna A, Kerbi S, Cheib A, Khairi H. Le cas Clinique du mois. Tuberculous ganglionaire axillaire une presentation inhabituelle [Tuberculous Axillary Lymphadenitis: An unusual presentation]. *Rev Med Liege.* 2007; 62:188-9. French. PMID:17566386.
  6. Jayabal, Pandiaraja, Arumugam, Shalini. A case of isolated axillary tuberculous lymphadenitis. *Nigerian Journal of Medicine* 2020;29:723-25. DOI: 10.4103/NJM.NJM\_162\_20
  7. Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS. Evidence based medicine: What it is and what it isn't. *BMJ.* 1996;312(7023):71-2. DOI: 10.1136/bmj.312.7023.71. PMID: 8555924; PMCID: PMC2349778

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/118584>