



Assessment of Physico-chemical Properties of Tada Shonga Irrigation Scheme Soil

M. A. Adejumobi ^{a*}, O. E. Onofua ^a and Y. A. Toyin ^b

^a *Department of Agricultural Engineering, Ladoké Akintola University of Technology, Ogbomosho, Nigeria.*

^b *Lower Niger River Basin Development Authority, P. M. B. 1529, Ilorin, Kwara State, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJRAF/2022/v8i130148

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/86190>

Original Research Article

Received 09 February 2022

Accepted 19 April 2022

Published 21 April 2022

ABSTRACT

This research was conducted to analyze the soil physico-chemical properties of Shonga Irrigation Scheme to ascertain its suitability for rice production. The study was carried out in the dry season of 2019. Disturbed soil samples were collected from the 0–15 and 15–30 cm soil layers in five identified areas in the irrigation scheme. The samples were sent to the Soil laboratory of the Lower Niger River Basin for physical and chemical analysis. Most of the soils were strongly acidic to moderately acidic with pH values ranging between 4.8 and 5.7. The organic carbon and organic matter ranged from 0.13 to 0.19% and 0.03 and 0.32%, respectively which implies low fertility. Calcium values ranged between 2.1 and 3.75 cmol/kg, with the low values related to the soil pH status. Magnesium values averaged 2.48 and 1.95 cmol/kg in the top and lower soil layers, respectively. Exchangeable Acidity (EA) values ranged between 0.20 and 3.40 cmol/kg in the surface horizon and between 0.2 and 5.4 cmol/kg in the sub-surface horizons. Available phosphorus had an average of 40.05 and 23.14 ppm in the surface and sub surface soil layers, respectively. Analysis of variance techniques was used for significant differences within chemical properties. For all tests, a threshold of $P=0.05$ was used to define statistical significance. Soil pH and available phosphorus were within the recommended land suitability requirement for rice cultivation. Organic carbon, organic matter, calcium and magnesium were not within the recommended range for rice production. Organic carbon and organic matter can be supplied through the incorporation of organic manure.

*Corresponding author: E-mail: maadejumobi@lautech.edu.ng;

Keywords: Soil properties; irrigation; irrigation scheme; soil samples.

1. INTRODUCTION

Globally, the increase in population has resulted into an increase in global food demand [1]. Hence, food security has become a major global problem. Water and soil resources needed to support agriculture have been under increasing pressure from urban, environmental and industrial users. This has led to the intensification of agricultural land and changing of land use [2]. Alternative strategies to meet the global food demand as well as to maximally utilize the limited land and water resources needs to be put into consideration. Irrigation provides one such solution by opening up more lands for crop production. Irrigation technology makes it possible for food to be produced throughout the whole year even in the absence of rain [3]. But irrigation is not without its challenges. Problems such as alkalization, acidification, salinization and waterlogging could be generated from irrigation and thereby making it hazardous to the soil environment [4].

The continuous use of soils may lead to changes in its physico-chemical properties, influencing both their interaction with the environment and their ability to produce crops. Thus physico-chemical deterioration remains the greatest threat to irrigation farming [5].

Though irrigation provides the needed soil moisture for the absorption of plant nutrients by plant roots, yet it could as well lead to the leaching and erosion of vital soil nutrients. Irrigated arable land is estimated to be around 301Mha worldwide [6]. Without periodic assessment and monitoring, the soil under irrigation operations could be adversely affected and eventually lead to a decrease in crop yield. Therefore this study was carried out to assess the soil physico-chemical properties of Shonga Irrigation Scheme.

2. METHODOLOGY

2.1 Description of the Study Area

The study area is located in Tada Shonga Irrigation Scheme, a distance of 110 km northeast of Ilorin in Edu Local Government Area of Kwara State, Nigeria. Its geographical coordinate is latitude 9° 1' 0" N North and longitude 5° 9' 0" East (Fig. 1). The study was

conducted in the dry season of 2019. The Tada Shonga Irrigation Scheme is operated by the Lower Niger River Basin Development Authority (LNRBDA). The climate in Tada Shonga irrigation sites is tropical continental with pronounced wet and dry seasons and steady high temperatures. The maximum rainfall in the area is during September and drops to zero in December. Its rainy season lasts for about 218 days, it usually starts in April and ends in October [7]. The Tada Shonga Irrigation Project, which is conceived as a pilot public-private partnership, has a proposed gross area of 3000 hectares for the irrigation scheme of which 2000ha is been utilized for rice production with a surface irrigation system. The scheme is designed as a lift irrigation system, involving the direct lifting of water from the River Niger and delivery through a network of canals. The scheme was handed over to LNRBDA by the Kwara State Government in 1978. Surrounding villages benefiting from the project include: Shonga, Tada, Chiji, Yimagi, Kpileagi, Kusogi, Kpataringi, Kochegei, Kpatako and Bafita. About 2.5 million people are serviced by the scheme.

2.2 Sample Collection

Disturbed soil samples were collected at five locations within the irrigation scheme. The five sampling points were labelled M1–M5. Soil samples were collected from the 0–15 (A1) and 15–30 cm (A2) soil layers at each sampling point with the aid of a soil auger. Soil samples were collected according to [8]. The collected soil samples were packed in different polythene bags to prevent atmospheric influences. The samples were sent to the laboratory of the Lower Niger River Basin for physical and chemical analysis.

2.3 Soil Sample Analysis

Soil samples were analyzed according to the USDA textural classes [9]. Physical properties of interest in soil analyses include texture, while chemical properties of concern include pH, organic carbon (OC) and organic matter (OM), exchangeable acidity (EA), exchangeable cations–calcium (Ca) and magnesium (Mg) and available phosphorus (AP). Mechanical analysis of soil was done by the Bouyoucous or hydrometer method [10] by sieve and sedimentation analysis; corresponding textural classes were determined.




Fig. 1. Location of the Tada-Shonga irrigation scheme
 Source: [7]

2.4 Statistical Analysis

Data were statistically analyzed for variance (ANOVA) using the SPSS (v. 20) statistical package. A threshold of $P = .05$ was used to define statistical significance.

3. RESULTS AND DISCUSSION

3.1 Soil Textural Analysis

The soil textural class for M1 was loam in both soil layers. M2 soil was clay loam for both the topsoil and subsoil. M3 and M4 had different soil types in the topsoil and subsoil. M3 was clay loam in the topsoil and silty clay loam in the subsoil. M4 was silty clay loam in the topsoil and clay loam in the subsoil. M5 had both soil layers as clay loam. Clay loam has been reported to be most suited for rice production [11].

3.2 Chemical Properties Analysis

The results of the selected soil chemical properties are presented in Fig. 2. The soil pH

ranged between 4.8 and 5.7 in the surface layer, and 5.0 to 5.5 in the sub-surface soil layer from the five sampling points. The average pH in the 0–15 cm surface soil layer was 5.24 while the sub-surface soil layer recorded 5.3. Soil pH characterizes the chemical environment of the soil and may be used as a guide to the suitability of soils for various pasture and crop species. Soil pH is also an indicator of the chemical processes that occur in the soil and is a guide to likely deficiencies and/or toxicities [12]. According to the classification by [13], the soil pH range in the soil samples indicates a very strong acidic to moderately acidic soil reaction. The relative acidity of the soil samples may be due to the leaching of basic cations, leaving the topsoil acidic as the soils were under irrigation, or due to crop uptake. The soil pH of Tada Shonga Irrigation Scheme falls within the recommended range for rice production (3.1 – 5.3) as stated by [11]. Therefore as regards pH, the five sampling locations of the scheme are suitable for rice cultivation as the scheme is dedicated to rice farming.

Table 1. Soil Textural Classification of Tada Shonga Irrigation Scheme

Sample Point	Sand (%)	Silt (%)	Clay (%)	Textural Class
M1A1	26	44	30	Loam
M1A2	28	54	18	Loam
M2A1	28	32	40	Clay Loam
M2A2	26	32	42	Clay Loam
M3A1	18	46	36	Clay Loam
M3A2	16	44	40	Silty Clay Loam
M4A1	20	48	32	Silty Clay Loam
M4A2	28	44	28	Clay Loam
M5A1	32	36	32	Clay Loam
M5A2	22	46	32	Clay Loam

Organic Carbon (OC) of the soil samples ranged from 0.13 to 0.19%. OC gives a direct measure of available nitrogen in the soil. Organic carbon for the soil is considered adequate if it is within the range of 0.20 – 21.0% for rice cultivation [11]. Hence OC of the soil samples was considered low.

Organic matter (OM) had values between 0.23 and 0.32% in the topsoil and values between 0.03 and 0.3% in the subsoil layer. An average value of 0.28% was recorded in the 0–15 cm depth and reduced to 0.22% at depth 15–30 cm. The OM of the scheme falls below the soil recommended range (0.34–36%) for rice cultivation [11]. The low SOM values in this irrigation site may be due to carbon loss as CO₂ as a result of yearly conventional tillage and removal of crop residue, causing immobilization of soil nutrients. The application of organic manure can help to improve the soil OM content.

Calcium (Ca) values ranged between 2.1 and 3.75 cmol/kg (average of 3.2 cmol/kg) for 0 – 15 cm soil layer and between 2.4 and 3.4 cmol/kg (average of 3.11 cmol/kg) for the 15 – 20 cm soil layer. Apart from sampling locations M2A1 and M5A2, the soil Ca values fall outside the recommended range (2.70 – 0.85 cmol/kg) for rice cultivation [11]. The low Ca could be related to the soil pH status. Acidic soils are usually calcium deficient, indicating that Ca-based fertilizer would be required to reverse this result so that the beneficial use of calcium in the soil including structure stabilization and combating soil acidity would be possible [14].

Magnesium (Mg) values were in the range 0.5–4.35 cmol/kg in the surface soil; and 0.4–3.3 cmol/kg in the sub-surface soil layer. The

average Mg value was higher in the surface layer (2.48 cmol/kg) as compared to the sub-surface layer (1.95 cmol/kg). The Mg values of the scheme soil were not within the recommended range (0.84–0.10 cmol/kg) [12] for rice cultivation. Further application of magnesium is not necessary since the value of the current data is adequate for the soil.

Exchangeable Acidity (EA) values ranged between 0.20 and 3.40 cmol/kg in the surface layers and between 0.2 and 5.4 in the sub-surface layers. According to [15], the range of EA values of the scheme soil were satisfactory for rice cultivation.

The available Phosphorus (AvP) had values ranging from 11.03 to 108.41 ppm, with an average value of 40.05 ppm in the 0–15 cm soil layer. At the 15–30 cm soil layer, AvP values were between 14.4 and 33.08 ppm, with an average value of 23.14 ppm. The AvP in the soil of the scheme (except M4A1) is within the acceptable range (8.60–83.0 ppm) for rice cultivation [11].

3.3 ANOVA of the Chemical Properties

ANOVA for statistical variance are presented in Table 2. There were no significant differences ($p > .05$) in the results between the two soil layers for all chemical properties considered. It is expected that the surface layer or topsoil is more fertile than subsoil, due to having more organic matter and because fertilizers are usually added to the topsoil only. However, the lack of significant differences between the soil layers could be due to erosion of nutrients in the topsoil as a result of continuous irrigation application.




Fig. 2. Soil Chemical Analysis of Tada Shonga Irrigation Scheme

Table 2. Analysis of Variance of Selected Chemical Properties

Chemical Properties		Sum of Squares	df	Mean Square	F	Sig.
Organic Carbon	Between Groups	.003	1	0.003	1.379	0.274
	Within Groups	.019	8	0.002		
	Total	.022	9			
Organic Matter	Between Groups	.010	1	0.010	1.383	0.273
	Within Groups	.056	8	0.007		
	Total	.065	9			
pH	Between Groups	.009	1	0.009	0.110	0.748
	Within Groups	.652	8	0.082		
	Total	.661	9			
Ca	Between Groups	.020	1	0.020	0.071	0.797
	Within Groups	2.292	8	0.287		
	Total	2.312	9			
Mg	Between Groups	.702	1	0.702	0.331	0.581
	Within Groups	16.988	8	2.124		
	Total	17.690	9			
EA	Between Groups	.784	1	0.784	0.179	0.683
	Within Groups	35.040	8	4.380		
	Total	35.824	9			
AvP	Between Groups	713.349	1	713.349	0.899	0.371
	Within Groups	6348.507	8	793.563		
	Total	7061.856	9			

4. CONCLUSION

Physico-chemical properties of Shonga irrigation scheme soil were assessed. Soil pH and available phosphorus were within the recommended land suitability requirement for rice cultivation. Organic carbon, organic matter, calcium and magnesium fell outside the recommended range for rice production. Organic carbon and organic matter can be supplied through the incorporation of organic manure.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Godde CM, Garnett T, Thornton PK, Ash AJ, Herrero M. Grazing systems expansion and intensification: drivers, dynamics, and trade-offs. *Glob. Food Sec.* 2018;16:93–105.
2. Drewry JJ, Carrick S, Penny V, Houlbrooke DJ, Laurenson S, Mesman NL. Effects of irrigation on soil physical properties in predominantly pastoral farming systems: A review. *New Zealand Journal of Agricultural Research.* Taylor & Francis. 2021;64(4):483–507.
3. Khalik AMSA, Moussa KF, Fattah MKA, Abdo AI. Effect of Irrigation Water Source on Some Soil Chemical Properties. *Plant Arch.* 2021;21(2):215–220. DOI:10.51470/PLANTARCHIVES.2021.v21.no2.034.
4. Ayers RS, Wescot DW. *Water Quality for Agriculture.* Food and Agricultural Organization of the United Nations, Rome. *Irrig. Drain. Pap.* 1994;29(29).
5. Gaya UI, Audu AA. Effect of Irrigation on Soil Properties in Jakara Kano, Nigeria,” *Casp. J. Env. Sci.* 2005;3(2):99–106.
6. Pranjali P, Kadiyan N, Chatterjee RS, Kumar D, Sati MS. Interpreting land subsidence impacts due to groundwater depletion using remote sensing-based GRACE gravity anomaly and DInSAR technique: a study on north-western parts of India. *Environ. Earth Sci.* 2021;80(17): 596. DOI: 10.1007/s12665-021-09905-y.
7. LNRDA. Progress report of irrigation scheme. Low. Niger River Basin Dev. Auth. Ilorin, Kwara State. 2021;1–400.
8. Ackerson JP. *Soil sampling guidelines.*
9. USDA, National soil survey handbook. Title 430-VI, Part 647. Washington, DC: Soil Survey Staff, Natural Resources Conservation Service, US Department of Agriculture, Government Printing Office; 1996.
10. Gee GW, Bauder JW. Particle-size analysis. A. Klute (ed.) *Methods of soil analysis.* Part 1. *Agron. Monogr.* 9. ASA and SSSA, Madison, WI. Part. Anal. A. Klute *Methods soil Anal.* Part 1. 2nd ed. *Agron. Monogr.* 9. ASA SSSA, Madison, WI. 1986;383-411.
11. Aondoakaa S, Agbakwuru P. An assessment of land suitability for rice cultivation in Dobi, Gwagwalada Area Council, FCT – Nigeria,” *Ethiop. J. Environ. Stud. Manag.* 2012;5(4):444–451. DOI: 10.4314/ejesm.v5i4.s2.
12. Kalev SD, Toor GS. The Composition of Soils and Sediments,” in *Green Chemistry,* Elsevier. 2018;339–357.
13. USDA-NRCS. *Soil quality concerns, pesticides,* National Soil Survey Center in cooperation with the Soil Quality Institute, NRCS, USDA and the National Soil Tilth Laboratory. *Agric. Res. Serv;* 1998.
14. Tesfai M, Dawod V, Abreha K. Management of Salt-affected Soils in the NCEW” Shemshemia” Irrigation Scheme in Upper Gash Valley of Eritrea. *Drylands Coordination Group;* 2002.
15. KMM, Idoga S. Characterization, classification and suitability ratings of soils for rainfed rice production in Rukubi, Doma, Nasarawa State, Nigeria. *Int. J. Environ. Agric. Biotechnol.* 2018;3(5): 1865–1873. DOI: 10.22161/ijeab/3.5.39.

© 2022 Adejumobi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/86190>