



## Heavy Metals and their Effects on Macroinvertebrates Present in the Ojo River, Lagos, Nigeria

G. A. Reis <sup>a\*</sup>, A. A. Olayeri <sup>a</sup>, A. O. Saba <sup>a</sup>, T. E. Falebita <sup>a</sup>, F. O. Alonge <sup>b</sup>, T. O. Abdulkareem <sup>c</sup>, T. F. Giwa <sup>b</sup>, A. O. Mudashiru <sup>a</sup> and U. O. Jibrin-yekini <sup>a</sup>

<sup>a</sup> Department of Zoology and Environmental Biology, Faculty of Science, Lagos State University, Ojo, Lagos State, Nigeria.

<sup>b</sup> Department of Botany, Faculty of Science, Lagos State University, Ojo, Lagos State, Nigeria.

<sup>c</sup> Department of Biochemistry, Faculty of Science, Lagos State University, Ojo, Lagos State, Nigeria.

### Authors' contributions

This work was carried out in collaboration among all authors. Authors GAR, AAO, AOS and TEF designed the study. Authors GAR, TOA, FOA, AOM, UOJY, AOS, TEF, TFG and AAO conducted the experiments and generated the data. Authors GAR, FOA, AAO, TOA and AOM analyzed the data. Authors FOA, AOM, UOJ and TFG managed the literature searches. Authors GAR, AAO, AOS and TEF drafted the manuscript. All authors read and approved the final manuscript.

### Article Information

DOI: 10.9734/AJEE/2022/v18i230310

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/88399>

Original Research Article

Received 07 April 2022  
Accepted 13 June 2022  
Published 17 June 2022

### ABSTRACT

**Aim:** To provide information regarding the presence of heavy metals in the tissues of crab (*Potamon fluviatile*), prawn (*Macrobrachium rosenbergii*), and crayfish (*Metanephrops australiensis*) obtained from the Ojo river of Lagos State, Nigeria.

**Study Design:** Commercially sold marine crustacean samples (crab, prawn, and crayfish) obtained from the Ojo river of Lagos, Nigeria, were assessed for the presence of heavy metals, and also the potential health risks for local consumers.

**Place and Duration of Study:** Ojo river, located close to Ojo local government secretariat, Ojo, Lagos State, Nigeria.

**Methodology:** Live samples of macroinvertebrates such as *Metanephrops australiensis*, *Potamon fluviatile*, and *Macrobrachium rosenbergii* were purchased from the fishermen at the riverside in Ojo and immediately transferred to the laboratory. The samples were oven-dried and ground into a

\*Corresponding author: E-mail: gbolahanreis@gmail.com;

fine powder, then subjected to sample digestion and finally atomic absorption spectrophotometer (AAS), to obtain the various heavy metal concentrations present in each sample.

**Results:** The result indicated variations in the metal body load among species. All the metals were below the FAO/WHO permissible limit for food consumption except for Cd in crayfish which was slightly beyond the set limit. The highest concentration examined were found in crayfish, followed by crab and prawn. Zinc and iron were of higher concentrations in the tissues of the macroinvertebrates while lead was the least concentrated metal present only in the tissues of crayfish and absent in the tissues of both crab and prawn.

**Conclusion:** The human health risk evaluation for the marine organisms indicated that both the crab and the prawn samples examined were safe for consumption, while crayfish may not be considered safe for consumption. Also, the potential health risk from consuming seafood exposed to these metals should not be ignored.

*Keywords: Heavy metals; crab; prawn; crayfish; crustacean; Ojo river; Lagos; FAO/WHO.*

## 1. INTRODUCTION

The aquatic ecosystem is a water-based environment where interactions between biotic and abiotic components occur. This ecosystem is divided into two types: the “freshwater ecosystem” and the “marine ecosystem” [1] as cited in [2]. The freshwater ecosystem covers about 1% of the Earth’s surface and comprises rivers, lakes, streams, ponds, reservoirs, groundwater, and wetlands. The marine ecosystem is the largest water ecosystem occupying over 70% of the Earth’s surface. This ecosystem is subdivided into the ocean, estuaries (brackish water), salt marshes, coral reefs, mangroves, and algal colonies [2,3].

Marine organisms are known as popular food sources to the inhabitants of the coastal region. Known to possess a high amount of protein and reduced quantity of saturated fats, marine organisms also offer other range of health benefits. However, these benefits could be compromised as a result of contaminants present in the marine ecosystem at levels beyond the tolerance of these organisms [4-6].

Contaminants in the marine ecosystem often remain in either soluble or suspension forms which tend to settle at the bottom of the sea, while some are taken up by organisms in the benthic region of the marine ecosystem. The inland waters exhibit more significant toxic effects arising from the indiscriminate deposition of waste as a result of their relatively small nature compared to the oceans and the seas. This justifies the heavily felt impact of waste deposition in the rivers, streams, lakes, and wetlands [7]. It was argued that about 80% of the pollutants from freshwater are directed into the marine environment [8]. This explains the reason

for the noticeably high level of pollutants in the ocean and low concentrations in many freshwaters.

Heavy metals are chemical elements capable of polluting the marine ecosystem. These metals are a group of metals and metalloids (including their compounds) that are of significant toxicity to the environment or ecotoxic. Heavy metals like manganese, iron, cobalt, copper, zinc, and molybdenum are elements essential to the growth and changes in the life cycle of any organism, however, these metals are toxic at high concentrations. While metals like mercury, lead, and cadmium are toxic even at low concentrations [6,7,9].

The main contributor of heavy metals to the surrounding environment especially the marine ecosystem is anthropogenic. This includes an increase in agricultural practices, urbanization, industrial development, industrial effluents, communal sewage, smelting, fuel combustion, mining, etc. inadequate consideration for environmental impact, among other factors [9-11].

The presence of heavy metals in the marine ecosystem poses a serious health problem to both the marine organisms and the consumers of these marine organisms in the long run [6,12]. Heavy metals have prolonged half-lives; hence they do not degrade easily and they often bioaccumulate in living tissues through metabolic and bioabsorption processes [10]. This may result in marine organisms exhibiting symptoms of toxicity. The toxicity effects of these metals rarely manifest immediately after the exposure of toxins to the environment or the organisms, the symptoms become apparent mostly after a few years of exposure [9].

The exposure time of an organism in the marine ecosystem to a toxin has a significant effect on the level of bioaccumulation in such an organism. Irrespective of how short the exposure period might be, a significant amount of the metal may have been deposited in such an organism [9,10]. Other factors accounting for the accumulation of heavy metals in the tissue of aquatic organisms are salinity, pH, life cycle, size and age, temperature, hardness, ecological needs, capture season, and feeding habits of such organisms [10].

Marine organisms such as fish, periwinkle, crab, and oyster have been implicated as accumulators of trace metals and other heavy metals [10,13,14]. Hence, the assessment of these aquatic organisms for the presence of heavy metals is used to identify the associated environmental risks and potential toxic effects. The coastal sediments are host sites for pollutant metals, coupled with the devastating effect of certain toxic metals on human health, metal pollution is a significant cause of aquatic pollution. Benthic organisms (e.g., macroinvertebrates such as crustaceans, mollusks, etc.) are likely to be most directly affected by sediment metal concentrations because the benthos is the major repository of the particulate materials that wash into the aquatic systems [6,14,15]. Therefore, understanding the distribution and bioaccumulation of heavy metals in macroinvertebrates in coastal regions is crucial for scientists and government regulatory agencies [6]. Aquatic invertebrates are common bioindicators of choice. They have reduced mobility and are easy to see without the aid of a microscope. These organisms recolonize easily with a life span of over a year, are easy to sample, and are an important indicator of integrating ecological conditions [16]. Hence, macroinvertebrates are the specimens of choice in this study.

Compared to other coastal areas, the most affordable animal protein available to the locals around the coastline of the Ojo river is sea foods like fish, prawns, crayfish, shrimps, crabs, oyster, periwinkle, etc. obtained from the river, which therefore makes the maintenance of the health of the river and its resources inevitable. Most crustaceans are edible, and they serve as a healthy fish substitute in the diets of some of the locals. They also serve as a means of livelihood for those involved in its harvesting and marketing. However, there is limited information

regarding the metal contents and or the bioaccumulation potential of edible macroinvertebrates from the Ojo division of Lagos, Nigeria, particularly the Ojo river, hence the need for this study.

This study aims to provide information regarding the presence of heavy metals in the tissues of crab (*Potamon fluviatile*), prawn (*Macrobrachium rosenbergii*), and crayfish (*Metanephrops australiensis*) obtained from the Ojo river of Lagos State, Nigeria.

## 2. MATERIALS AND METHODS

### 2.1 Sample Collection

Live samples of macroinvertebrates such as *Metanephrops australiensis*, *Potamon fluviatile*, and *Macrobrachium rosenbergii* were purchased from the fishermen at the riverside in Ojo. The Ojo river is brackish, and it is located in the Ojo Local Government Area of Lagos State, Nigeria which is between latitude 40, 58' to 32.578"N and longitude 80, 20' to 30.124"E, and a surface area of about 264km. The crab, prawn, and crayfish bought were kept in plastic containers and transported immediately to the laboratory for analysis.

### 2.2 Sample Treatment

In the laboratory, a clean acid-washed porcelain crucible was kept in a drying oven at a temperature of 105°C for 24 hours to dry the samples. The dried samples were later ground into a fine powder form. The samples were kept till further analysis [17].

### 2.3 Sample Digestion

Five grams of ground powder samples were weighed into a clean crucible labeled according to the sample number and the dry-ashing process was carried out in a muffle furnace by a stepwise increase of the temperature up to 550°C and then left to ash at the same temperature for 6 hours. The samples were removed from the furnace and allowed to cool. The ash was made wet with water and 2.5 ml of concentrated HNO<sub>3</sub> was added. The crucible was covered with a watch glass and placed on a hot plate. The digestion was performed at a temperature of 90 to 95°C for 1 hour. The ash was dissolved in 5 ml of 9.25% HCl and digested again on a hot plate until the white fumes ceased and the sample volume reduced to 2 ml. After

cooling, 20 ml of distilled water was added and filtered using Whatman filter No.41. The filtered sample was then diluted up to the mark of 50 ml standard volumetric flask, and stored in a polyethylene container until analysis. All samples were prepared identically in triplicates. Blanks were prepared to check for background contamination by the reagents used [17,18].

### 2.4 Preparation of Standards and Analysis of Samples

Working standard solutions of Copper (Cu), Iron (Fe), Zinc (Zn), Chromium (Cr), Lead (Pb), Cadmium (Cd), and Manganese (Mn) were prepared from 1000 ppm element stock standard solutions in 2N HNO<sub>3</sub>. A calibration blank with three series of calibration standard solutions was used to calibrate the instrument, and an atomic absorption spectrophotometer (AAS) (Model: ELICO SL-194) was used to measure the elements present in the digested samples [17].

### 3. RESULTS AND DISCUSSION

Sea foods are essential sources of essential amino acids and protein for humans. Other nutrients inherent in seafood are vitamins, essential minerals with relatively low calories, and fat contents. Seafood like fish, crustaceans, and mollusks are the primary source of animal protein for the inhabitants of coastal areas [19]. However, because these marine organisms can absorb chemical contaminants from the

environment into their living tissues, the nutritional value and consumer health risk from seafood must be taken into consideration. Tables 1 and 2 below show the average concentration of heavy metals in crayfish, prawn, and crab obtained from the Ojo river and their comparison with the FAO/WHO standards. Crayfish contained more heavy metals than prawns and crabs (Fig. 1), lead was only detected in crayfish (Fig. 2).

From the result shown in Fig. 1, heavy metals in the macroinvertebrates vary in concentrations. Crayfish decreased in the following order: Fe > Zn > Cu > Cr > Mn > Cd > Pb, prawn decreased in the following order Zn > Fe > Mn > Cd > Cu > Cr > Pb, while crab decreased in this order; Zn > Fe > Cd > Cu > Cr > Mn > Pb. The phenomenon that different metals are accumulated at varying concentrations in different species in the environment was observed in this study. This phenomenon was also reported by Ekpo *et al.*, (2014) who assessed heavy metal concentrations in water, sediments, crab and periwinkle from the Uta Ewa Creek of Ikot Abasi, Akwa Ibom State, Nigeria [20]. The difference in the levels of accumulation of heavy metals in the different species examined in this study could be primarily attributed to the differences in the physiological role of each species. The regulatory ability of individual specie, behaviour and feeding habits are other factors that could have influenced the accumulation of metals in the different species [20].




Fig. 1. Heavy metal concentration in crayfish, crab, and prawn from Ojo river

**Table 1. Mean concentration of heavy metals (Mean±SD, mg/kg dry weight) in macroinvertebrates**


Samples	Metals (mg/kg)						
	Pb	Cu	Zn	Cr	Cd	Fe	Mn
Crayfish ( <i>Metanephrops australinsis</i> )	0.142±0.010	8.190±0.100	11.310±0.160	1.820±0.006	0.510±0.006	11.800±0.100	1.220±0.003
Crab ( <i>Potamon fluvidale</i> )	ND	0.035±0.003	3.728±0.170	0.033±0.002	0.085±0.005	1.099±0.144	0.013±0.007
Prawn ( <i>Macrobrachium rosenbergii</i> )	ND	0.022±0.002	1.850±0.150	0.015±0.002	0.032±0.003	1.620±0.099	0.660±0.003

\*ND-Not detected

**Table 2. Mean concentration of heavy metals (Mean ± SD, mgkg<sup>-1</sup> dryweight) in macroinvertebrate samples compared with FAO/WHO permissible limit**

Metals	Macroinvertebrate samples			FAO/WHO Permissible limit
	Crayfish ( <i>Metanephrops australiensis</i> )	Crab ( <i>Potamon fluviatile</i> )	Prawn ( <i>Macrobrachium rosenbergii</i> )	
Cu	8.190±0.100	0.035±0.003	0.022±0.002	73.3 [21]
Fe	11.800±0.100	1.099±0.144	1.620±0.099	425.5 [21]
Zn	11.310±0.160	3.728±0.170	1.850±0.150	99.4 [21]
Cr	1.820±0.006	0.033±0.002	0.015±0.002	2.3 [21]
Pb	0.142±0.010	ND	ND	0.3 [21]
Cd	0.510±0.006	0.085±0.005	0.032±0.003	0.2 [21]
Mn	1.220±0.003	0.013±0.007	0.660±0.003	0.4 [21]

\*ND - Not detected



**Fig. 2. The mean concentration of heavy metal within crayfish, crab, and prawn**

The mean concentrations of heavy metals assessed in this study were mostly below the permissible limit set by FAO/WHO except for the case of Cd in crayfish (Table 2). Cadmium concentration beyond the FAO/WHO permissible limit for food may inhibit metabolic enzymes when consumed over a long period [17]. Also, the mean concentration of Pb present in crayfish was below the permissible limit but was absent in both the crab and prawn assessed in this study (Table 2). This was similar to the report by Olowu et al., (2010), who reported the same manner of accumulation in crab and prawn samples where Pb was found to be absent in both the crab and the prawn samples. The absence of Lead may be attributed to the lack of heavy industrial activities around the sample site [17].

The metal body loads of the macroinvertebrates decreased in the following order: crayfish > crabs > prawns (Fig. 2), this may be due to the functional differences in the body of the macroinvertebrates, and this variation could also be an indication of the degree to which individual species absorb particulate matter from the surrounding water and sediments while feeding. Crayfish and crabs are bottom feeders and are generally expected to absorb more metals than surface feeders like prawns. This is in agreement with an earlier report made by Olowu et al. (2010), who assessed the level of heavy metals in crabs and prawns in the Ojo river, Lagos, Nigeria using atomic absorption spectrophotometer [17].

Several factors could predicate metal accumulation in marine organisms. These factors could be a result of living habits, feeding strategies, metabolic activities, and living environments in which these organisms are found [19]. The absence of lead in the crab and prawn could be related to activities around the location of the river [17]. The Ojo local government area consists mostly of farmlands, residential houses, and warehouses for furniture supplies. There are few or no industrial activities around the study location.

#### 4. CONCLUSION

In summary, from the evaluation of the results obtained in this study, the macroinvertebrates examined from the Ojo river possess heavy metals at a range within the safe limits set by the FAO/WHO for consumption except for Cd in crayfish. Both crab and prawn could be considered safe for consumption, however, the presence of Pb and the concentration of Cd in crayfish is an indication that it is unfit for consumption as Pb and Cd belong to the group of heavy metals that are toxic even in lower concentrations. There is a need for continuous monitoring of these macroinvertebrates especially crayfish to prevent potential health risks due to bioaccumulation of toxic metals over an extended consumption period. It is recommended that more sensitization be done to the public on the dangers of aquatic pollution and further assessments be done at regular intervals

to keep monitoring the metal body loads of the aquatic biota.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Field JG, Harris RP, Hofmann EE, Perry RI, and Werner F. Marine Ecosystems and Global Change; 2010.
- Bashir I, Lone F, Bhat RA, Mir SA, Dar ZA, and Dar SA. Concerns and threats of contamination on aquatic ecosystems, in Bioremediation and biotechnology. Springer. 2020;1-26.
- Irfan S, Alatawi AMM. Aquatic ecosystem and biodiversity: a review. Open Journal of Ecology, 2019;9(01):1.
- Bosch AC., O'Neill B, Sigge GO, Kerwath SE, and Hoffman LC. Heavy metals in marine fish meat and consumer health: A review. Journal of the Science of Food and Agriculture. 2016;96(1):32-48.
- Raknuzzaman M, Ahmed MK, Islam MS, Habibullah-Al-Mamun M, Tokumura M, Sekine M, et al. Trace metal contamination in commercial fish and crustaceans collected from coastal area of Bangladesh and health risk assessment. Environmental Science and Pollution Research. 2016; 23(17):17298-17310.
- Hao Z, Chen L, Wang C, Zou X, Zheng F, Feng W. et al. Heavy metal distribution and bioaccumulation ability in marine organisms from coastal regions of Hainan and Zhoushan, China. Chemosphere. 2019;226:340-350.
- Jenyo-Oni A. and Oladele A. Heavy metals assessment in water, sediments and selected aquatic organisms in Lake Asejire, Nigeria. European Scientific Journal. 2016;12(24).
- Mani T, Hauk A, Walter U, Burkhardt-Holm P. Microplastics profile along the Rhine River. Scientific Reports. 2015;5(1):1-7.
- Jakimska, A, Konieczka P, Skóra K, Namieśnik J. Bioaccumulation of Metals in Tissues of Marine Animals, Part I: The Role and Impact of Heavy Metals on Organisms. Polish Journal of Environmental Studies. 2011;20(5).
- Siraj M, Shaheen M, Sthanadar AA, Khan A, Chivers D, Yousafzai A. A comparative study of bioaccumulation of heavy metals in two fresh water species, *Aorichthys seenghala* and *Ompok bimaculatus* at River Kabul, Khyber Pakhtunkhwa, Pakistan. Journal of Biodiversity and Environmental Sciences. 2014;4(3):40-54.
- Ghorade I, Jadhavar V, Patil S. Assessment of heavy metal content in Amba river water (Maharashtra). World Journal of Pharmacy and Pharmaceutical Sciences (WJPPS). 2015;4(5):1853-1860.
- Arojoye OA, Oyagbemi AA, and Afolabi JM. Toxicological assessment of heavy metal bioaccumulation and oxidative stress biomarkers in *Clarias gariepinus* from Igbokoda River of South Western Nigeria. Bulletin of Environmental Contamination and Toxicology. 2018;100(6):765-771.
- Jakimska A, Konieczka P, Skóra K, and Namieśnik J. Bioaccumulation of metals in tissues of marine animals, Part II: metal concentrations in animal tissues. Polish Journal of Environmental Studies. 2011; 20(5).
- Wokoma O. Bioaccumulation of trace metals in water, sediment and crab (*Callinectes*) from Sombreiro River, Niger Delta, Nigeria. Internal J. Sci. Technol. Res. 2014;3(12):295-299.
- Rajeshkumar S, Liu Y, Zhang X, Ravikumar B, Bai G, and Li X. Studies on seasonal pollution of heavy metals in water, sediment, fish and oyster from the Meiliang Bay of Taihu Lake in China. Chemosphere. 2018;191:626-638.
- Greenfield R. Chapter 20 - Introduction to wetland monitoring, in Fundamentals of Tropical Freshwater Wetlands, T. Dalu and R.J. Wasserman, Editors. Elsevier. 2022;641-660.
- Olowu R, Ayejuyo O, Adejoro A, Adewuyi G, Osundiya M, Onwordi C, et al. Determination of heavy metals in crab and prawn in Ojo Rivers Lagos, Nigeria. E-Journal of Chemistry. 2010;7(2):526-530.
- Abalaka SE. Heavy metals bioaccumulation and histopathological changes in *Auchenoglanis occidentalis* fish from Tiga dam, Nigeria. Journal of Environmental Health Science and Engineering. 2015;13(1):1-8.
- Liu Q, Liao Y, Shou L. Concentration and potential health risk of heavy metals in seafoods collected from Sanmen Bay and its adjacent areas, China. Marine Pollution Bulletin. 2018;131:356-364.
- Ekpo F, Ukpong E. Assessment of Heavy Metals Concentrations in water,

- Sediments and some common sea foods *Callinectes amnicola* and *Tympanotonus fuscatus* (crabs and periwinkles) from Uta Ewa Creek of Ikot Abasi, Akwa Ibom State, Nigeria. *Global Journal of Applied Environmental Sciences*. 2014;4(1):99-106.
21. Mensah E, Kyei-Baffour N, Ofori E, Obeng G. Influence of human activities and land use on heavy metal concentrations in irrigated vegetables in Ghana and their health implications, in *Appropriate Technologies for Environmental Protection in the Developing World*. Springer. 2009;9-14.

© 2022 Reis et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/88399>