



Level of Mediations on Pain and Satisfaction Level of Patients Undergoing Abdominal Surgery

Ayesha Javaid ^{a*}, Mah Gull ^a and Iqra Tasleem ^b

^a Fatima Jinnah Medical University, Lahore, Pakistan.

^b Khawaja Muhammad Safdar Medical College, Sialkot, Pakistan.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JPRI/2022/v34i28A36015

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/84339>

Original Research Article

Received 17 January 2022

Accepted 24 March 2022

Published 28 March 2022

ABSTRACT

The main objective of the study was to analyse the level of mediations on pain and satisfaction level of patients undergoing abdominal surgery. This cross sectional study was conducted at Fatima Jinnah Medical University Lahore. The study was performed with 124 patients undergoing abdominal surgery under general anesthesia aged 18-65, with no diagnosis of cancer, no chronic pain, hospitalized for at least 24 hour following the surgical procedure, undergoing elective surgery and agreeing to participate. The data was collected from 124 patients with mean age 18 to 40 years. The data was collected from both genders. Of a total of 124 abdominal-surgery patient subjects, 58.1% were male, 94.3% were married, 61.3% had undergone surgery, and 41.1% had undergone total gastrectomy. Among the 124 participants, 47 were included in the control group, 36 in experimental group. Regarding their general characteristics, no significant differences in any aspects were observed. It is concluded that preoperative nursing intervention for pain has positive effects for patients undergoing abdominal surgery.

Keywords: Abdominal surgery; gastrectomy; mediations; anesthesia.

1. INTRODUCTION

Post-operative care is the process of providing care to patient after surgery. Post-operative care is necessary to reduce the occurrence of complications like shock, hemorrhage, pneumonia, respiratory depression, deep vein thrombosis, urinary tract infection, pressure sores, incisional hernia, bowel obstruction, infection and to enhance recovery of the surgical wound, and bring back patient to normal level of health and functioning [1-2]. Post-operative care must be provided to patient for immediate recovery and prognosis of the patient after surgery. Post operative nursing care include ,providing interventions such as continuous observation and monitoring of vital signs, maintenance of personal hygiene , early ambulation ,steam inhalation, care of surgical wound and incentive spirometry³. Post-operative care is an essential aspect in post-operative period as it provides opportunities for nurses to care for patients in a comprehensive and holistic manner so as to achieve early recovery of patient health status [4-5].

Anxiety is an individual experience and it is a concept that is difficult to describe with words. No matter how major or minor an operation is, it tends to raise a certain level of anxiety in every patient. Hospitalization for surgical procedure can be experienced as a threat or stressor and may produce anxiety in patients [6]. The main objective of the study was to analyse the level of mediations on pain and satisfaction level of patients undergoing abdominal surgery.

2. MATERIALS AND METHODS

This cross sectional study was conducted at Fatima Jinnah Medical University Lahore during 2020 to 2021. The study population consisted of patients undergoing abdominal surgery (benign prostate hypertrophy, appendicitis, acute cholecystitis, inguinal hernia, acute abdomen, uterine myoma, ovarian cyst, uterine bleeding, polyp etc.) in the surgical department. The study

was performed with 124 patients undergoing abdominal surgery under general anaesthesia aged 18-65, with no diagnosis of cancer, no chronic pain, hospitalized for at least 24 hour following the surgical procedure, undergoing elective surgery and agreeing to participate.

The data obtained were transferred to computer for analysis using SPSS version 15.00 (SPSS, Inc., Chicago, IL, USA) software. A p value <0.05 was considered significant. Numbers, percentages, and mean plus standard deviation were used at analysis.

3. RESULTS

The data was collected from 124 patients with mean age 18 to 40 years. The data was collected from both genders. Of a total of 124 abdominal-surgery patient subjects, 58.1% were male, 94.3% were married, 61.3% had undergone surgery, and 41.1% had undergone total gastrectomy. Among the 124 participants, 47 were included in the control group, 36 in experimental group. Regarding their general characteristics, no significant differences in any aspects were observed.

4. DISCUSSION

Despite progress in reducing pain, a universal experience, individuals still report experiencing it at various levels. For most patients, admission to hospital for surgery can be very stressful. Studies in this area support that requirements of patients to be informed in the preoperative period are not met, and anxiety can arise from lack of information [7]. In this study, all the patients who did not have adequate information about their disease and operation (51.7% in the study group -before they were instructed- and 45% in the control group) stated that they wished to get information from the healthcare personnel [8]. Emotional and psychological surgical preparation plays an important role in many areas of nursing [9].

Table 1. Differences in patients Knowledge regarding Pain Management

Variable	Pre	Post		F	p
		2 weeks	4 week		
	M ± SD	M ± SD	M ± SD		
Pain management	31.96 ± 3.62	32.50 ± 3.50	37.93 ± 2.36***	14.46	<.001

***p < .001

The powerful social factors affecting the reactions of women after hysterectomy are indicated as the educational status, income level, cultural structure, age at hysterectomy, short decision period before the operation, little support from the spouse and existence of a mental disorder preoperatively. In our study, no relationship was found between age groups and the level of anxiety ($p > .05$) [10].

5. CONCLUSION

It is concluded that preoperative nursing intervention for pain has positive effects for patients undergoing abdominal surgery. The intervention used in this study could serve as a guide for patients to improve the pain care of these patients.

DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

CONSENT

As per international standard or university standard, Participants' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Yilmaz M. Effect of the planned pre-operative teaching on the occurrence of post-operative complications and patient satisfaction. HEMAR-GE. 2002;4:40–51.
2. Guler H, Taksin L. The effect of planned education on coping with problems in post-hysterectomy period. Journal of Cumhuriyet University School of Nursing. 2001;5:9–18.
3. Karayurt O. The effects of different preoperative teaching methods on state anxiety preoperative and postoperative period and level of the pain postoperatively. Journal of Cumhuriyet University School of Nursing. 1998;2: 20–26.
4. Ozdemir F, Pasinlioglu T. The effects of training and progressive relaxation exercises on anxiety level after hysterectomy. The New Journal of Medicine. 2009;25:102–107.
5. Oner N, Le Compte A. State-trait anxiety inventory booklet. 2. İstanbul: Bogazici University Pub. 1985:21.
6. Farooqi YN. Depression and anxiety in patients undergoing hysterectomy. Journal of Pakistan Psych Society. 2005;2:13–6.
7. Karazeybek EA, Ozbayir T. Determination of the efficiency of the education of the patients who have had hysterectomy operation. Journal of Ege University School of Nursing. 2005;21:1–11.
8. Beatrice SK, Toivo H, Marten K, Kaarlo L, Pirkko N. Depression, anxiety, hostility and hysterectomy. Journal of Psychosomatic Obstetrics & Gynecology. 2005;26: 193–204.
9. Wang XQ, Lambert CE, Lambert VA. Anxiety, depression and coping strategies in post-hysterectomy Chinese women prior to discharge. International Nursing Review. 2007;54:271–279.
10. Helmy YA, Hassanin IM, Elraheem TA, Bedaiwy AA, Peterson RS, Bedaiwy MA. Psychiatric morbidity following hysterectomy in Egypt. Int J Gynaecol Obstet. 2008;102:60–4.

© 2022 Hussain et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/84339>