



Urbanorum spp. in Municipalities in the Metropolitan Region of Vale do Paraíba - SP, Brazil: Determination of the Epidemiological Profile

Hilary Rodrigues Bonito^{a†} and Matheus Diniz Gonçalves Coelho^{a*#}

^a Centro Universitário FUNVIC, São Paulo, Brazil.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/85980>

Received 09 February 2022

Accepted 18 April 2022

Published 21 April 2022

Original Research Article

ABSTRACT

Aim: The present study aims to define the epidemiological profile of *Urbanorum* spp. in cities of the Vale do Paraíba metropolitan region.

Materials and Methods: 2777 coproparasitological results were obtained using the technique based on the Hoffman, Pons & Janner protocol, in which 252 samples presented *Urbanorum* spp.

Results: The results showed that the city with the highest incidence was Guaratinguetá (15.3%) followed by Pindamonhangaba (10.2%) and Taubaté (9.9%). Regarding the epidemiological profile, there was a predominance of positive samples of *Urbanorum* in female sex, with 61.9% of the total positives and in the age group between 41 and 59 years, accounting for 34.5% of the samples.

Conclusion: It is concluded that there is a high incidence of *Urbanorum* spp. in the Vale do Paraíba region, being more relevant in adult women, aged between 41 and 50 years, with uniform distribution among the municipalities evaluated, but with greater occurrence in the municipality of Guaratinguetá.

Keywords: Coproparasitology; epidemiological profile; protozoa; protozoology; *Urbanorum* spp.

[†]Pharmacist Graduated;

[#]Professor and Researcher;

^{*}Corresponding author: E-mail: profmatheuscoelho@gmail.com;

1. INTRODUCTION

Intestinal parasitosis are diseases of high persistence, as they are based on a nutritional dependence of the parasite on the host, according to which the parasite cannot radically damage its host, since its obtaining of food is exclusively from the host and the damage or even the death of the same, would not be advantageous. Despite this need, the pattern for most parasitic species leads to manifestations such as anemia, irritability, sleep disturbances, nausea, vomiting, and diarrhea. In children, it can cause more serious cases, such as the development or worsening of malnutrition, cognitive and physiological delay, hepatosplenomegaly, anorexia, rectal prolapse, bowel and intestinal obstruction, and even death, regardless of age [1-3].

Currently, it is estimated that 3.5 billion people worldwide are affected by parasites. Family income, food, housing, basic sanitation, access to drinking water and health education are factors closely related to these diseases [2,4]. In this sense, population growth linked to socio-environmental issues can not only maximize the appearance of parasitic diseases, but also allows the creation of ecosystems favorable to the appearance of new species of parasites. Therefore, in 1991, in the city of Bucaramanga in Colombia, it was observed in a patient a correlation between clinical signs and symptoms with the presence of a possible species of the protozoan, then called *Urbanorum* spp [5].

Urbanorum spp, which is classified as belonging to the Phylum Sarcostomatophora, has an exoskeleton covered by a double membrane, consisting of one or two exit pores, through which its pseudopods are released, which help in its movement. Its size can vary from 80 to 100 microns, being considered large compared to other protozoa such as *Entamoeba coli*, *Endolimax nana* and *Giardia lamblia*. It has a round hyaline structure that is stained with Lugol. Patients affected by the protozoan may present symptoms such as secretory, liquid diarrhea with an acidic pH, without mucus, blood and leukocytes, as well as colic in the lower right hypochondrium, suggesting an infectious process derived from the colon [5,6].

The treatment of patients affected by this protozosis is through the use of commonly indicated antimicrobial drugs such as metronidazole and secnidazole. Its probable reproduction is by endodiogenesis and the

hypothetical evolutionary cycle of the parasite is started by the ingestion of water or food contaminated with fecal matter that contains mature cysts, which, releases the trophozoites in the colon, causing diarrhoea. Some trophozoites become cysts again, and both are passed in the feces. In between, the trophozoites are destroyed and only the cysts, resistant to the environment and can continue the cycle contaminating water and / or food [5-7].

One of the complications related to the parasitism caused by the protozoan studied is the Acute Diarrheal Syndrome that causes an increase in the number of stools, with watery stools, which may be accompanied by vomiting, fever and abdominal pain. Such disease can be considered as a risk factor capable of triggering malnutrition and dehydration, especially in children and population with nutritional gap, so prevention is of the utmost importance [8].

The epidemiological triad of *Urbanorum* spp is described by Díaz and Perlaza [9] and Bermeo and Sánchez [10], which encompasses three factors, the first being the agent: the protozoan directly attacks the large intestine. The second factor is the environment: feces find the soil, spreading its transmission through contaminated and poorly sanitized food, untreated water and direct contact with contaminated feces. And the third factor is the host, which, depending on its health status, can develop acute diarrheal syndrome, commonly observed in patients infected by this protozoan.

Studies report that in the city of Barrancabermeja in Colombia, the prevalence of *Urbanorum* spp was between 10 and 16% of the samples analyzed. But other countries have also reported the appearance of the protozoan [5,6]. In Peru, a study by Villafuerte et al. [6] shows images of the parasite, found in a 67-year-old woman, a resident of San Juan de Lurigancho, in Lima, who presented abdominal pain and dyspepsia for 9 months [6].

The first case report in Brazil occurred in 2017, in the city of Buriti in Maranhão. A 41-year-old patient weighed between 55 and 60 kg, lived in a rural area, without basic sanitation or treated water. She referred constant abdominal cramps and fever, being diagnosed with acute diarrhea due to symptoms. Laboratory tests were requested, including the coproparasitological examination, performed according to the Hoffman, Pons and Janer (HPJ) protocol, identifying the parasite [11].

In 2018, the second case report in Brazil was published in the city of Livramento, Rio Grande do Sul, a 72-year-old patient with Sjögren's syndrome. The patient did not present diarrhoea, signs of melena, hematochezia or enterorrhagia. The presence of *Urbanorum* spp was confirmed by the parasitological examination of feces, performed by the HPJ protocol [12]. In the same year, two other patients were diagnosed with *Urbanorum* spp. in Brazil [13-14].

The most recent case report in Brazil occurred in São José dos Pinhais-Paraná, being a 56-year-old male patient who used metformin for the treatment of diabetes, sought medical care for abdominal colic mainly in Brazil. The coproparasitological examination allowed the identification of *Urbanorum* spp. [15].

Finally, a survey conducted in a clinical analysis laboratory where 5,786 samples were analyzed, these being from different regions of Brazil, identified 84 positive results for *Urbanorum* spp., of which 43% were from the Midwest region of the country and 57% from Southeast, more precisely, from Greater São Paulo [16].

Being an emerging disease, the lack of information on this probable parasite raises the need to design surveys that aim to better understand its epidemiological profile, so in this study we seek to evidence the occurrence of *Urbanorum* spp., in the results of coproparasitological examinations of a clinical analysis laboratory, which serves residents of different localities in the metropolitan region of Vale do Paraíba - SP, Brazil, and also to verify the profile of the temporal and spatial distribution of the parasite in question, as well as its profile

distribution per person, taking into account the sex and age of the patients.

2. MATERIALS AND METHODS

This is a descriptive study, in which retrospective information was used from the results of coproparasitological examinations, carried out in a clinical analysis laboratory located in the municipality of Taubaté, which analyzes samples of patients from several municipalities in the metropolitan region of Vale do Paraíba, SP - Brazil, and which, for the analysis of the samples, uses the spontaneous sedimentation method, based on the HPJ protocol, being a method commonly used in the laboratory. The main sample was composed by 2,777 results of coproparasitological examinations of patients from the municipalities of Taubaté, Pindamonhangaba, Caçapava, São José dos Campos, Jacareí, Guaratinguetá and Campos do Jordão. Images of the parasite were obtained by making sample slides that were preserved in formalin and stored refrigerated in the laboratory.

The study used results of examinations carried out from February to August 2020. The data was analyzed in epidemiological terms, correlating the results with data regarding age, sex and municipality of origin. Statistical analyzes were performed using bioestat 5.0 software as a support tool.

4. RESULTS AND DISCUSSION

The decision to make the positive diagnosis for *Urbanorum* spp. was based on the morphological identification of structures compatible with the species (Fig. 1) and the comparative visual analysis with the findings of other researchers (Fig. 2).




Fig. 1. Sampling of evolutionary forms compatible with *Urbanorum* spp., identified in this study




Fig. 2. Pattern of evolutionary forms diagnosed as *Urbanorum* spp.: A- in the present study; B- in the findings of Villafuerte *et al.* [6] and C- in the findings of Casarin *et al.* [20]




Fig. 3. Incidence of intestinal protozoa and helminths detected in relation to the total number of positive samples in Vale do Paraíba, SP – Brazil (2021)

*Significant difference ($p < 0.0001$ - chi-square) in relation to the other species diagnosed

It was observed that 13.97% of the samples were positive for some evolutionary form of helminths or protozoa, which represents a low positivity. These results corroborate those presented by several researchers [17,18]. Among which Barbosa *et al.* [17] who evaluated the occurrence of coproparasitological test results from clinical analysis laboratories, which serve patients in the Midwest region of Minas Gerais, in the cities of Pará de Minas and São Gonçalo do Pará, and observed low positivity in 2014 (24%), possibly

reflecting adequate socioeconomic and sanitary conditions and, consequently, less exposure to risk factors for the transmission of enteroparasitosis in this region.

With respect to the proportional distribution of the diagnosed species, it was observed that there was a significant predominance ($p < 0.0001$, chi-square) of positive samples for protozoa, with 378 (97.4%) positive reports, compared to only 2.5% positive samples for helminths.

Urbanorum spp. presented the highest incidence coefficient in the period in which the study was carried out, representing 9.0% (252 positive samples, of a total of 2,777 samples evaluated, and 64,94% of the positive samples, being significantly higher ($p < 0.0001$, chi-square) than the coefficients of the other species identified, as can be observed in Fig. 3.

The low positivity for helminths has been observed in several surveys and is possibly linked not only to health improvements, but also to the practice of antiparasitic treatment in the routine of doctors who attend the Unified Health

System - SUS, in Brazil [17-19]. According to Belo *et al.* [19] the free distribution of Albendazole and its indiscriminate use when purchased in pharmacies, which does not occur for specific drugs for protozoa, are factors that contribute to a selective decrease in the occurrence of helminths.

Specifically in relation to the protozoan *Urbanorum* spp. a higher incidence was observed in the month of June, which was significantly higher ($p < 0.0001$) than that observed in the other months, as shown in Fig. 4.




Fig. 4. Temporal distribution of the incidence of *Urbanorum* spp. in municipalities of the Paraíba Valley, SP - Brazil (2020)

* Significantly higher value ($p < 0.0001$ - chi-square) than that observed in February, March and August




Fig. 5. Spatial distribution of the incidence of *Urbanorum* spp. in municipalities of Vale de Paraíba, SP - Brazil (2021)

* significant difference ($p < 0.05$ - Chi-square) in relation to the municipalities of Campos do Jordão, Jacareí and São José dos Campos

Regarding the spatial distribution of *Urbanorum* spp., it was possible to detect a significantly higher incidence ($p < 0.05$ - chi-square) in the municipality of Guaratinguetá, compared to the values observed in the municipalities of Campos do Jordão, Jacareí and São José dos Campos, as shown in Fig. 5.

Regarding the distribution by sex, it was observed that of the total positive samples for *Urbanorum* spp., 96 were from male patients (38.1%) and 156 from female patients (61.9%), with a significant predominance of women $P (<0.0001$ - Chi-square), corroborating with Casarin *et al.* (2019), who detected a predominance in the female sex, in the proportion of 1: 11.5 in relation to the male sex [20].

The age range of affected patients was wide, from 10 months to 91 years. The age group that was most abundantly positive was that of adults between 41 and 59 years, totaling 87 positive samples (34.5%), of which 50 samples belonged to female patients (57.5%) and 37 women to males (42.5%). The other data related to the distribution of incidence by sex and age group are shown in Fig. 6.

The distribution by age groups observed in the present study also agrees with the results obtained by Casarin *et al.* [20], who conducted an epidemiological survey to determine the occurrence of this species, and with similarity of the present study, identified a predominance of

positive results in adult patients. Regarding the distribution by gender, it was observed that there was no significant difference in the different age groups, with the exception of the age group of 19 to 40 years, in which a predominance of *Urbanorum* spp. significantly higher ($p <0.0001$ -Chi-square) in female patients [20].

Regarding the concomitant occurrence with other species, only in four reports it was positive for more than one species together with *Urbanorum* spp., being 3 samples with *Endolimax nana* and 1 with *Entamoeba coli*, species of protozoa that despite not being pathogenic, may possibly help in a better understanding of the mechanisms involved in the transmission of the protozoan in question, since such protozoa are transmitted primarily by ingestion of water or food contaminated by feces [21].

Finally, it should be inferred that, despite the unpublished nature of the results presented here, which may be useful for a better understanding of the epidemiological aspects related to *Urbanorum* spp., it is of fundamental importance that advanced tools be used, in order to test the biological condition of this. According to Díaz and Perlaza [9], research related to the standardization of biological culture, analysis of cell structure and molecular and genetic analysis, are of fundamental importance to determine its biological veracity, as well as to inspire studies that better determine its pathology, epidemiological and therapeutic aspects.




Fig. 6. Distribution by sex and age of the incidence of *Urbanorum* spp. in reports of coproparasitological examinations of patients treated in a clinical analysis laboratory in Vale do Paraíba, SP - Brazil (2021)

* significant difference ($p <0.0001$ - Chi-square) in relation to the rest of the age groups, except for the range between 19 and 40 years

5. CONCLUSION

It is concluded that there is a high incidence of *Urbanorum* spp. in the Vale do Paraíba region, being more relevant in adult women, between 41 and 50 years old, with uniform distribution among the evaluated municipalities, but with greater occurrence in the municipality of Guaratinguetá.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. De Melo MCB, Klem VGQ, Mota JAC, Penna FJ. Intestinal parasites. Rev Med Minas Gerais (1 Suppl. 1): S3-S12, 2004.
2. Neves DP, de Melo AL, Linardi PM, Vitor RWA. Human Parasitology. 11th ed.: Atheneu; 2004. 494 p.
3. Mascarini LM. A historical approach to the trajectory of parasitology. Ciência & Saúde Coletiva 2003;8(3):809-814. DOI: 10.1590/s1413-81232003000300015
4. Orlandini MR, Matsumoto LS. Prevalence of intestinal parasites in schoolchildren. State University of Northern Paraná, Campos Luiz Meneghel (UENP/CLM), 2010. [Cited on May 9, 2020]. Available: <http://www.diaadiaeducacao.pr.gov.br/portals/pde/arquivos/1655-8.pdf>
5. Rivero de Rodríguez Z. Es *Urbanorum* spp. a parasite?. Kasmera [Internet]. 2016 Jun [Cited on May 9, 2020] ; 44(1): 5-6. Available: http://ve.scielo.org/scielo.php?script=sci_arttext&pid=S0075-52222016000100001&lng=es.
6. Villafuerte RIM, Collado LAZ, Velarde CN. *Urbanorum* spp. IN EL PERU. Rev Peru Med Exp Salud Publica. 33(3):593-595, 2016. DOI: 10.17843/rpmesp.2016.333.2351
7. Ayol GLP, Guaman TJI. El *Urbanorum* spp. intestinal protozoan parasite. [Bachelor Thesis] State University of Milagro, Ecuador 2018. [Cited on June 15, 2020] Available at: <http://repositorio.unemi.edu.ec/handle/123456789/4159>
8. Guadalupe Vargas MJ, Evia Flores DI. Primary health care in diarrheal syndrome by *Urbanorum* spp. [Bachelor Thesis] State University of Milagro, Ecuador 2017. [Cited on June 15, 2020] Available: <http://repositorio.unemi.edu.ec/handle/123456789/3640>
9. Díaz KLG, Perlaza KFA. Epidemiology of acute diarrheal syndrome caused by the protozoan *Urbanorum* spp. [Bachelor Thesis] State University of Milagro, Ecuador 2018. [Cited on June 17, 2020] Available: <http://repositorio.unemi.edu.ec/handle/123456789/3636>
10. Bermeo ASF, Sánchez, LFH. Epidemiological chain and source of infection in acute diarrheal syndrome by the protozoan *Urbanorum* spp. [Bachelor Thesis] State University of Milagro, Ecuador 2018. [Cited on June 17, 2020] Available: <http://repositorio.unemi.edu.ec/handle/123456789/3643>
11. De Aguiar RPS, Alves LL. *Urbanorum* spp.: First Report in Brazil. American Journal of Case Reports; 19:486-490, 2018. DOI 10.12659/AJCR.908653
12. Lopez JCB, Nunes LS. 2018. *Urbanorum* spp.: second case report in Brazil proceedings of the 10th international teaching, research and extension fair - SIEPE Universidade Federal do Pampa | Santana do Livramento. [Cited on June 20, 2020] Available: <http://seer.unipampa.edu.br/index.php/siepe/article/download/39078/23894>
13. Perini LD, Lida EK, João EM, Braz GL, Madrugada HZA, Silva FSP, et al. *Urbanorum* spp., a new parasite? Case report. XVII Brazilian Digestive System Week – SBAD. São Paulo: Revista GED – Gastroenterology Endoscopy Digestive. 2018;37:322.
14. Turano CT, Mischiatti MN, Moura TV, De Lazaro BPB, Da Silva FPM. *Urbanorum* spp.: Case report of a new parasite in Brazil. XVII Brazilian Digestive System Week – SBAD. São Paulo: Revista GED – Gastroenterology Endoscopy Digestive. 2018;37:340.
15. Kruger EMM. *Urbanorum* spp.: new parasite in Brazil. Rev Bras Med Fam Comunidade. 2020;15(42):2157. DOI: 10.5712/rbmfc15(42)2157
16. Leão FMD, Siniauskas A, Corbucci R, Kiffer CRV. *Urbanorum* spp. in Brazil: are we facing a new epidemic parasitosis?.

- The Brazilian Journal of Infectious Diseases. 2018;22:124-125.
DOI: 10.1016/j.bjid.2018.10.238
17. Barbosa MCF, Duarte EV, Silva GM, Vieira JM, Silva LTC, Batista LT, *et al.* Investigation of the incidence of parasites in Pará de Minas-MG and region. SynThesis Revista Digital FAPAM 2016;7(7):171-178.
 18. Silva AA, Silva, PVR. da Rocha, TJM. Intestinal parasites: Frequency and epidemiological aspects in users of a private laboratory. Diversitas Journal. 2018;3(2):245-256.
Available:<https://doi.org/10.17648/diversitas-journal-v3i2.628>
 19. Belo VS, Oliveira RB, Fernandes PC, Nascimento BWL, Fernandes FV, Castro CLF, *et al.* Factors associated with the occurrence of intestinal parasites in a population of children and adolescents. Rev Paul Pediatr. 30(2):195-201, 2012.
DOI: 10.1590/S0103-05822012000200007
 20. Casarin JN, Duarte SMS and Sampaio JS. 2019. First reports from *Urbanorum* spp. Meeting in SUS patient floes in a private Imperatriz laboratory during 2018. International Journal os Development Research. 9(10):20676-30678.
 21. Silva-Díaz H. "*Urbanorum* spp.": Controversy of its biological condition and acceptance as a new intestinal parasite. Rev Exp Med 30;3(1):03-04, 2017.
[Cited on December 1, 2020].
Available:<http://rem.hrlamb.gob.pe/index.php/REM/article/view/73>

© 2022 Bonito and Coelho; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/85980>