



---

# **Histological Manifestations of Effects of Eliozeu Dumpsite Leachate on the Reproductive Tract of Female Wistar Rats**

**Kinikanwo I. Green <sup>a</sup> and Weleh I. Ikechukwu <sup>b\*</sup>**

<sup>a</sup> *Department of Obstetrics and Gynaecology, Faculty of Clinical Sciences, College of Health Sciences, University of Port Harcourt, Rivers State, Nigeria.*

<sup>b</sup> *Department of Human Physiology, Faculty of Basic Medical Sciences, College of Health Sciences, University of Port Harcourt, Rivers State, Nigeria.*

### **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

### **Article Information**

#### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/84821>

**Original Research Article**

**Received 12 January 2022**

**Accepted 20 March 2022**

**Published 07 April 2022**

---

## **ABSTRACT**

Reproduction is an important biological trait to produce new individual organisms and is the fundamental for the life of an individual as well as the survival and development of the system. The response of reproductive organs to toxic substances varies from that of other target organs and may serve as an ideal for the deleterious effects of environmental toxicity on animals and human health. The incidence of infertility and risen maladies are increased in the last five decades. Various human activities have released into the environment plenty toxic substances which could be found in dumpsite leachate. The present study focused on experimental studies of histological effects of Eliozeu dumpsite leachate on reproductive tract of female using Wistar rats as model. Twenty five (25) female Wistar rats were divided into five groups of five animals each; the leachate was collected from the dumpsite and water from near-by borehole also collected. Group 1 which served as control group received 1ml of commercial bottle water, group 2 received 1ml of borehole water 1kilometer from the dumpsite, groups 3, 4 and 5 received different concentration of the leachate in 10%, 50% and 100% for thirty days, the animals were sacrificed after being anesthetized with chloroform vapor, the ovaries and uterus were collected and put in 10% formaldehyde, sectioned

---

\*Corresponding author: E-mail: [lyke\\_wele@yahoo.com](mailto:lyke_wele@yahoo.com);

and stained with haematoxylin and eosin (H&E) for histopathological studies. The slides were placed in a light microscope and analyzed by a consultant Anatomy pathologist. The results revealed that the Elioazu dumpsite leachate does not have any effect on the morphology of the uterus but have deleterious effect on the histomorphology of the ovaries. Elioazu dumpsite leachate has shown deleterious effect on female ovaries, which is an indication of toxicity that may cause infertility to the experimental animals. Therefore, it is recommended that further studies aimed at corroborating this finding be carried out especially on humans and residents around dumpsite should always embark on reproductive tract medical check-up.

*Keywords: Elioazu dumpsite leachate; histological effect; reproductive tract.*

## 1. INTRODUCTION

In developing countries, the dumpsite is the most cost-effective method of disposing of municipal solid garbage [1]. However, leachate, a liquid effluent, may leak from such a stockpile and contaminate the neighbouring water body [2]. Uncontrolled overflow, rainfall, and infiltration into the underlying ground water aquifer or surrounding surface water can all expose the environment to dumpsite leachate. Leachate is increasingly recognised as a possible health hazard to both ecosystems and human populations [3]. Heavy metal concentrations were found in high concentrations in leachates at the Delta State Teaching Hospital Oghara dumpsite, according to studies [4]. Characterization and toxicological assessment of leachates from the Air Hitman Sanitary Landfill in Malaysia using *Pangasius Sutchis* and *Clarias batrachus* found that the leachate enclosed significant amounts of ammonia, dissolved organic matter, some semi-volatile organic compounds, and monocyclic aromatic hydrocarbons, all of which are highly toxic to both species of fish, and that these compounds could be used as indicators of leachate contamination in freshwater. Ammonical-nitrogen in the leachate is thought to be the main cause of fish mortality. However, benzene, toluene, and ethyl benzene may have played a role in the toxicity of the leachate [5]. Copper, chromium, lead, arsenic, cadmium, manganese, nickel, ammonia, chloride, and phosphate concentrations in the leachates samples of the Elioazu dumpsite were found to be beyond the allowed levels, according to Weleh and coworkers [6]. Heavy metals that may be present in leachate have been shown to have haematotoxic potential in rats [7]. In rats, raw and simulated leachates from the Olusosun municipal solid waste landfill were found to have immunotoxic potential (Subhasini et al, 2009). Subhasini et al. found changes in the biochemical and histopathological profile of the liver in Swiss Mice treated with distillery soil

leachate at concentrations of 5%, 10%, and 20% [9]. Hepatotoxicity and oxidative stress were also detected in rats in a similar survey in Olusosun municipal landfill leachate in Ojota, Lagos State, Nigeria. Neurologic lesions, neurodegeneration of purkinje cells with loss of dendrites, perineural vacuolations of the neuronal cytoplasm (spongiosis) and neuronal necrosis in the brain of Wistar rats exposed to Olusosun and Aba-Eku municipal landfill leachate has also been studied [6]; these reported brain tissue alterations correlate with a significant decrease in body weight gain and Superoxide dismutase activity but increase in absolute and relative brain weight gain, Malondialdehyde concentration and Catalase activity in both brain and serum in treated rats [6].

The Elioazu Dumpsite is one of Rivers State's largest dumpsites, receiving untreated deposits of both domestic and industrial wastes from all over Port Harcourt, despite its potential direct and indirect hazards, given that residents in the dumpsite's vicinity rely heavily on borehole water for drinking. Secondly, despite the fact that infertility remains prevalent in our environment and premium placed on child bearing, there is paucity of literature on the possible effect of leachate on reproduction *visa-viz* reproductive tract. This study therefore attempts to evaluate the impacts of Elioazu dumpsite leachate on the histology of the reproductive tract using Wistar rats as experimental model. Experimental animals and data from the toxicological studies in Wistar rats is a way of forecasting human risk assessment.

## 2. MATERIALS AND METHODS

### 2.1 Experimental Animals

Twenty-five female Wistar rats (10-12 weeks old, weighing about 200g) were obtained from the Animal House unit of Faculty of Basic Medical Sciences, University of Port Harcourt, Nigeria.

The animals were cared for according to the general recommendation and provision of the University Ethical committee. Rats were housed in a climate controlled room (12 hours' dark and light cycle), with free access to drinking water and standard rodent chow (*ad libitum*).

## 2.2 Leachate Collection and Preparation

Raw leachate sample was collected from leachate well on the dumpsite at Eliozu in Obio/Akpor local government area of River State, Nigeria in clean 5litres plastic containers. The sample was taken to the laboratory of the Department of Chemistry, University of Port Harcourt where it was filtered and prepared accordingly as described by [9]. The different concentrations of the leachate were gotten by serial dilution of leachate with distilled water.

## 2.3 Experimental Design

Following two weeks of acclimatization, the rats were divided into five groups of five Wistar rats

each. Group 1 served as the control and received 1ml of commercial bottled water; Group 2 received 1ml of water obtained from borehole about 1km from the dumpsite; while Groups 3, 4 and 5 received 1ml of 10% of leachate concentration, 1ml of 50% of leachate concentration and 1ml of 100% of leachate concentration respectively. All administrations were given once daily using an orogastric cannula for 30 consecutive days. On the 31<sup>st</sup> day however, the animals were anaesthetized using chloroform vapor and the ovaries and uterus were harvested for histological assay.

## 2.4 Histopathological Examination

Uterus and ovary collected were fixed with 10% formalin, embedded in paraffin, sectioned and stained with hematoxylin-eosin (H&E). Experienced pathologist performed pathological evaluation.

## 2.5 Histology of the Uterus and Ovary




Plate 1 (H & E) x100




Plate 2 (H & E) x100




Plate 3 (H & E) x100




Plate 4 (H & E) x100




Plate 5 (H & E) x100




Plate 6 (H & E) x100

### 3. RESULTS

The result of photomicrograph of the Wistar rats uterus stained with H&E at the same magnification (X100) displayed in Plates 1 to 3 normal Perimetrium (PM), Myometrium (MM), Endometrium (EM) containing endometrial glands, Simple columnar epithelium lining the endometrium in plate 1 (control). Plate 2 (borehole water) shows the same morphology when compared with the control and plate 3 (100% leachate) also showed the same morphology, without any alteration of the uterus. However, the result of photomicrograph of the Wistar rats ovary stained with H&E at the same magnification (X100) were displayed in Plate 4 to 6. Plate 4 (control) shows normal morphology of the ovary with Medulla (MD), Cortex containing primary follicle (PF), and corpus Luteum (CL), plate 5 also shows the same morphology of the ovary while plate 6 shows distorted ovary with swelling of the Graafian follicles. This alteration in the histology of the ovary examination implies a remarkable alteration of the ovary but not in the uterus by leachate from Elioizu.

### 4. DISCUSSION

In developing countries paucity of suitable technological choices could be the reason solid waste are dumped [1]. Exposure to leachate causes health risk. The physicochemical properties of Elioizu dumpsite leachate is higher compared with the accepted values of National Environmental Standard Regulations Enforcement Agency as described by [6].

The present study showed that there were no significant morphological changes in the examined uterus; this could be as a result of the short duration of exposure to the dump site leachate. We observed abnormalities in the ovary of the experimental animals that received pure concentration of leachate. There were cellular hypertrophy, degenerative and atrophic changes and swelling of ovarian graffian follicle when compared with the control group. This is in support with earlier report by [9], on the impact of Elioizu Port Harcourt landfill leachate on some reproductive hormones in female Wistar rats, where it was observed that Elioizu leachate altered female reproductive hormones. This finding indicates that the leachate may have deleterious effects on the oocyte of the ovaries of adult Wistar rats, in extension may contribute to the causes of female infertilities in the human population. It is similar with an earlier report by

Eweka and Om'Imaboh's [10], on the histological studies of effect of Monosodium Glutamate on the ovaries of adult Wistar rats. Cellular swelling is known to occur either directly or indirectly by denaturation of volume-regulating ATPases or indirectly by disruption of the cellular energy transfer pathway required for ionic regulation [11]. This study further supports an earlier report by Ayandiran *et al.*, (2017). The abnormality observed in the ovary is also similar with an earlier report by [12], who reported that raw and simulated solid waste leachate from three dumpsite in Nigeria induce dose dependent increase in frequency of sperms with abnormal morphology in mice. There were similar reports in mice treated with leachates from electronic waste and hospital waste incinerated bottom ash [13]. The causative agents of this reproductive abnormality could be lead and mercury as seen in the physicochemical analysis of Elioizu dumpsite leachate [6].

Further studies on heavy metals revealed that arsenic, lead, cadmium cause health implications including disturbances in reproduction [14].

In a study, effect of long-term exposure of female Wistar rats to low levels of lead: ovary and uterus histological architecture changes, observed that lead values and the micromorphological structure changes found in the study in the ovaries, the uterus and fallopian tubes demonstrated the deleterious effect of lead, this study is similar with our work on ovary but differs in the morphology of the uterus [15].

Also record shows deleterious effect of the ovaries and uterus exposed to cadmium, lead and mercury on the structure and function of reproductive organs. It further explained that cadmium affects the maturation of follicles, degradation of the corpus luteum [16].

This observation suggests that leachate from Elioizu dumpsite is capable of inducing damage in germ cells in experimental animals indicating ovarian cyst and data from the toxicological studies in Wistar rats may be a way of forecasting human risk assessment.

### 5. CONCLUSION

The reproductive tracts of Wistar rats exposed to Elioizu dump site leachate has been examined using histological method. The examination showed no abnormality in the morphology of the uterus. However, the ovary of the experimental

rats' exhibited morphological abnormalities which is an indication that dumps site leachate may contribute to the causes of female infertility. Therefore, it is recommended that further studies aimed at corroborating this finding be carried out especially on human population and medical check-ups be carried out on regular basis on the reproductive tracts.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

The ethical approval for this study was sought and obtained from the institution research ethics committee on 3<sup>rd</sup> July, 2018. The approval reference number is UPH/R&D/REC/04.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Vaccari M, Torretta V, Collivignarelli C. Effect of improving environmental sustainability in developing countries by upgrading solid waste management techniques: A case study. *Sustainability*. 2012;4:2852-2861.
2. Chatham-Stephens K, Caravanos J, Ericson B, Landrigan P, Fuller R. The pediatric burden of disease from lead exposure at toxic waste sites in low and middle income countries. *Environmental Research*. 2014;132:379-383.
3. Al Sabbagh MK, Velis CA, Wilson DC, Cheeseman CR. Resource management performance in Bahrain: A systematic analysis of municipal waste management, secondary material flows and organizational aspects. *Waste Management Research*. 2012;30:813-824.
4. Nwaka PO, B Anegbe, O Adeniyi, IG. Okunzuwa, A Jidonwo. Impact of leachate on physicochemical properties of soil, within the vicinity of Oghara medical dumpsite, Delta State, Nigeria. *Physical Science International Journal*. 2018;17(1):1-14.
5. Tsai CL, Krogmann U, Strom PF. Effect of forced aeration and mechanical turning on leachate quantity and quality from glass cullet stockpiles. *Journal of Environmental Engineering*. 2010;136(8):854-9.
6. Weleh II, Weleh IE, Green IK. Impact of (ELIOZU) Dumpsite Leachates on Some Haematological Parameters in Wistar Rats. *International Journal of Research and Reports in Hematology*. 2021;4(2):70-78.
7. Afolayan, OS, Ogundele FO, Odewumi SG. Spatial variation in landfills leachate solution in urbanized Area of Lagos State, Nigeria. *American International journal of contemporary Research*. 2012;2(8):178-184.
8. Subhasini Sharma, Kalpana Sharma, Oivedita Yadav, KP. Sharma. Alterations in Biochemical and Histopathological Profile of Liver in Distillery Soil Leachate Treated Swiss Albino Mice (*Mus Musculus L.*). *Pharmacologyonline*. 2009;3:1047-1053
9. Green I. Kinikanwo, Oriji K. Vadunume, Weleh I. Ikechukwu. Impact of ElioZu Port Harcourt Landfill Leachate on Some Reproductive Hormones in Female Wistar Rats, *International Journal of Research and Reports in Gynaecology*. 2020;3(1):20-24.
10. Eweka AO, Om'Imaboh's FAE. Histological studies of the effects of monosodium glutamate on the ovaries of adult wistar rats. *Annals of Medical and Health Sciences Research*. 2011;1(1):37-43.
11. Hinton DE, Laure DJ. Liver Structural alterations accompanying chronic toxicity in fishes; potential biomarkers of environmental contamination. In: McCarthy JF, *Biomarkers of Environmental Contamination* Boca Raton: Lewis. 1990; 17-57.
12. Bakere AA, Alimba CG, Alabi OA. Genotoxicity and mutagenicity of solid leachates: A review. *African Journal of Biotechnology*. 2013;12(27): 4206-4220.
13. Akinbola Temitayo, Adetutu Adeyemi, Olajumoke A. Morenikeji and Adekunle Bakere. Hospital waste incinerator bottom ash leachate induced cytogenotoxicity in *Allium cepa* and reproductive toxicity in mice. *Toxicology and Industrial Health*. 2011;27(6):505-514.
14. Piotr Rzymyski, Katarzyna Tomczyk, Pawel Rzymyski, Barbara Poniedzialek, Tomasz Opala and Maciej Wilczak. Impact of heavy metals on the female reproductive system.

- Annals of Agriculture and Environmental medicine. 2015;22 (2): 259-264.
15. Eugenia Dumitrescus, Victoria Chiurciu, Florin Muselin, Roxana Popescu, Diana Brezovan, Romeo T. Cristina. Effects of long-term exposure of female rats to low levels lead: ovary and uterus histological architecture changes. Turkish Journal of Biology. 2015;284-288.
16. Peter Massányi, Martin Massányi, Roberto Madeddu, Robert Stawarz and Norbert Lukác. Effects of Cadmium, Lead and Mercury on the Structure and Function of Reproductive Organs. Toxic. 2020;8:94.

© 2022 Green and Ikechukwu; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/84821>