



Preliminary Phytochemical Profile, *in vitro* Antioxidant and Sun Protective Activities of *Alhagi pseudalhagi* and *Elaeagnus angustifolia* L.

**Mohammad Humayoon Amini^{1*}, Amina Ahmady², Aqa Mohammad Zhakfar²,
Mohammad Nasim Sediqi¹ and Gulalai Babak²**

¹Department of Pharmacognosy, Faculty of Pharmacy, Kabul University, Kabul, Afghanistan.

²Department of Pharmaceutics, Faculty of Pharmacy, Kabul University, Kabul, Afghanistan.

Authors' contributions

This work was carried out in collaboration among all authors. Authors MHA, AMZ and MNS collected the plants raw materials. Authors MHA and AA conceived, designed and performed the experiments, analyzed and interpreted the data and drafted the manuscript. All authors read, revised and approved the final version of the manuscript before submission.

Article Information

DOI: 10.9734/JPRI/2019/v31i430308

Editor(s):

(1) Dr. Q. Ping Dou, Professor, Barbara Ann Karmanos Cancer Institute, Departments of Oncology, Pharmacology and Pathology, School of Medicine, Wayne State University, USA.

Reviewers:

(1) Maria Bintang, IPB University, Indonesia.

(2) Moldovan Zenovia, University of Bucharest, Romania.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/51341>

Original Research Article

Received 08 July 2019
Accepted 16 September 2019
Published 12 November 2019

ABSTRACT

Background and Objectives: *Alhagi pseudalhagi* and *Elaeagnus angustifolia* are among the medicinal plants used traditionally in Afghanistan. The aim of present work was to study *A. pseudalhagi* herbs and *E. angustifolia* leaves for their preliminary phytochemical profile, total phenolic content (TPC), total flavonoid content (TFC), *in vitro* antioxidant activity and sun protective factor (SPF).

Methods: *Alhagi pseudalhagi* herb crude extract (APCE) and *E. angustifolia* leaves crude extract (EACE) were prepared using Soxhlet apparatus. Standard qualitative phytochemical tests were performed to detect various phytochemicals in the crude drugs. Both of APCE and EACE were purified as per previously described method to obtain *A. pseudalhagi* herb purified extract (APPE) and *E. angustifolia* leaves purified extract (EAPE). The TPC, TFC, and SPF of the extracts were determined, Following prescribed methods. To evaluate *in vitro* antioxidative potential of the

*Corresponding author: E-mail: m_hamayon@yahoo.com;

extracts, DPPH radical scavenging assay method was performed. The experiments were performed in three replications.

Results: Carbohydrates, phenolics, flavonoids, sterols, tannins, and saponins were present in all tested extracts. The TPC of tested extracts ranged from 71.14 ± 2.10 to 203.76 ± 2.07 mg gallic acid equivalent (GAE) / g dried extract, as recorded for APCE and EAPE, respectively. TFC values of 33.24 ± 3.23 and 135.85 ± 3.83 mg rutin equivalent (RUE) / g dried extract were found for EACE and EAPE, respectively. TFC values of 59.61 ± 3.40 and 121.06 ± 3.37 were recorded for APCE and APPE, respectively. DPPH radicals scavenging capacity and SPF of the tested extracts were positively correlated with their TPC and TFC. The EAPE and APPE produced IC_{50} value of 4.5 and 14.31, respectively. The higher SPF of 29.73 ± 0.31 was produced by EAPE, while APPE exhibited SPF of 17.83 ± 0.88 , as calculated for 2mg/ml methanol solution of the extracts.

Conclusion: Based on present study, EAPE revealed higher TPC and TFC values and hence, showed potent *in vitro* antioxidant potential and higher SPF value, as compared with *A. pseudalhari* herb extracts. Therefore, EAPE could be highly recommended to be used in development of herbal sunscreen formulations.

Keywords: *Alhagi pseudalhari*; *Elaeagnus angustifolia*; antioxidant; sun protective factor; total flavonoid content; total phenolic content.

1. INTRODUCTION

Medicinal plants (MPs) have been used by mankind as curing agents since immemorial time, and still they are known as natural remedies for divers array of maladies. According to WHO, about 80% of world population still rely on traditional herbal medicines in case of their primary health care needs [1,2]. Herbal extracts owing to their complex compositions exhibit different biological and pharmacological effects; such as anti-inflammatory, antioxidant, antiapoptotic immunomodulatory, sun blocking, anti-cancer activities, and so on. Different botanical extracts are reported for their free radical scavenging capacity, anti-photo aging effects, and their role in treatment of different skin diseases and improvement of skin appearance [3-6].

Excessive solar UV radiations (UVR) that cause formation of excess Reactive Oxygen Species (ROS), are known to be responsible for various skin damages such as; sunburn, skin pigmentation, premature aging, photo carcinogenesis, and skin cancers [7-9].

Applying topical sunscreens is widely encouraged in order to protect skin from the adverse consequences of solar UVR. An effective sunscreen besides containing active screening agents, should contain antioxidants to scavenge ROS and to efficiently prevent skin from photo aging and skin cancers. Plants' antioxidants e.g. phenolic compounds and flavonoids are suitable candidates to be incorporated in sunscreen formulations [10].

Currently, plants as well as marine organisms are considered to be valuable sources of potential anti-photoaging and photoprotective agents for development of cosmeceutical and topical pharmaceutical formulations [11].

Alhagi pseudalhari (Syn. *A. maurorum*) belongs to family Fabaceae, is a small thorny shrub with stiff and spiny stem and branches. The plant is called *Shuthur khar* in Afghanistan and is used in traditional medicine of the country for treatment of kidney disease, urinary tract infections, gonorrhoea, bleeding, wounds, hypotension, and gastrointestinal disorders [12]. *Alhagi pseudalhari* is reported to be rich in several bioactive compounds e.g. flavonoids, flavone glycosides, alhagidin, alhagitin, proanthocyanidins, triterpenes, tannins, and phenolic compounds [13,14].

Elaeagnus angustifolia L. belongs to family Elaeagnaceae, is a small to medium sized tree or shrub, 2 – 5 m tall and grow abundantly in different parts of Afghanistan [15]. *Elaeagnus angustifolia* is used in traditional medicines as antipyretic, diuretic, tonic, antidiarrheal, anti-inflammatory, analgesic, and as a medication against kidney stones [16]. Different parts of the plant including its leaves are rich in various phytochemicals such as; polysaccharides, sitosterols, glycosides, terpenoids, coumarins, saponins, carotenoids, vitamins, tannins, phenolics and flavonoid compounds that act as natural antioxidants [17-19].

Unfortunately, because of the long-lasting wars and other political issues, study of MPs grow in

Afghanistan is neglected, and hence, publications regarding phytochemical profile and biological activity of Afghani MPs are very scarce. Considering worldwide interests into research on MPs and the huge demand of natural sunscreens, the main objectives of present research were to evaluate the preliminary phytochemical profile, total phenolic content (TPC), total flavonoid content (TFC), *in vitro* antioxidant potential, and sun protective factor (SPF) values of *A. pseudalhagi* herb and *E. angustifolia* leaves collected from Afghanistan.

2. MATERIALS AND METHODS

2.1 Plant Materials

Elaeagnus angustifolia leaves were collected from Paghman district of Kabul on 7th of June, 2017, and *A. pseudalhagi* herb was collected at blooming time from Kabul University campus on 2nd July, 2017. Collected materials were botanically authenticated by Prof. Mohammad Nasim sediqi, Head of Pharmacognosy department, Faculty of Pharmacy, Kabul University. Voucher specimens of the collected *A. pseudalhagi* and *E. angustifolia* bearing No. 1955 and No.1956 respectively, were deposited in herbarium of faculty of Pharmacy, Kabul University, for further use as references. Collected materials were shade dried and after being powdered, passed through mesh size 1400 μ . Powdered materials were separately packed in polyethylene bags for further use.

2.2 Preparation of Plants Extracts

Conventional extraction method using soxhlet extractor was used in preparation of the crude extracts. 20 g of the dried and powdered *A. pseudalhagi* herb and *E. angustifolia* leaves were separately extracted with 70% methanol (MeOH), at 70°C, until the marc was exhausted. The extracts were filtered through Watmann No. 1, and then were concentrated under vacuum, at 40°C. Concentrated extracts were further dried in oven at 60°C, until getting their constant weights (complete drying). The dried *A. pseudalhagi* herb crude extract (APCE) and *E. angustifolia* leaves crude extract (EACE) were kept in refrigerator at 4°C for further uses.

2.3 Purification of Extracts

To purify the extracts from chlorophyll and other inert lipophilic compounds, the method described by Jarzycka et al. with slight modifications were followed [20]. Accurately weighed amounts of

both APCE and EACE were first dissolved in 70% MeOH, then extracted 4 times with equal amounts of its volume petroleum ether. The MeOH fractions were dried at 60°C in drying oven. The dried extracts were then separately dissolved in hot distilled water, followed by addition of ascorbic acid (0.5 mg/g dried extract) to the mixture. Each of the mixture was left in refrigerator for 24 hours, and then, was extracted five times with diethyl ether. The last aqueous fractions were further extracted five times with ethyl acetate. In both cases, the ethyl acetate fraction was evaporated to obtain the dried *A. pseudalhagi* herb purified extract (APPE) and *E. angustifolia* leaves purified extract (EAPE). APPE and EAPE were used for further studies. The yield percentages of the crude and purified extracts were calculated and shown in Table 1.

2.4 Preliminary Phytochemical Screening

Methanol solutions (2 mg/ml MeOH 70%) of the extracts, e.g. APCE, EACE, APPE, and EAPE were separately prepared and subjected to preliminary phytochemical screening to detect diverse categories of phytochemicals in the extracts. Standard procedures and methods previously described were followed [21-23].

2.4.1 Detection of carbohydrates

2.4.1.1 Molisch's test

About 3 ml of test solution (TS) was mixed with 3 drops of Molisch's reagent (alpha-naphthol 20% in ethyl alcohol) in a clean test-tube. Then about 2 ml of conc. H₂SO₄ was carefully added along the side of the test-tube. Formation of a reddish violet or dark purple colour ring at the junction of the two layers indicated presence of carbohydrates.

2.4.1.2 Resorcinol test

A few small crystals of resorcinol were dissolved in about 2 ml of TS in a test-tube. Then an equal volume of conc. HCl was added to the test-tube, followed by warming on water-bath for a few minutes. Formation of a rose colour indicates presences of ketoses.

2.4.2 Detection of alkaloids

2.4.2.1 Dragendorff's test

Three drops of Dragendorff's reagent (potassium iodide + bismuth nitrate) were added on 2 ml of

TS. Formation of an orange red/brown colour precipitate indicates presence of alkaloids.

2.4.2.2 Hager's test

Few drops of Hager's reagent (saturated picric acid solution) were added on 2 ml of TS. Formation of a yellow precipitate reveals a positive result.

2.4.2.3 Mayer's test

Few (2 - 3) drops of Mayer's reagent (solution of Potassium mercuric iodide) were added on 2 ml of TS. Formation of a white, pale yellow or cream precipitate indicates presence of alkaloids.

2.4.2.4 Wagner's test

About 2 – 4 drops of Wagner's reagent (Iodine solution) were added on 2 ml of TS in a test-tube. Formation of a brown or reddish-brown precipitate indicates a positive result.

2.4.3 Detection of glycosides

2.4.3.1 Legal's test

About 2 ml of TS were treated with 2 ml of sodium nitroprusside 2% solution in pyridine, and few drops of 20% sodium hydroxide in MeOH. The mixture was observed for formation of a pink to deep blood red colour.

2.4.3.2 Keller-Kiliani test

About 3 ml of TS were mixed with 2 ml glacial acetic acid and 1 ml ferric chloride (5%) solution. The mixture was heated on water-bath, then cooled and transferred to a test-tube containing 2 ml of concentrated H₂SO₄, and observed for formation of a reddish-brown colour at the junction of the liquids, and gradually the acetic acid layer turns blue.

2.4.4 Detection of anthraquinone glycosides

2.4.4.1 Borntrager test

Few mg of the test extract was treated with 5 ml HCl 0.5 N, then it was mixed with 5 ml chloroform for 15 minutes in a separatory funnel. Then, the mixture was placed aside for being separated into two phases. The chloroform phase was taken in a clean test-tube, followed by addition of 2 ml of 10% ammonia solution. The mixture was shaken well; then, it was stand till the two phases

were separated. Appearance of a pink or reddish colour in the aqueous phase indicated a positive result.

2.4.5 Detection of phenols

2.4.5.1 Ferric chloride test

About 2 ml of TS was placed in a clean test-tube, followed by addition of 2 – 3 drops of 1% ferric chloride solution. Formation of an intense greenish-black colour indicates presence of phenolic compounds.

2.4.6 Detection of flavonoids

2.4.6.1 Alkali reagent test

About 2 ml of diluted TS was placed in a test-tube. A few drops of NaOH 1N solution were added drop-wise in the test-tube. Increasing intensity of the yellow colour produced in the test-tube that would be colorless by addition of few drops of dilute HCl, indicates presence of flavonoids.

2.4.6.2 Shinoda/Pew test

A small amount of magnesium turnings was added into a test-tube containing about 3 ml of the TS. After addition of 3 drops of concentrated HCl in the test-tube, it was kept aside for completion of the reaction. After a while it was observed for formation of a reddish-pink or rose colour for a positive result.

2.4.7 Detection of tannins

2.4.7.1 Gelatin test

About 2 ml of TS was treated with 2 ml aqueous solution of 1% gelatin containing 10% sodium chloride. Formation of a white buff colour precipitate (a milky colour) indicates presence of tannins in the test sample.

2.4.8 Detection of saponins

2.4.8.1 Foam test

One ml of the TS after being diluted with 2 ml distilled water in a test-tube was vigorously shaken for 15 minutes. Formation of persistent foam reveals presence of saponins in the test sample.

2.4.8.2 Fontan-kendale test

To differentiate between triterpenic and steroidal saponins, 1 ml of the TS was placed into two

clean test-tubes (numbered 1 and 2), separately. Then, 5 ml of NaOH 0.1 N and 5 ml of HCl 0.1 N were mixed with the contents of test-tube No. 1 and No. 2, respectively. Both of the test-tubes were simultaneously shaken well, and were observed for the amount and stability of the foam produced in both test-tubes. Production of a persistent foam in test-tube No. 1 (containing mixture of test sample and diluted alkali) than in test-tube No 2 (containing mixture of test sample and diluted HCl) indicates presence of steroidal saponins in the sample.

2.4.9 Detection of sterols

A small amount of the dried extract was washed with petroleum ether and then with acetone. The residue was extracted with small amount (10 ml) of chloroform and the later was tested for detection of steroids.

2.4.9.1 Liebermann – Burchard's test

About 5 ml of the chloroform extract of the sample was placed in a test-tube, followed by addition of few drops of acetic anhydride. Then 2 ml of conc. H₂SO₄ were carefully added from the side of the test-tube. Formation of a brown to blood red colour at the junction of two phases indicates a positive result.

2.4.9.2 Salkowski reaction

About 5 ml of the chloroform extract was placed in a test-tube, followed by carefully addition of about 2 ml of conc. H₂SO₄ from the side of test-tube. Formation of yellow-red or brown colour ring at the junction, which turns red after a short time, indicates presence of sterols in the sample.

Results of phytochemical screening of the tested extracts are presented in Table 2.

2.5 Determination of Total Polyphenolic Content (TPC)

Total phenolic content of APCE, APPE, EACE and EAPE was determined as per Folin-Ciocalteu spectrophotometric method previously described [24,25].

Briefly; 1 ml of extract MeOH solution having concentration of 300 µg/ml, was mixed with 1 ml Folin-Ciocalteu's Reagent (FCR) diluted (1/10) with distilled water in a clean test-tube. The mixture was shaken well, and placed at room ambience for 6 minutes, followed by addition of 3

ml Na₂CO₃ 7.5% aqueous solution. Blank was concurrently prepared as per the same procedure by substituting the extract with distilled water. Aliquots of gallic acid (GA) at concentrations of 5, 10, 20, 30, 40, and 50 µg/ml in MeOH, were used for establishing a standard calibration curve. All of the mixtures were shaken well, and incubated for 60 minutes at room ambience, subsequently centrifuged at 3000 rpm for 10 minutes. The supernatants were checked for their absorbances at 765 nm wavelength against the blank, using a spectrophotometer (Shimadzu UVmini 1240). TPC of the tested samples was calculated by use of the regression formula produced in GA calibration curve, and were expressed as mg GAE/g dried extract, as shown in Table 3.

2.6 Determination of Total Flavonoid Content (TFC)

The APCE, APPE, EACE, and EAPE were screened for their TFC values, applying slightly modified Aluminum trichloride colorimetric method previously described [26,27]. Briefly; AlCl₃ solution 2% (w/v) in MeOH was freshly prepared as reagent solution. Two ml of sample extract MeOH solution (300 µg/ml) were well mixed with 2 ml of the reagent in clean test-tubes. Two ml of Rutin (RU) aliquots at concentration range of 5 – 50 µg/ml were used for establishing a standard calibration curve. All of the mixtures were shaken well, and incubated at room ambience for 30 minutes in the dark for reaction completion. Then, absorbance of the mixtures was spectrophotometrically measured at 415 nm wavelength against the blank sample consisting of the mixture of 2 ml of sample/standard solution with 2 ml MeOH without AlCl₃. The TFC values were calculated based on the regression formula produced from RU standard curve, and were expressed as mg RUE/g dried extract, as shown in Table 4.

2.7 Determination of *in Vitro* Antioxidant Activity

For determination of *in vitro* antioxidant activity of tested extracts, DPPH assay method was used, as previously described [26,28]. Ascorbic acid serial dilutions at concentration range of 0.25 – 2 µg/ml in MeOH were used as standard antioxidant compound. Accurately, 4 ml of the test extracts serial dilutions in MeOH having concentrations of 1.25, 2.5, 5, 10, 15, 20, and 25 µg/ml were separately mixed with 0.5 ml DPPH solution (0.5 mM) in MeOH. Likewise, 4 ml of

serial dilutions (0.25, 0.50, 0.75, 1.00, 1.5, and 2.0 µg/ml) of ascorbic acid in MeOH were mixed with 0.5 ml DPPH solution (0.5 mM) in MeOH, to establish a standard calibration curve. As a positive control, 4 ml of pure MeOH was mixed with 0.5 ml DPPH solution (0.5 mM). All of the resultant mixtures were thoroughly shaken, and incubated in the dark at room temperature for 30 min for reaction completion. Then, absorbance of the mixtures were screened at 517 nm wavelength against methanol as blank, using a UV-spectrophotometer (Shimadzu UVmini 1240). Radical scavenging activity (RSA) of the test extracts as well as standard ascorbic acid was calculated as percentage inhibition (% inhibition) of DPPH free radicals, applying the following formula:

$$\% \text{ inhibition} = 100 \times [(Ac - As) / Ac]$$

Where, Ac is the absorbance of control, and As is absorbance of sample/standard

Antioxidative potential of the standard ascorbic acid and tested extracts were expressed as inhibition concentration of 50% DPPH free radicals (IC₅₀), which were calculated from their plotted graph of % inhibition. Fig. 1, represents the % inhibition of DPPH free radicals by the tested extracts, while Fig. 2 represents the IC₅₀ values of the tested extracts.

2.8 Determination of Sun Protective Factor (SPF)

The SPF is most commonly used parameter for expression of sun protective capacity of sunscreens [10,29]. SPF is UV energy required to produce a minimal erythema dose (MED) on protected skin, divided by the UV energy required to produce a MED on unprotected skin. SPF is calculated using the following formula:

$$SPF = \frac{\text{Minimal erythema dose in sunscreen protected skin}}{\text{Minimal erythema dose in nonsunscreen protected skin}}$$

MED is the lowest time interval or dosage of UV light irradiation, required to produce a minimal, perceptible erythema or reddening on unprotected skin [30,31].

SPF values of tested extracts were determined by *in vitro* spectrophotometric method based on measuring transmittance of the extracts diluted solutions in MeOH. Transmittance of the extracts dilutions were measured in the range of 292.5 to 337.5 nm, at five nanometers intervals. SPF

values of extracts were calculated using the following formula adopted from previous works [32,33].

$$SPF = \frac{\sum_{292.5}^{337.5} E(\lambda) \times \varepsilon(\lambda)}{\sum_{292.5}^{337.5} E(\lambda) \times \varepsilon(\lambda) \times T(\lambda)}$$

Where E (λ) is the spectral irradiance of terrestrial sunlight; and ε (λ) is the erythema action spectrum at λ, and T (λ) is the measured transmittance at λ. The values of E (λ) and ε (λ) are constant, and derived from previous works [32]. To establish a standard calibration curve, SPF values of five different concentrations of the test extracts (20, 30, 40, 50, 70 mcg/ml for EAPE and 40, 60, 80,100, 120 mcg/ml for other extracts) were measured and their related standard curves were plotted. Regression formula of the different extracts are presented in Table 5. SPF values for the tested extracts MeOH solution (2 mg/ml) were calculated using the regression formula obtained from their plotted standard curves, and the results are shown in Table 5.

2.9 Statistical Analysis

All of the experiments were performed in triplicates (n=3) and values were expressed as Mean±SD. Statistical analyses were performed using Ms. Excel 2016 package. One-way analysis of variance (ANOVA) and Student t-test were used to assess the differences between different variables. All analyses were performed at the 5% significant level (p<0.05).

3. RESULTS AND DISCUSSION

3.1 Yield Percentages of the Extracts

The yield percentage of the extracts for *A. pseudalhagi* herb and *E. angustifolia* leaves are presented in Table1.

Table 1. Data showing the extractive % (w/w) of the plants

S. no.	Extract type	% extractive Mean±SD*
1	APCE	31.834±1.64
2	EACE	35.51±1.62
3	APPE	5.308±0.10
4	EAPE	4.475±0.08

* Mean ± Standard Deviation of the three experiments (n=3),

APCE; *A. pseudalhagi* herb crude extract,
APPE; *A. pseudalhagi* herb processed extract,
EACE; *E. angustifolia* leaves crude extract,
EAPE; *E. angustifolia* leaves processed extract

As shown in Table 1, the highest yield is recorded for EACE (35.51±1.62%), followed by APCE (31.834±1.64%), while yields % for APPE and EAPE were recorded to be 5.308±0.10 and 4.475±0.08, respectively.

As statistical analysis of the data showed, the percentage yields of the purified extracts are significantly lower than that of their relative crude extracts ($p < 0.05$). This could be attributed to elimination of some extra materials such as chlorophyll and other lipophilic substances during washing of the crude extract by petroleum ether. Such purified extracts are proved to pose more potent biological efficacy because of removal of the inert compounds thereof.

3.2 Phytochemical Screening of the Extracts

Phytochemicals present in the tested extracts are shown in Table 2.

As shown in Table 2, different classes of phytochemicals such as; carbohydrates, flavonoids, phenolic compounds and saponins were found to be present in all tested extracts. Based on qualitative phytochemical tests, *A. pseudalhagi* herb contains triterpenic saponins while *E. angustifolia* leaves contain steroidal saponins. However, further works and quantitative tests are required for accurate determination of the phytochemicals present in the tested extracts. Based on available literature, presence of diverse array of secondary

metabolites is reported in *Alhagi* species including *A. pseudalhagi* [14,34,35]. Similarly, *E. angustifolia* L. is reported to contain different compounds such as; alkaloids, flavonoids, polysaccharides, saponins, tannins, vitamins, and metal elements in different plant parts [17,18,36,37].

3.3 Total Phenolic Content (TPC) of the Extracts

In present study, extracts MeOH solution at (300 µg/ml) were screened for their TPC values, applying Folin-Ciocalteu method previously described [25,38]. TPC of tested samples were calculated from the regression equation ($y=0.0201x + 0.024$, $R^2=0.995$) obtained from GA standard curve, and were expressed in terms of GA equivalent (mg of GAE/g of dried extract), as shown in Table 3.

As shown in Table 3, the order of TPC in tested extracts was highest for EAPE > EACE > APPE > APCE. Consequently, among tested extracts, both of the EAPE and EACE exhibited highest TPC values of 203.76±2.07 and 82.64±2.57 mg GAE/g dry extract, respectively. The data showed that purified extracts revealed higher TPC values than that of crude extracts. Plants phenols, by virtue of their aromatic rings, show an intense absorption in the UV region of the spectrum [22]. Hence, they would serve as choice compounds to be incorporated in sunscreen formulations.

Table 2. Phytochemicals detected in the tested extracts

S. no.	Phytochemicals	APCE	APPE	EACE	EAPE
01	Carbohydrates				
	Molisch's test	++	+	++	+
	Resorcinol test	+++	+	++	+
02	Phenols				
	FeCl ₃ Test	++	+++	++	+++
03	Flavonoids				
	Alkali R. Test	+	+++	++	+
	Shinoda Test	+	++	++	+
04	Tannin				
	Gelatin test	++	-	++	-
05	Saponin				
	Foam test	+	++	+	-
	Fontan-Keldale test	triterpenic	triterpenic	steroidic	-
06	Sterols				
	Liebermann – B test	+	+	+	+
	Salkowski test	+	+	+	+

APCE; *A. pseudalhagi* herb crude extract, APPE; *A. pseudalhagi* herb processed extract, EACE; *E. angustifolia* leaves crude extract, EAPE; *E. angustifolia* leaves processed extract, '+' Present, '++' moderately present, '+++' intensely present, '-' absent

Table 3. Data showing the TPC and TFC values of tested extracts

S. no.	Extract type	TPC values *(mg GAE/g dried extract)
1	APCE	71.14±3.10
2	APPE	80.43±3.69
3	EACE	82.64±2.57
4	EAPE	203.76±2.07

* The values are recorded as Mean±SD, (n=3)
 APCE; *Alhagi pseudalhagi* herb crude extract,
 APPE; *A. pseudalhagi* herb processed extract,
 EACE; *Elaeagnus angustifolia* leaves crude extract,
 EAPE; *E. angustifolia* leaves processed extract

3.4 Total Flavonoid Content (TFC) of the Extracts

Determination of TFC of botanicals is of high importance in predicting biological activities, such as antioxidative potential of tested crude drugs. In present study, TFC of tested extracts were calculated from the regression equation ($y=0.023x - 0.072$, $R^2=0.9996$) obtained from RU standard curve. The TFC values were expressed in terms of rutin equivalent (mg of RUE/g of dried extract), and shown in Table 4.

As shown in Table 4, the order of TFC of tested extracts was highest for EAPE > APPE > APCE > EACE, respectively. TFC of the tested extracts

varied from 33.24±3.23 to 135.85±3.83 mg RUE/g dried extract. The highest flavonoids content (135.85±3.83 mg RUE/g extract) was recorded for EAPE.

Table 4. Data showing TFC values of tested extracts

S. no.	Extract type	TFC values *(mg RUE/g dried extract)
1	APCE	59.61±3.40
2	APPE	121.06±3.37
3	EACE	33.24±3.23
4	EAPE	135.85±3.83

* The values are recorded as Mean±SD, (n=3)
 APCE; *Alhagi pseudalhagi* herb crude extract,
 APPE; *A. pseudalhagi* herb processed extract,
 EACE; *Elaeagnus angustifolia* leaves crude extract,
 EAPE; *E. angustifolia* leaves processed extract

3.5 In vitro Antioxidant Activity of the Plant Extracts

Evaluation of antioxidative potential of botanicals serves as a choice approach to judge their pharmaceutical as well as pharmacological values. In present work, sample extracts at concentration range of 2.5 – 25 µg/ml in MeOH were evaluated for their DPPH free radical scavenging activity. Fig. 1 represents the % inhibition of DPPH free radicals by the tested extracts.




Fig. 1. Graph showing % inhibition activity of tested extracts in DPPH assay
 EAPE; *E. angustifolia* leaves processed extract, EACE; *E. angustifolia* leaves crude extract,
 APPE; *A. pseudalhagi* herb processed extract, and APCE; *A. pseudalhagi* herb crude extract

As shown in Fig. 1, the RSA of the tested samples was dose-dependent. That is to say, the % inhibition of DPPH free radicals were increasing by increase in concentration of test extracts solutions used in the assay.

The antioxidant potentials of standard ascorbic acid and tested extracts were expressed as their IC₅₀ values. IC₅₀ actually refers to the concentration (µg/ml) of standard or sample extracts required for inhibition of 50% DPPH radicals used in the assay. Lower IC₅₀ value of a test sample indicates higher RSA and higher antioxidant activity of the same sample [26,27,39].

In vitro antioxidative potential of tested extracts were expressed as their IC₅₀ values which were calculated from their plotted graphs of % inhibition vs. concentrations (Fig. 1). Fig. 2 represents the calculated IC₅₀ values for both ascorbic acid and the tested extracts.

As shown in Fig. 2, IC₅₀ for ascorbic acid is 2.32 µg/ml, while IC₅₀ of the tested extracts are in range of 4.5 – 21.36 µg/ml. Among the tested extracts, EAPE exerts significantly higher free RSA (IC₅₀ = 4.5, and *p* < 0.05), while RSA of the remaining extracts decreased in order of APPE>EACE>APCE with IC₅₀ values of 14.31<14.36<21.36, orderly.

In DPPH assay, the stronger activity of EAPE could be attributed to its purity from other inert phytochemicals such as chlorophyll, etc. and its richness in phenolics compounds. Therefore, a

direct correlation exists between TPC and DPPH radical scavenging potential of EAPE. Our data were in compliance with those of many previous works which reported positive correlation between DPPH radical scavenging activities and TPC values of tested plant extracts [26,40]. As per another study, high phenolic content and strong antioxidant properties of ethylacetate fraction of defatted methanolic extract of *Allium* species was reported [41]. In another study, methanol extracts of cinnamon and cloves exhibited strong antioxidant properties owing to their high phenolic and flavonoids content [38]. Positive correlation between TPC and *in vitro* antioxidant potential of *Aloe vera* leaves is also reported [42].

3.6 Sun Protective Factor (SPF) of the Extracts

SPF values of the tested extracts' MeOH solution (2 mg/ml) are presented in Table 5.

As per the results, the SPF values were between 5.47 ± 0.03 and 29.73 ± 0.3. As data showed, both of EAPE and APPE exhibited higher SPF than EACE and APCE. This issue could be attributed to the purity of EAPE and APPE from excess of chlorophyll and other inert compounds. The higher amounts of both polyphenols and flavonoid compounds in EAPE and APPE, and elimination of their inert chemicals, are claimed to be the reasons for their increased UV absorption. Compared to SPF value of APPE (17.83 ± 0.88), the EAPE revealed significantly higher SPF value of 29.73 ± 0.31 (*p* < 0.05).




Fig. 2. Graph showing the IC₅₀ values for the standard and tested extracts
 Vit. C; vitamin C, EAPE; *E. angustifolia* leaves processed extract, EACE; *E. angustifolia* leaves crude extract, APPE; *A. pseudalhagi* herb processed extract, and APCE; *A. pseudalhagi* herb crude extract

Table 5. Data showing regression equations and SPF value for the tested extracts 2 mg/ml MeOH solutions

S. no.	Extract type	Regression equation & R ²	SPF* (Conc. 2 mg/ml)
1	APCE	y = 2.5048x – 2.7942 (R ² =0.93)	5.47 ± 0.03
2	APPE	y = 14.614x – 22.361 (R ² =0.81)	17.83 ± 0.88
3	EACE	y = 4.8386x – 6.5119 (R ² =0.87)	10.69 ± 0.16
4	EAPE	y = 15.201x – 20.306 (R ² =0.95)	29.73 ± 0.31

* Data were recorded as Mean±SD, (n=3)

APCE; *Alhagi pseudalhagi* herb crude extract, APPE; *A. pseudalhagi* herb processed, EACE; *E. angustifolia* leaves crude extract, EAPE; *E. angustifolia* leaves processed extract

The *In vitro* spectrophotometric method used in present study, was based on measuring transmittance of diluted solution of the tested extracts. Since botanical extracts contain a wide range of phytochemicals, such as: phenolic compounds, flavonoids, carotenoids, vitamin E, vitamin C and etc., hence these compounds are responsible for UV absorption and sun protective properties of botanicals [10,29,43].

In present study, positive correlations were found between DPPH radical scavenging capacity and SPF values of the tested extracts. Likewise, there were positive correlations between SPF and TPC, and also between SPF and TFC of the tested extracts, as shown in Table 6.

As shown in Table 6, the strongest linear correlation ($r = -0.9603$) was found between SPF and IC₅₀ values of the tested extracts. While the negative value of r is because of the inverse relationship between IC₅₀ and SPF values of the extracts. It indicates if smaller is the value of IC₅₀ of a sample, greater would be the SPF value of the same sample, and vice versa. However, phenolics and flavonoids are very well known for their radicals scavenging potential as potent natural antioxidants [27,39].

Table 6. The correlation coefficients (r values) obtained between SPF values and IC₅₀, TPC and TFC of tested extracts

The values of pearson correlation coefficients	
SPF vs. IC ₅₀	$r = -0.9603$
SPF vs. TPC	$r = 0.8999$
SPF vs. TFC	$r = 0.8476$

* Data were calculated using Ms excel 2016 Data analysis tool pack

SPF; Sun Protective Factor, IC₅₀; Inhibition concentration of 50% of DPPH radicles, TPC; Total Phenolic content, TFC; Total Flavonoids content, r = Pearson correlation coefficient

Overall, our findings were in compliance with many previous works, which have reported that both phenolics and flavonoid compounds act as UV absorbing agents [10,32,44]. In addition, phenolics and flavonoids as strong antioxidant agents, play important roles in prevention of UV induced damage such as; photo aging and skin cancers [20,45]. Recently, antioxidants present in botanicals and marine organisms are deemed to serve as compounds of choice for developing natural sunscreen formulations [10,11].

4. CONCLUSION

Considering the huge demand for safe and effective plants derived UV-filters, *A. pseudalhagi* herb and *E. angustifolia* leaves were screened for their antioxidant potential and sun protective factor. Among the tested extracts, purified extract of *E. angustifolia* leaves being rich in phenolics and flavonoids, demonstrated strong DPPH radical scavenging activity and a high SPF of 29.73 ± 0.31 for its MeOH solution (2 mg/ml). In comparison with *E. angustifolia* leaves, *A. pseudalhagi* herb purified extract revealed lower contents of phenolics and flavonoids, and hence produced greater IC₅₀ value (14.31) and lower SPF (17.83 ± 0.88). As per data found in present study, *E. angustifolia* leaves could be recommended as natural source of the sun protective agents beneficial in developing potent herbal sunscreen formulations.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENTS

The authors gratefully acknowledged Family Health International (FHI)-360, USAID for financial support (under Cooperative Agreement

No. AID-306-A-13-00009-00), and Ministry of Higher Education, Islamic Republic of Afghanistan for providing facilities to conduct this research. The content of this publication does not necessarily reflect the views, analysis or policies of FHI-360 or USAID, nor does any mention of tradenames, commercial products, or organizations imply endorsement by FHI 360 or USAID.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Beyene B, Beyene B, Deribe H. Review on application and management of medicinal plants for the livelihood of the local community. *J Resources Devel and Manag.* 2016;22:33-39.
2. Uritu CM, Mihai CT, Stanciu GD, Dodi G, Alexa-Stratulat T, Luca A, Leon-Constantin M, Stefanescu R, Bild V, Melnic S, Tamba BI. Medicinal plants of the Family Lamiaceae in pain therapy: A review. *Pain Res Manag.* 2018;1-44.
3. Dong-Ping X, Li Y, Meng X, Zhou T, Zhou Y, Zheng J, Zhang JJ, Li HB. Natural antioxidants in foods and medicinal plants: Extraction, assessment and resources. *Int J Mol Sci.* 2017;1-32.
4. Cavinato M, Waltenberger B, Baraldo G, Grade CVC, Stuppner H, Jansen-Durr P. Plant extracts and natural compounds used against UVB-induced photoaging. *Biogerontology.* 2017;18:499-516.
5. Raju GS, Moghal MMR, Dewan SMR, Amin MN, Billah MM. Characterization of phytoconstituents and evaluation of total phenolic content, anthelmintic, and antimicrobial activities of *Solanum violaceum* Ortega. *Avicenna J Phytomed.* 2013;3(4):313-320.
6. Subramanya SB, Venkataraman B, Meeran MFN, Goyal SN, Patil CR, Ojha S. Therapeutic potential of plants and plant derived phytochemicals against acetaminophen-induced liver injury. *Int J Mol Sci.* 2018;1-43.
7. Chanchal D, Swarnlata S. Herbal photoprotective formulations and their evaluation. *The Open Nat Prod J.* 2009;2: 71-76.
8. Nunes AR, Rodrigues ALM, Queiroz DB, Vieira IGP, Neto JFC, Junior JTC, Tintino SR, Morais SM, Coutinho HDM. Photoprotective potential of medicinal plants from Cerrado biome (Brazil) in relation to phenolic content and antioxidant activity. *J Photochem Photobiol B: Biol.* 2018.
DOI: 10.1016/j.jphotobiol.2018.10.013
9. Tienaho J, Poikulainen E, Sarjala T, Muilu-Mäkelä R, Santala V, Karp M. A bioscreening technique for ultraviolet irradiation protective natural substances. *Photochem Photobiol.* 2018;94(6):1273-1280.
10. Muzaffer U, Paul VI, Prasad NR, Karthikeyan R, Agilan B. Protective effect of *Juglans regia* L. against ultraviolet B radiation induced inflammatory responses in human epidermal keratinocytes. *Phytomedicine.* 2018;42:100-111.
11. Kostyuk V, Potapovich A, Albuhaydar AR, Mayer W, Luca CD, Korkina L. Natural substances for prevention of skin photoaging: Screening systems in the development of sunscreen and rejuvenation cosmetics. *Rejuvenation Res.* 2018;21(2):91-101.
12. Amini MH, Hamdam SM. Medicinal plants used traditionally in Guldara district of Kabul, Afghanistan. *Int J Pharmacogn Chinese Med.* 2017;1(3):000118.
13. Srivastava B, Sharma H, Dey YN, Wanjari MM, Jadhav AD. *Alhagi pseudalhagi*: A review of its phyto-chemistry, pharmacology, folklore claims and Ayurvedic studies. *Int J Herb Med.* 2014;2(2):47-51.
14. Varshochi F, Asadollahi K. *Alhagi*: Traditional and modern medicine effective against kidney stones. *J Nephrofarmacol.* 2017;6(1):15-16.
15. Breckle SW, Rafiqpoor MD. *Field Guide Afghanistan: Flora and vegetation.* Germany: Scientia Bonnensis; 2010.
16. Wang B, Qu H, Ma J, Sun X, Wang D, Zheng Q, Xing D. Protective effects of *Elaeagnus angustifolia* leaf extract against myocardial ischemia/reperfusion injury in isolated rat heart. *J Chem.* 2014;1-6.
17. Hamidpour R, Hamidpour S, Hamidpour M, Shahlari M, Sohraby M, Shahlari N, Hamidpour R. Russian olive (*Elaeagnus angustifolia* L.): From a variety of traditional medicinal applications to its novel roles as active antioxidant, anti-inflammatory, anti-mutagenic and analgesic agent. *J Tradit Complement Med.* 2016;1-6.

18. Incilay G. Volatile composition, antimicrobial and antioxidant properties of different parts from *Elaeagnus angustifolia* L. J Essent. Oil-Bear Plants. 2014;17(6): 1187-1202.
19. Okmen G, Turkcan O. The antibacterial activity of *Elaeagnus angustifolia* L. against mastitis pathogens and antioxidant capacity of the leaf methanolic extracts. J Anim Vet Adv. 2013;12(4):491-496.
20. Jarzycka A, Lewinska A, Gancarz R, Wilk KA. Assessment of extracts of *Helichrysum arenarium*, *Crataegus monogyna*, *Sambucus nigra* in photoprotective UVA and UVB; photostability in cosmetic emulsions. J Photochem Photobiol B: Biol. 2013;128:50-57.
21. Evans WC. Trease and evans pharmacognosy, 16th Ed. London: Saunders; 2009.
22. Harborne JB. Phytochemical methods: A guide to modern techniques of plant analysis. 3rd Ed. India: Springer; 1998.
23. Satheesh KB, Suchetha KN, Bhat VS, Sharmila KP, Bekal MP. Preliminary phytochemical screening of various extracts of *Punica granatum* pell, whole fruit and seeds. Nitte University J Health Sci. 2012;2(4):34-38.
24. Blois MS. Antioxidant determinations by the use of a stable free radical. Nature. 1958;1199-1200.
25. Singleton VL, Orthofer R, Lamuela-Raventos M. Analysis of total phenols and other oxidation substances and antioxidants by means of Folin-Ciocalteu Reagent. Methods Enzymol. 1999;299: 152-178.
26. Amini MH, Kalsi V, Kaur B, Khatik GL, Lobo R, Singh G, Agarhari UC, Yele S, Sutte A. Phytochemical screening and antioxidant activity of *Heracleum afghanicum* Kitamura leaves. Res J Pharm Technol. 2017;10(10):3498–3502.
27. Borkatoky M, Kakoty BB, Saikia LR. Influence of total phenolic content and total flavonoid content on the DPPH radical scavenging activity of *Eclipta alba* (L.) Hassk. Int J Pharm Pharm Sci. 2013;5(Suppl. 1):224-327.
28. Amiri H. Essential oils composition and antioxidant properties of three *Thymus* species. Evid Based Complement Alternat Med. 2012;1-8.
29. Salvador A, Chisvert A, (Eds.). Analysis of cosmetic products. 1st Ed. Amsterdam: Elsevier; 2007.
30. Conant L, Beck KM, Liao W. A rapid and cost-effective device for testing minimal erythema dose. Dermatol Ther (Heidelb). 2018;1-7.
31. Dutra EA, Oliveira DACC, Hackmann ERMK, Santoro MIRM. Determination of sun protection factor (SPF) of sunscreens by ultraviolet spectrophotometry. Brazil J Pharm Sci. 2004;40(3):381-385.
32. Ebrahimzadeh MA, Enayatifard R, Khalili M, Ghaffarloo M, Saeedi M, Charati JY. Correlation between sun protective factor and antioxidant activity, phenol and flavonoid contents of some medicinal plants. Iran J Pharm Res. 2014;13(3): 1041-1047.
33. Khazaeli P, Mehrabani M. Screening of sun protective activity of the ethyl acetate extracts of some medicinal plants. Iran J Pharm Res. 2008;7(1):5-9.
34. Asghari MH, Fallah M, Moloudizargari M, Mehdikhani F, Sepehrnia P, Moradi B. A systematic and mechanistic review on the phytopharmacological properties of Alhagi Species. Anc Sci Life. 2016;36(2): 65-71.
35. Muhammad G, Hussain MA, Anwar F, Ashraf M, Gilani AH. *Alhagi*: A plant genus Rich in bioactives for pharmaceuticals. Phytother Res. 2014;29(1):1-13.
36. Khan SU, Khan A, Shah AA, Shah SM, Hussain S, Ayaz M, Ayaz S. Heavy metals content, phytochemical composition, antimicrobial and insecticidal evaluation of *Elaeagnus angustifolia*. Toxicol Ind Health. 2013;1-8.
37. Sun N, Liu H, Liu x, Zhang Y, Liu X, Wang S, Xu X, Tian W. Immunological activities of polysaccharide extracted from *Elaeagnus angustifolia* L. CYTA J Food. 2018;16(1):995-1002.
38. Gupta D. Comparative analysis of spices for their phenolic content, flavonoid content and antioxidant capacity. Am Int J Res Form Appl Nat Sci. 2013;4(1):38-42.
39. Valyova M, Stoyanov S, Yuliana M, Ganeva Y. Evaluation of *in vitro* antioxidant activity and free radical scavenging potential of variety of *Tagetes erecta* L. flowers growing in Bulgaria. Int J Appl Res Nat Prod. 2012;5(2):19-25.
40. Mazandarani M, Majidi Z, Moghaddam PZ, Abrodi M, Hemati H, Fathiazad F. Essential oil composition, total phenol, flavonoid, anthocyanin and antioxidant activities in different parts of *Artemisia*

- annua* L. in two localities (North of Iran). J Med Plants By-prod. 2012;1:13-21.
41. Abdel-Gawad M, Abdel-Aziz M, El-Sayed M, El-Wakil E, Abdel-Lateef E. *In vitro* antioxidant, total phenolic and flavonoid contents of six *Allium* species growing in Egypt. J Microbiol Biotechnol Food Sci. 2014;3(4):343-346.
42. Kumar S, Yadav A, Yadav M, Yadav JP. Effect of climate change on phytochemical diversity, total phenolic content and *in vitro* antioxidant activity of *Aloe vera* (L.) Burm.f. BMC Res Notes. 2017;1-12.
43. Korac RR, Khambholja KM. Potential of herbs in skin protection from ultraviolet radiation. Pharmacog Rev. 2011;5(10): 164-173.
44. Roh E, Kim JE, Kwon JY, Park JS, Bode AM, Dong Z, Lee KW. Molecular mechanisms of green tea polyphenols with protective effects against skin photoaging. Crit Rev Food Sci Nut. 2017;57(8):1631-1637.
45. Kim SH, Jung EY, Kang DH, Chang UJ, Hong YH, Suh HJ. Physical stability, antioxidative properties, and photo-protective effects of a functionalized formation containing black garlic extract. J Photochem Photobiol B: Biol. 2012;117: 104-110.

© 2019 Amini et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/51341>