



CaO Nanoparticles Seed Invigoration on Germination and Seedling Growth of Rice var. TRY 3

S. Gopinath ^{a++*}, K. Raja ^{b#}, V. Manonmani ^{c†‡},
P. Jeyaprakash ^{a†‡}, T. Ramesh ^{d#‡}, C. Vanniarajan ^{e^}
and S. Kavitha ^{f#}

^a Department of Seed Science and Technology, Tamilnadu Agricultural University, Coimbatore, India.

^b Department of Plant Breeding and Genetics, ADAC&RI, Trichy, India.

^c Department of Seed and Technology, TNAU, Coimbatore, India.

^d Department of Agronomy, ADAC&RI, Trichy, India.

^e ADAC&RI, Trichy, India.

^f Seed Centre, TNAU, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i102877

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105869>

Original Research Article

Received: 26/06/2023

Accepted: 31/08/2023

Published: 04/09/2023

ABSTRACT

Rice (*Oryza sativa* L.), a vital cereal and staple food crop in South East Asia. It faces significant challenges in cultivation worldwide, due to soil salinity. Seed enhancement techniques like priming have shown resilience against salt stress. In this study, the potential of nano-priming with calcium

⁺⁺ M.Sc Scholar;

[#] Associate Professor;

[†] Professor;

[‡] Head;

[^] Dean;

*Corresponding author: E-mail: gopinathkk98@gmail.com;

oxide nanoparticles (CaO NPs) to enhance rice seed germination and growth was investigated. CaO NPs were synthesized from organic waste (eggshells) using a sol-gel technique and characterized using Raman and UV-Vis spectroscopy. Paddy seeds were primed with varying concentrations of CaO NPs, and their germination and seedling growth were assessed. The results showed that nano-priming with 40 ppm of CaO NPs significantly improved germination, root and shoot length, dry matter production, and seedling vigour as compared to the control. However, higher concentrations of CaO NPs had a detrimental effect on germination. The study suggests that CaO NP seed priming has the potential to enhance rice germination and seedling development.

Keywords: Rice; calcium oxide nanoparticle; nano-priming; seedling vigour.

1. INTRODUCTION

Rice (*Oryza sativa* L. 2n=24) important cereal and staple food crop in South East Asia. Worldwide China is the leading producer of rice followed by India. In India 43 million hectares are under rice cultivation with an average yield of 2.56 tons/ha. Tamilnadu ranks second in rice production next to Punjab, and accounts 2.2 million ha under rice production with an average productivity of 2.8 tonnes/ha. Water constraint, imbalanced fertilizer and increasing soil salinity/alkalinity are the major constraints for rice production. Worldwide, Particularly, Bangladesh, China and India have significant challenges in cultivation due to soil salinity. It is the second most prevalent issue after drought. The sensitivity of rice to salt, especially during the seedling and reproductive stages, greatly affects its growth and productivity [1, 2]. Soil salinity above EC 4 dSm⁻¹ is moderate for rice, while salinity exceeding 8dSm⁻¹ is considered severe [3]. Salinity can lead to various morphological, physiological, and biochemical changes in rice plants, including severe cases of plant mortality [4]. Despite efforts in classical breeding to enhance rice's resistance to salt stress, success has been limited, and most current rice varieties are unable to tolerate high levels of salinity [5, 6].

Use of quality seeds [7] and quality enhancement technique such as seed priming are viable techniques, aimed at altering the physiological and biochemical properties of seeds to enhance their tolerance against abiotic stress. It involves inducing extensive physiological reorganization through hydration and dehydration processes. Priming leads to various physicochemical changes that modify the protoplasmic characteristics and enhance the physiological activity of the embryo and associated structures. As a result, the seeds exhibit improved water absorption, facilitated by increased cell wall elasticity, and the development of a stronger and more efficient root system [8]. Seed priming with

carbon nanoparticles (CNPs) and chitosan (CS) helps in enhancing seedling growth and preserving pigment levels. Further, there are scientific reports evidenced the effect of NPs priming on improved germination and seedling growth [9] and [10].

Nano-priming of paddy seeds with calcium oxide nanoparticles can result in several potential effects. Firstly, it has been observed that such priming can enhance the germination process, promoting faster and more efficient seed germination [11]. CaO NPs stimulate metabolic activity, leading to increased nutrient uptake and root development, ultimately resulting in enhanced seedling vigour [12]. Moreover, calcium oxide nanoparticle priming has the potential to enhance stress tolerance in paddy plants. Studies have demonstrated that nano-priming with calcium oxide nanoparticles can confer improved resistance to abiotic stresses such as drought and salinity [13]. By enhancing the plants' ability to withstand adverse environmental conditions, nano-priming can contribute to better survival rates and overall growth performance. This study aimed to reveal effect of CaO NPs on improved germination growth of rice.

2. MATERIALS AND METHODS

CaO nanoparticles were synthesized via sol-gel technique from organic waste (egg shells) [14]. CaO NPs were characterized for Raman and UV-Vis [15]. Paddy seeds of var. TRY 3 were surface sterilized with 1% sodium hypochlorite, and primed with CaO nanoparticles at 10ppm, 20ppm, 30ppm, 40ppm, 50ppm 60ppm, 70ppm, 80ppm, 90ppm and 100ppm solutions at 1:1 ratio (seed: solution) for 12hr, and seeds were shaded dried back to original moisture content. The CaO NPs primed seeds were evaluated for germination and seedling growth under controlled environment.

2.1 Germination Percentage

The germination test for NPs primed paddy seeds was conducted following the ISTA guidelines, utilizing the roll towel method in germination chamber set at a constant temperature of $25\pm 2^\circ\text{C}$ and a relative humidity of 95 ± 2 per cent. On the 14th day, the number of normal seedlings, abnormal seedlings, hard seeds, and dead seeds were recorded. The mean germination percentage was then calculated based on the number of normal seedlings.

2.2 Seedling Length

During the final count, ten random normal seedlings were selected from each replication of treatments. The measurement taken was the length from the tip of the main leaves to the bottom of the primary root, which is referred to as the "seedling length." These measurements were recorded in centimeters. Mean values were then calculated based on the measurements of these ten normal seedlings for each treatment.

2.3 Dry Matter Production

Dry matter production was determined by selecting ten normal seedlings for measurement. After removing the cotyledon and seed coat, the seedlings were carefully folded and placed inside a paper cover. They were then subjected to shade drying for 24 hours followed by further drying in a hot air oven at 80 degrees Celsius for an additional 4 hours. After this, the seedlings were allowed to cool in a desiccator for 30 minutes. Using an electronic weighing balance, the weight of the seedlings dried in the hot air oven was recorded. This weight measurement was used to calculate the mean value, expressed in grams per 10 seedlings.

2.4 Vigour Index

The vigor index of the seedlings was determined following the procedure described by [16]. This index was calculated by multiplying the germination percentage with the total seedling length, which includes both the root and shoot lengths. The resulting vigor index values were calculated and presented as whole numbers.

$$\text{Vigor Index} = \text{Germination (\%)} \times \text{Total Seedling Length (cm)}$$

2.5 Statistical Analysis

The statistical analysis of the experimental data involved the use of analysis of variance (ANOVA)

as a factorial combination of treatments. Mean values were then compared and separated based on the least significant difference (LSD) only if the F test of ANOVA for treatments showed significance at the 0.05 probability level. Prior to the analysis, the percentage data was transformed using the arcsine transformation. In cases where the F test was found to be non-significant, the notation "NS" was used to indicate this result.

3. RESULTS AND DISCUSSION

3.1 Raman Spectroscopy of Organic Waste (Eggshell) Derived CaO

The presence of oxide species was successfully detected by using raman spectroscopy technique. The characteristics bands at 281 cm^{-1} , 358 cm^{-1} , $1,358\text{ cm}^{-1}$, $1,675\text{ cm}^{-1}$ and 963 cm^{-1} conforming to a function of the calcium oxide content in the sample. The peaks at 281 cm^{-1} in the spectrum of eggshell-derived CaO confirmed the lattice vibration region. A function assigned to the Ca-O bonds is the sharp band at 358 cm^{-1} in the spectra of ES-CaO (Fig 1). According to a chemical perspective, the highly reactive calcium oxide formed from eggshell powder exposed to ambient air is responsible for the incidence of substantial peaks in the spectra for carbonate anion. The increased Ca^{2+} content of the equivalent carbonates that decomposed at high temperatures due to the calcination of powdered eggshell and produced highly reactive CaO can be used to explain the persistence of the Raman signal found in the ES-CaO spectrum [15].

3.2 UV-VIS Spectroscopy of Organic Waste (Eggshell) Derived CaO

With wavelengths ranging from 200 to 800 nm, the UV-visible absorbance and reflectance spectra of powdered CaO made from chicken eggshells were captured. It was determined by the perceived spectrum that the CaO made from eggshell has enough transmission throughout the IR and visible spectrum. This result is in accordance with [15, 17] (Fig. 2).

3.3 Germination Percentage

Germination and growth attributes were observed between hydroprimed seeds and nanoprimed seeds. Seeds were primed with various concentrations of CaO nanoparticles and observed for germination and growth parameters along with a control group. The results demonstrated that, on average, seeds primed

with 40ppm nanoparticle concentration exhibited the highest germination rate of 98 percent, whereas the control group only showed 86 percent germination (Fig. 3). Nanoprimering with 40ppm CaO NP shows 13.9 per cent increase in germination.

Table 1. Effect of seed nano priming with different concentrations of CaO nanoparticles on germination percentage and growth parameters of paddy

Priming treatments	Germination per cent	Root length	Shoot length	Dry matter production	Vigour index
Control	86(70)	10.74	18	0.29	2473
Hydropriming	88(71)	10.99	18.03	0.30	2533
10ppm	88(71)	11.003	18.07	0.36	2562
20ppm	92(78)	13.33	18.55	0.45	2937
30ppm	94(80)	13.96	19.72	0.69	3164
40ppm	98(84)	14.80	20.19	1.27	3426
50ppm	94(75)	13.15	19.54	0.46	3074
60ppm	92(71)	12.64	18.85	0.42	2897
70ppm	90(73)	12.48	18.72	0.41	2810
80ppm	88(72)	12.32	18.31	0.41	2695
90ppm	88(72)	12.11	17.5	0.36	2609
100ppm	82(66)	11.77	17.12	0.36	2367
Mean	90(74)	12.44	18.55	0.48	2796
SEd	2.1554	0.3243	0.4810	0.0143	73.2730
CD(0.05)	4.4485	0.6693	0.9927	0.0295	151.2284

Data in the parenthesis () denotes arc sine transformed values





Fig. 1. Raman spectroscopy analysis of CaO NPs



UV-visible spectra of CaO prepared from waste eggshells transmittance

Fig. 2. UV- visible spectra analysis of CaO NPs




Fig. 3. Effect of seed nano priming with different concentrations of CaO nanoparticles on germination and vigour index




Fig. 4. Effect of seed nano priming with different concentrations of CaO nanoparticles on seedling length and dry matter production

The observed increase in germination can be attributed to the effective regulation of ion homeostasis (Na⁺/K⁺), as proposed by [18]. Nanopriming seems to facilitate increased water uptake by penetrating the seed coat and creating more channels for water entry [19]. This accelerated water uptake enhances α-amylase

activity, which, in turn, increases the amount of soluble sugars, thus promoting respiratory metabolism for seed germination, as suggested by [20]. Additionally, the elevated level of Ca²⁺ triggered by nanopriming synthesizes polyamines, which have been shown to improve seed germination, as discussed by [21].

However, treatment with higher concentration of CaO NP showed reduction in germination percentage. Treatment with 100ppm CaO NP reduced germination by 4.65 per cent compared to control. Elevated calcium levels within seeds can disrupt the balance of ions, thereby influencing the water imbibition process. The presence of excessive calcium can result in the formation of insoluble calcium pectate, which negatively affects water uptake and impedes the expansion of cell walls crucial for seed germination [22]. This phenomenon may explain the gradual decrease in germination observed as the concentration of CaO nanoparticles increases.

3.4 Seedling Length

CaO nanoparticle at 40ppm recorded significantly higher root length (14.80). Priming with 40ppm CaO NP recorded approx. 38 per cent increase in root length as compared to control (Fig. 4).

Seed priming using CaO NP had a notable impact on the shoot length. Seeds treated with 40 ppm of CaO NP demonstrated a considerable increase in shoot length (20.19), surpassing the control group by 12 percent (Fig. 4). This, in turn, lowers the root osmotic potential, hindering water absorption and transportation to the shoots. As a consequence, embryo expansion and seedling emergence are adversely affected [23]. In sorghum, there was a drastic increase in shoot length when Ca^{2+} was supplied to seeds [24]. As the concentration of CaO increases, there is a consistent reduction in the length of seedlings. For instance, at a CaO concentration of 100 ppm, the seedling shoot length measured 17.12 cm, the shoot length decreased by 15 per cent compared to the shoot length observed at 40 ppm CaO concentration. In sorghum, at higher levels of Ca^{2+} there was a linear decrease in seedling growth [24]. The CaO NPs treated plants increase the shoot length and showed a positive impact on seedling development [25], and [26] reported that CaO NPs treated plants increasing the stem firmness and decreasing stem bending *Gerbera (Gerbera jamesonii)*.

3.5 Dry Matter Production

Seed priming with CaO NPs, as increased the dry matter production of seedlings. CaO at 40ppm has highest dry matter production of 1.27g/10 seedlings, as compared to control

which as dry matter production of 0.29 g/10 seedlings (Fig. 4).

3.6 Vigour Index of Seedling

Seedling vigour has been observed to increase by 38.5 per cent compared to control. Seedling vigour for control was 2473 while for 40ppm nanoparticle concentration it was 3426 (Fig. 3). The increase in vigour can be attributed to higher and faster germination, and better seedling growth. Calcium is known to affect cell division and growth, which may have attributed to higher seedling growth, thus recorded increase in vigour [27].

4. CONCLUSION

The study presented here focused on seed nanopriming as a potential enhancement technique for improving the germination and growth parameters of rice seedlings. Employing calcium oxide nanoparticles (CaO NP) for priming exhibited promising results. The germination, root and shoot growth, and overall vigour of rice seedlings were positively influenced by CaO NP priming, particularly at a concentration of 40 ppm. This nanoparticle-mediated seed priming strategy holds significant potential to enhance ion homeostasis, facilitate water uptake, and trigger beneficial physiological responses, leading to improved germination and seedling development.

By harnessing the capabilities of nanotechnology and seed priming techniques, there is a potential avenue for enhancing the growth and yield of rice. Nonetheless, while the results are promising, additional studies and field trials are essential to validate the effectiveness of CaO NP seed priming across diverse rice varieties and real-world agricultural conditions. Moreover, a comprehensive understanding of the underlying molecular mechanisms driving the observed improvements in germination and growth attributes is necessary to fine-tune and optimize this technique. With concerted efforts in research and application, seed priming using nanotechnology could potentially revolutionize rice cultivation and ultimately contributing to global food security and sustainable agriculture.

ACKNOWLEDGEMENT

The authors highly acknowledge fellowship provided by ICAR- RF for the study. The author also thank Anbil Dharmalingam Agricultural college and Research Institute, Department of

Seed Science & technology and Centre for Nanotechnology for their support during the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Lutts S, Kinet J, Bouharmont J. Changes in plant response to NaCl during development of rice (*Oryza sativa* L.) varieties differing in salinity resistance. *Journal of Experimental Botany*. 1995;46(12):1843-1852.
2. Khan MA, Abdullah Z. Salinity–sodicity induced changes in reproductive physiology of rice (*Oryza sativa*) under dense soil conditions. *Environmental and Experimental Botany*. 2003;49(2):145-157.
3. Ghassemi F, Jakeman AJ, Nix HA. Salinisation of land and water resources: 429 human causes, extent, management and case studies. CABI Publishing, Wallingford. 1995;526.
4. Gupta S, Schillaci M, Walker R, Smith PM, Watt M, Roessner U. Alleviation of salinity stress in plants by endophytic plant-fungal symbiosis: Current knowledge, perspectives and future directions. *Plant and Soil*. 2021;461:219-244.
5. Rabbani G, Rahman A, Mainuddin K. Salinity-induced loss and damage to farming households in coastal Bangladesh. *International Journal of Global Warming*. 2013;5(4): 400-415.
6. Solis CA, Yong MT, Vinarao R, Jena K, Holford P, Shabala L, Chen ZH. Back to the wild: On a quest for donors toward salinity tolerant rice. *Frontiers in Plant Science*. 2020;11: 323.
7. Jerlin R, Menaka C, Raja K, Moorthy KR, Tamilkumar P. Standardization of sieve size for grading of olitorius jute seeds. *Asian Journal of Agricultural Research*, 2010;4(1):15-19.
8. Krishnasamy V, Srimathi P. Seed Management of rainfed agriculture. Land and use planning and watershed management in Rainfed Agriculture (ed.) Balusamy, M., CR Chinnamuthu and A. Velayutham. Centre of Advanced studies. Department of Agronomy, Tamilnadu Agricultural University, Coimbatore. 2001;140.
9. Raja K, Sowmya R, Sudhagar R, Sathya MP, Govindaraju K, Subramanian KS. Biogenic ZnO and Cu nanoparticles to improve seed germination quality in blackgram(*Vigna mungo*). *Materials Letters*. 2019;235:164-167.
10. Mukiri C, Raja K, Senthilkumar M, Subramanian KS, Govindaraju K, Pradeep D, Ranjan S. Immobilization of beneficial microbe *Methylobacterium aminovorans* in electrospun nanofibre as potential seed coatings for improving germination and growth of groundnut *Arachis hypogaea*. *Plant Growth Regulation*. 2021;1-9.
11. Dhage S, Biradar DP. Nanofertilizers: Perspective to enhance growth, yield and NUE of crops. *Indian Journal Pure and Applied Biosciences*. 2020;8(6):339-49.
12. Alam A, Ullah H, Cha-um S, Tisarum R, Datta A. Effect of seed priming with potassium nitrate on growth, fruit yield, quality and water productivity of cantaloupe under water-deficit stress. *Scientia Horticulturae*. 2021;288:110354.
13. Zhang W, et al. Silicon promotes growth and root yield of *glycyrrhiza uralensis* under salt and drought stresses through enhancing osmotic adjustment and regulating antioxidant metabolism. *Crop Protection*. 2018;107:1-11.
14. Habte L, Shiferaw N, Mulatu D, Thenepalli T, Chilakala R, Ahn JW. Synthesis of nano-calcium oxide from waste eggshell by sol-gel method. *Sustainability*. 2019;11(11): 3196.
15. Jaiswal KK, Dutta S, Pohrmen CB, Verma R, Kumar A, Ramaswamy AP. Bio-waste chicken eggshell-derived calcium oxide for photocatalytic application in methylene blue dye degradation under natural sunlight irradiation. *Inorganic and Nano-Metal Chemistry*. 2020;51(7):995-1004.
16. Abdul-Baki AA, Anderson JD. Vigor determination in soybean seed by multiple criteria 1. *Crop science*. 1973;13(6):630-633.
17. Marquis G, Ramasamy B, Banwarilal S, Munusamy AP. Evaluation of antibacterial activity of plant mediated CaO nanoparticles using *Cissus quadrangularis* extract. *Journal of Photochemistry and Photobiology B: Biology*. 2016;155:28-33.
18. Fu C, Khan MN, Yan J, Hong X, Zhao F, Chen L, Wu H. Mechanisms of nanomaterials for improving plant salt tolerance. *Crop and Environment*; 2023.

19. Khodakovskaya M, Dervishi E, Mahmood M, Xu Y, Li Z, Watanabe F, Biris AS. Carbon nanotubes are able to penetrate plant seed coat and dramatically affect seed germination and plant growth. *ACS nano*. 2009;3(10):3221-3227.
20. Sami F, Yusuf M, Faizan M, Faraz A, Hayat S. Role of sugars under abiotic stress. *Plant Physiology and Biochemistry*. 2016;109:54-61.
21. Mazhar MW, Ishtiaq M, Maqbool M, Atiq Hussain S, Casini R, Abd-ElGawad AM, Elansary HO. Seed Nano-Priming with Calcium Oxide Maintains the Redox State by Boosting the Antioxidant Defense System in Water-Stressed Carom (*Trachyspermum ammi* L.) Plants to Confer Drought Tolerance. *Nanomaterials*. 2023;13(9):1453.
22. Bewley JD, et al. Seeds: Germination, structure, and composition. *Seeds: physiology of development and germination*. 1994;1-33.
23. Aroca R, R Porcel, Ruiz-Lozano JM. Regulation of root water uptake under abiotic stress conditions. *Journal of Experimental Botany*. 2012;63(1):43-57.
24. Mulaudzi T, Hendricks K, Mabiya T, Muthevhuli M, Ajayi RF, Mayedwa N, Iwuoha E. Calcium improves germination and growth of Sorghum bicolor seedlings under salt stress. *Plants*. 2020;9(6):730.
25. Ayesha Farooq, Sumera Javad, Khajista Jabeen, Anis Ali Shah, Asma Ahmad, Adnan Noor Shah, Mohammed Nasser Alyemeni, Walid F.A Mosa, Asad Abbas. Effect of calcium oxide, zinc oxide nanoparticles and their combined treatments on growth and yield attributes of *Solanum lycopersicum* L. *Journal of King Saud University – Science*. 2023;35(5).
26. Alberto García-González, Lluvia de Abril Alexandra Soriano-Melgar, María Luisa Cid-López, Gladis Yakeline Cortez-Mazatán, Esmeralda Mendoza-Mendoza, Luis Alonso ValdezAguilar, René Darío Peralta-Rodríguez, Effects of calcium oxide nanoparticles on vase life of gerbera cut flowers, *Scientia Horticulturae*. 2023;291.
27. Javadi A, Khomari S, Sofalian O. Seed vigor and boron and calcium nutrition influence oilseed rape germinability and seedling growth under salt stress. *Journal of Plant Nutrition*. 2016;39(12):1688-1696.

© 2023 Gopinath et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/105869>