



Performance of Cotton (*Gossypium hirsutum* L.) Varieties under High Density Planting System in Alfisols of Telangana, India

K. Indudhar Reddy ^{a*}, A. V. Ramanjaneyulu ^b, A. Geetha ^c,
G. Seshu ^d and D. Vishnuvardhan Reddy ^{e++}

^a AICRP on WIA, PJTSAU, Hyderabad, India.

^b AICRP on Agroforestry, PJTSAU, Hyderabad, India.

^c College of Agriculture, Rajendranagar, PJTSAU, Hyderabad, India.

^d College of Agriculture, Jagtial, PJTSAU, Hyderabad, India.

^e PJTSAU, Hyderabad, India.

Authors' contributions

This work was carried out in collaboration among all authors. Author KIR designed the study, performed the statistical analysis, wrote the protocol, and wrote the first draft of the manuscript. Authors AVR and AG managed the analyses of the study. Authors GS managed the literature searches and DVR supervised the total work. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i214040

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107839>

Original Research Article

Received: 19/08/2023

Accepted: 26/10/2023

Published: 31/10/2023

ABSTRACT

Aims: Cotton productivity is lower in India than the global average due to inadequate soil moisture during boll formation stage due to cessation of south west monsoon. Cotton is mainly grown under rainfed conditions in marginal and shallow soils, cultivating short duration cotton varieties with high density could improve productivity in the rainfed regions. So, this experiment was conducted to evaluate the three cotton varieties and a hybrid under four high density spacings.

⁺⁺ Retd. Professor;

*Corresponding author: E-mail: indu4reddy@gmail.com;

Study Design: The experiment was laid out in split plot design and replicated thrice.

Place and Duration of Study: The field experiment was conducted for three years during Kharif 2014, 2015 and 2016 at Regional Agricultural Research Station, Palem, Nagarkurnool District of Telangana state, India. The experimental site was located at 16.51703° North latitude and 78.2469° East longitude and an altitude of 478 m above mean sea level.

Methodology: Treatments consisted of 4 spacings (45×10 cm, 60×10 cm, 75×10 cm and 90×10 cm) under main plots and 3 non bt cotton varieties (Suraj, WGCV-48 and ADB-39) and 1 Bt Cotton hybrid Jadoo was included as 4th treatment under subplots during 2015 and 2016. The soil at the site was Alfisol with low organic carbon (0.24%) and N (210 kg ha⁻¹), high in available P (75 kg P₂O₅ ha⁻¹) and available K (455 kg K₂O ha⁻¹). The crop was cultivated under rainfed conditions. Intercultivation and hand weeding was done and kept weed free. Gross plot size of 9 m × 3 m was maintained for each treatment. Nitrogen dose of 40 kg N ha⁻¹, 20 kg P₂O₅ ha⁻¹ and 20 kg K₂O ha⁻¹ was applied. The single plant data was collected on five tagged plants in each plot of each treatment and yield data was collected from the net plot area.

Results: Seed cotton yield was significantly influenced by the spacing and the varieties. The pooled data of 3 years revealed that cotton crop under 90×10 cm spacing (2054 kg ha⁻¹) recorded significantly higher seed cotton yield than the crop under 75×10 cm (1854 kg ha⁻¹), 60×10 cm (1723 kg ha⁻¹) and 45×10 cm (1603 kg ha⁻¹). The pooled data of 3 years revealed that the seed cotton yield in hybrid Jadoo (2189 kg ha⁻¹) was significantly higher than the three varieties. The seed cotton yield in all the three varieties (Suraj, WGCV-48 and ADB-39) was on par with each other during all the years under study.

Conclusion: High density spacing of 90×10 cm, recorded significantly higher seed cotton yield than the other tested spacing. Among the varieties and the hybrid, cotton Jadoo outperformed all the cotton varieties and seed cotton yield of cotton varieties (Suraj, WGCV-48 and ADB-39) was on par with each other.

Keywords: Cotton; crop geometry; HDPS; spacing; varieties.

1. INTRODUCTION

Cotton (*Gossypium hirsutum* L.) is an important commercial crop in India with an area of 13.29 Mha producing 35.25 MBales of cotton during 2020-21 [1]. Due to its high market demand and the remunerative market prices, it is also called as 'White Gold'. India is the world's leading producer of cotton contributing to 30 percent of total world's cotton production [2]. Although yields in India are well below the global average with lint productivity of 451 kg ha⁻¹ during 2020-21. In India, Maharashtra, Telangana and Gujarat are the major states cultivating cotton and contributes to 70 percent of area and 65 percent of production in India during 2020-21 [1].

Cotton is cultivated primarily under rainfed conditions in India. In India, rainfall commences in June and recedes by end of September [3]. Boll formation in long duration cotton varieties and hybrids starts in October and reaches a peak by the end of November. Thus low or inadequate soil moisture especially in shallow soils negatively affects the boll formation and retention resulting in low yields. Hence, in view of low productivity and inadequate soil moisture conditions, cultivating short duration varieties

with high density for improving productivity in the rainfed regions is an option [4]. Hence, this experiment was proposed to evaluate the three cotton varieties under four high density treatments.

2. MATERIALS AND METHODS

A field experiment was conducted for three years during Kharif 2014, 2015 and 2016. at Regional Agricultural Research Station, Palem, Nagarkurnool District of Telangana state--509215 (India). Treatments consisted of 4 spacings (45 × 10 cm, 60 × 10 cm, 75 × 10 cm and 90 × 10 cm) under main plots and 3 non bt cotton varieties (Suraj, WGCV-48 and ADB-39) and 1 Bt Cotton hybrid Jadoo was included as 4th treatment under subplots during 2015 and 2016. The experiment was laid out in split plot design and replicated thrice. The experimental site was located at 16.51703° North latitude and 78.2469° East longitude and an altitude of 478 m above mean sea level. The soil at the site was Alfisol with low organic carbon (0.24%) and N (210 kg ha⁻¹), high in available P (75 kg P₂O₅ ha⁻¹) and available K (455 kg K₂O ha⁻¹). The crop was cultivated under rainfed conditions. Intercultivation with guntaka and hand weeding

was done and kept weed free. Gross plot size of 9 m × 3 m was maintained for each treatment. Nitrogen dose of 40 kg N ha⁻¹, 20 kg P₂O₅ ha⁻¹ and 20 kg K₂O ha⁻¹ was applied. The single plant data was collected on five tagged plants in each plot of each treatment and yield data was collected from the net plot area. The data was analyzed statistically duly following the technique of analysis of variance (ANOVA) as suggested by Gomez and Gomez [5].

3. RESULTS AND DISCUSSION

3.1 Plant Height

Perusal of the data from Table 1 revealed that plant height at harvest was not significantly influenced by the spacings during the three years under study. However, among the varieties the plant height was significantly different. The findings were contradictory with the findings of Ram and Giri [6] and Paslawar et al. [7]. During 2014, the plant height of WGCV-48 (127.5cm) was significantly higher than that of Suraj and ADB-39 varieties. During 2015, the plant height of hybrid Jadoo (121.3 cm) was significantly higher than the varieties and followed by WGV-48 (104.6cm), Suraj (101.5cm) and ADB-39 (86.4cm). Lowest plant height was recorded in ADB-39 variety. During 2016, the plant height of WGCV-48 was significantly higher than that of remaining varieties and also the hybrid Jadoo. WGCV-48 was followed by ADB-39, Jadoo and Suraj. The height of cotton hybrid Jadoo during 2016 was lesser than that of varieties was due to the low available soil moisture during crop period due to less rainfall. This showed that the hybrids are less tolerable to stresses than the varieties. The pooled data of 3 years revealed that WGCV-48 (111.5cm) recorded significantly higher plant height than that of Suraj (98.8cm) and ADB-39 (95.9cm) but on par with that of hybrid Jadoo (106.8cm). All the varieties have performed in a similar trend under tested spacings. Hence the interaction effect was found non-significant.

3.2 Number of Harvested Bolls Per Plant

The number of harvested bolls per plant was significantly influenced by the spacings during all the years under study (Table 1). However, the varieties recorded significantly different number of harvested bolls per plant. During 2014, cotton crop under 90×10 cm spacing recorded significantly higher number of harvested bolls per plant than that of the rest of treatments. During 2015, crop under 90×10 cm spacing recorded

significantly higher number of harvested bolls per plant than that of crop under 45×10 cm but on par with the crop under 60×10 cm and 75×10 cm. During 2016, crop under 90×10 cm spacing recorded significantly higher number of harvested bolls per plant than that of crop under 45×10 cm and 60×10 cm but on par with the crop under and 75×10 cm. The pooled data of 3 years revealed that cotton crop under 90×10 cm spacing recorded significantly higher number of harvested bolls per plant than that of the rest of treatments. Similar results were reported by Pradeep et al. [8] and Narayana and Aparna [9]. The number of bolls per plant in all the three varieties was on par with each other during all the years under study. However, the number of bolls per plant in hybrid Jadoo was significantly higher than the three varieties during 2015 and 2016. The pooled data of 3 years revealed that the number of bolls per plant in hybrid Jadoo (13.50) was significantly higher than the three varieties.

3.3 Boll Weight

Perusal of the data from Table 2 revealed that boll weight was not significantly influenced by the spacings during the three years under study. Similar findings were reported by Moola and Giri [10] and Pradeep et al. [8]. However, among the varieties the boll weight was significantly different. During 2014, among the varieties, Suraj (4.5 g) recorded significantly higher boll weight than ADB-39 (2.9 g) but on par with WGCV-48 (4.2g). During 2015 and 2016, cotton hybrid Jadoo recorded significantly higher boll weight than the three varieties. Among the varieties, Suraj recorded significantly higher boll weight than ADB-39 but on par with WGCV-48. From the pooled data of 3 years, cotton hybrid Jadoo (5.4g) recorded significantly higher boll weight than the three varieties. Among the varieties, Suraj (4.6g) recorded significantly higher boll weight than ADB-39 (3.2g) but on par with WGCV-48 (4.4g).

3.4 Seed Cotton Yield

Seed cotton yield was significantly influenced by the spacings and the varieties. During all the 3 years under study, cotton crop under 90×10 cm spacing recorded significantly higher seed cotton yield than that of under 45×10 cm, 60×10 cm and 75×10 cm. The pooled data of 3 years revealed that cotton crop under 90×10 cm spacing (2054 kg ha⁻¹) recorded significantly higher seed cotton yield than the crop

Table 1. Plant height (cm) and Number of harvested bolls per plant as influenced by different spacings and varieties

	Plant height (cm)				Number of harvested bolls per plant			
	2014	2015	2016	Pooled	2014	2015	2016	Pooled
Spacing								
S1-45 × 10 cm	111.2	96.5	88.3	98.7	7.11	7.23	8.42	7.59
S2-60× 10 cm	112.3	98.3	94.4	101.7	7.66	8.40	9.10	8.39
S3-75 × 10 cm	116.9	93.6	94.2	101.6	8.22	8.13	10.6	8.98
S4-90× 10 cm	121.6	102.2	95.1	106.3	10.8	8.75	11.1	10.22
S.Em±	5.8	6.4	3.0	4.8	0.76	0.21	0.46	0.28
C.D at 5%	NS	NS	NS	NS	2.64	0.64	1.24	0.91
Varieties								
V1-Suraj	111.2	101.5	83.8	98.8	9.00	8.94	9.62	9.28
V2-WGCV-48	127.5	104.6	102.4	111.5	7.75	8.12	9.20	8.66
V3-ADB-39	107.8	86.4	93.4	95.9	8.66	8.33	10.20	9.27
V4-Jadoo	-	121.3	92.3	106.8	-	12.60	14.40	13.50
S.Em±	2.7	2.9	1.9	2.5	0.59	0.73	0.84	0.81
C.D at 5%	8.2	9.6	5.7	7.8	NS	2.78	2.90	2.24

Table 2. Boll weight (g) and Seed cotton yield (kg ha⁻¹) as influenced by different spacings and varieties

	Boll weight (g)				Seed cotton yield (kg ha ⁻¹)			
	2014	2015	2016	Pooled	2014	2015	2016	Pooled
Spacing								
S1-45 × 10 cm	3.7	3.8	4.7	4.0	1492	1515	1802	1603
S2-60× 10 cm	4.2	4.0	4.7	4.3	1578	1648	1944	1723
S3-75 × 10 cm	4.0	3.9	4.7	4.2	1780	1625	2156	1854
S4-90× 10 cm	3.6	4.1	5.0	4.2	1927	1861	2373	2054
S.Em±	0.1	0.2	0.2	0.2	26	28	71.6	38
C.D at 5%	NS	NS	NS	NS	90	88	112	101
Varieties								
V1-Suraj	4.5	4.4	5.0	4.6	1683	1462	2180	1775
V2-WGCV-48	4.2	4.3	4.7	4.4	1745	1533	2207	1828
V3-ADB-39	2.9	2.9	3.8	3.2	1656	1501	2316	1824
V4-Jadoo	-	5.2	5.6	5.4	-	1683	2695	2189
S.Em±	0.1	0.2	0.2	0.2	49	36	85	61
C.D at 5%	0.4	0.5	0.6	0.5	NS	91	251	190

under 75×10 cm (1854 kg ha⁻¹), 60×10 cm (1723 kg ha⁻¹) and 45×10 cm (1603 kg ha⁻¹). The seed cotton yield in all the three varieties (Suraj, WGCV-48 and ADB-39) was on par with each other during all the years under study. However, the seed cotton yield in hybrid Jadoo was significantly higher than the three varieties during 2015 and 2016. The pooled data of 3 years revealed that the seed cotton yield in hybrid Jadoo (2189 kg ha⁻¹) was significantly higher than the three varieties. The seed cotton yield recorded by all the varieties were in a similar trend under tested spacings. Hence the interaction affect was found non-significant.

4. CONCLUSION

The present study on the performance of cotton varieties under high density planting system revealed that among the tested high density spacings, 90×10 cm spacing recorded significantly higher seed cotton yield than the other spacings. Among the varieties and the hybrid, cotton Jadoo outperformed all the cotton varieties (Suraj, WGCV-48 and ADB-39). However, among the varieties all the three varieties recorded seed cotton yield which was on par with each other and there was no significant difference between the varieties.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Anonymous. Agricultural Statistics at a glance- 2022 Government of India Ministry of Agriculture & Farmers Welfare Department of Agriculture & Farmers Welfare Economics & Statistics Division. 2022:59.
2. Anonymous. Economic research service, USDA.2023. <https://www.ers.usda.gov>
3. IMD. Frequently asked on monsoon. 2023. Accessed 26th September, 2023. Available:https://mausam.imd.gov.in/imd_latest/monsoonfaq.pdf
4. Venugopalan MV, Kranthi KR, Blaise D, Lakde, Shubhangi, Shankaranarayanan K. High density planting system in cotton-The Brazil Experience and Indian Initiatives. Cotton Research Journal. 2013;5(2):172-185.
5. Gomez KA, Gomez AA. Statistical Procedures for Agricultural Research. 2nd Edition, John Wiley and Sons, New York. 1984:680.
6. Ram M and Giri AN. Response of newly released cotton (*Gossypium hirsutum*) varieties to plant densities and fertilizer levels. Journal of Cotton Research and Development. 2006;20(1):85-86.
7. Paslawar AN, Deotalu AS, Nemade PW. High density planting of cotton variety akh – 081 under rainfed condition of vidharbha. Plant Archives.2015;15(2):1075-1077.
8. Pradeep Kumar AS, Karle Deshraj Singh, Lalita Verma. Effect of High Density Planting System (HDPS) and Varieties on Yield, Economics and Quality of Desi Cotton. International journal of Current Microbiology and Applied Science. 2017;6(3):233-238.
9. Narayana E, Aparna D. Performance of cotton varieties (*Gossypium arboreum* L.) under different spacings and nitrogen levels in black cotton soils of coastal Andhra Pradesh. Journal of Cotton Research and Development. 2011;25(1):59-62.
10. Moola R, Giri AN. Response of newly released cotton (*Gossypium hirsutum*) varieties to plant densities and fertilizer levels. Journal of Cotton Research and Development.2006;20(1):85-86.

© 2023 Reddy et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107839>