



Effect of Integrated Nutrient Management on Growth, Yield and Quality in Garlic (*Allium sativum* L.)

Bhag Singh ^{a++*} Hem Raj Sharma ^{a#} and Impa H. R. ^{a++}

^a Department of Vegetable Science, Dr YSP UHF, Nauni, Solan (H.P.) 173 230, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i224174

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/105491>

Original Research Article

Received: 29/06/2023

Accepted: 05/09/2023

Published: 05/12/2023

ABSTRACT

Garlic has the charisma of a potent remedy and has held its repute as a therapeutic panacea since the dawn of civilization. So to increase its productivity and medicinal value integrated approach will be required. Allicin is the main bioactive compound, which is responsible for the medicinal value produced by garlic. It is present in the bulb in the form of alliin, which is converted to allicin when the bulb is cut or crushed. The study was conducted on the garlic variety "Kandaghat Selection". The experiment was laid out in Randomized Complete Block Design (RCBD) with three replications comprising fourteen treatments was carried out at the Experimental farm of Horticulture Research and Training Station and KVK, Kandaghat, Dr. YS Parmar University of Horticulture and Forestry, Nauni, Solan (HP) during the Rabi season of 2018-19. Out of the fourteen treatments comprising of 100% recommended dose of NPK + 50 kg S/ha + 5% Jv @ 1 L/ m² (T₇) recorded significantly higher plant height, number of leaves per plant, bulb weight, bulb diameter, number of cloves per bulb, bulb yield, dry matter content of bulb and TSS. Phytomedicine evaluation by HPLC showed that all bulbs from all treatments are rich in allicin content and their recorded values were higher than pharmaceutical grade.

⁺⁺ Research Scholar;

[#] Professor & Head;

^{*}Corresponding author: E-mail: bhagsingh777@gmail.com;

Keywords: INM in garlic; sulphur; allicin; jeevamrit.

1. INTRODUCTION

Garlic (*Allium sativum* L.) is popularly grown *Allium* species after onion belonging to the family Alliaceae. It is originated in Central Asia and mainly used for food as well as medicinal purposes [1]. Garlic is grown for its therapeutic benefits, which are increasing in popularity around the world. It reduces total plasma cholesterol, blood pressure, and platelet aggregation [2]. The majority of garlic's therapeutic properties are due to a sulfur component known as allicin [3]. The minimal allicin content required to assure the medicinal and economic viability of garlic powder products, according to British Pharmacopoeia 1998, is 4.5 mg/g. Globally, it occupies an area of 1,577.77 thousand hectares and a production of 28,164.05 thousand metric tonnes with an average productivity of 17.85 tonnes per hectare (FAO, 2017). In India, it is grown in an area of 319 thousand hectares with a production of 1862 thousand metric tonnes (NHB, 2019). In Himachal Pradesh garlic is mainly grown as a cash crop covering an area of 4.95 thousand hectares and production of 8.49 thousand metric tonnes [4]. Sulphur which is the fourth major plant nutrient after nitrogen, phosphorus and potassium is essential for building up sulphur-containing amino acids (cystine, cysteine and methionine) in plant cells, particularly in the early stage of plant growth [5]. Sulphur is a critical component in excellent garlic production; consequently, a lack of optimal supply in different plant sections inhibits crop growth at any stage, resulting in yield reduction. Recent research has shown that amino acids can influence plant growth and development physiological activities either directly or indirectly. Furthermore, amino acids are widely known as bio-stimulants with good effects on plant growth and yield, as well as greatly mitigating the effects of abiotic stressors. Sulphur is a constituent of the enzyme nitrite reductase which is responsible for the reduction of NO_2^- in chloroplasts and thus reduces the accumulation of cancerous compounds like nitrates in vegetables. Sulphur-deficient plants also displayed inadequate use of macro and micronutrients. Sulphur is an important macronutrient that, at optimal concentrations, stimulates plant development [6]. Many types of living creatures are activated by organic manures, which emit phytohormones that may drive plant growth and nutrient absorption. As a result, organic manures play an important role in

the growth and development of garlic [7]. Organic manure works as a great substrate for soil microbes and raises the fraction of labile carbon and nitrogen, directly driving microorganism population and activity. According to Marathe et al. [8] the presence of organic manures increased the microbial population since there was more organic carbon and mineralized nutrients available for their growth and continued cellular development. Jeevamrut contains a high concentration of beneficial microflora, which helps to maintain and accelerate plant growth, resulting in stronger vegetative growth and a higher quality output [9].

2. MATERIALS AND METHODS

A field experiment was conducted at the Experimental farm of Horticulture Research and Training Station and KVK, Kandaghat, Dr YS Parmar University of Horticulture and Forestry, Nauni, Solan (HP) during the rabi season of 2018-19. This experiment was laid out in Randomized Block Design (RBD) with fourteen treatments and three replications. The size of the experimental plot was 2.0 m × 2.0 m with spacing of 20 cm × 10 cm. The cultivar of garlic used for the present study was 'Kandaghat Selection'. In each treatment, ten plants were randomly selected from each replication for recording morphological and biochemical parameters. Total soluble solids (TSS) was estimated using a hand refractometer. Allicin content of bulbs of different treatments from each replication was determined with HPLC by using the method of British Pharmacopoeia 1998. Baghalian et al. (2005) provide a detailed description of this approach. Treatment details.


2.1 Preparation of Jeevamrit

Jeevamrit drenching at fortnightly intervals and first drenching after 15 days of sowing was given. Table 1 shows the standardized techniques of preparing the Jeevamrit as suggested by Sreenivasa et al. [10].

Table 1. Ingredients of Jeevamrit used for drenching

Ingredient	Quantity
Cow dung	10kg
Cow urine	10L
Jaggry	2 kg
Pulse Flour	2 kg
Fertile Soil	1 kg
Water	200 L

Flow chart for the preparation of Jeevamrut



2.2 Statistical Analysis

The experimental results were statistically analysed by using a general linear model of the standard errors of the mean. The data obtained in Randomized Complete Block Design (RCBD) for each parameter were tested by ANOVA using MS-Excel and OPSTAT. The difference between the treatments was compared by critical difference (CD) at a 5 per cent level of probability (confidence), wherever the results were significant. The calculated F-values were compared with tabulated F-values. When the F-test was significant, CD was then calculated to find out the comparative effectiveness among different treatments.

3. RESULTS AND DISCUSSION

3.1 Growth and Yield Parameters

The plant height, number of leaves per plant, bulb weight, number of cloves per bulb, bulb diameter and bulb yield per hectare of garlic were significantly affected by the different treatments of integrated nutrient management. The plant height of garlic was significantly affected by the application of sulphur and jeevamrit (Table 2). S is the chief constituent of several enzymes and amino acids that are required for chlorophyll synthesis and it increases the uptake of N which is a chief constituent of chlorophyll [11]. S deficiency causes the accumulation of soluble N within the plant but prevents utilization of N [12]. Thus,

increasing the level of S in turn improved plant growth by meeting higher nutritional demand for plant growth. The results also support the findings of Jawagadi *et al.* [13] who reported that integrated nutrient management improved soil fertility by increasing organic carbon, available N, S, Mn and Fe and improvement in nutrient availability resulted in a significant increase in plant height.

The number of leaves in a plant is directly correlated with the leaf area for photosynthetic activity and ultimately with yield. The photosynthates that are formed in the leaves are ultimately stored in the bulb as garlic is an underground crop. So, there is a direct correlation between leaf number, photosynthates manufactured in the leaves and the carbohydrates stored in the bulb. An increase in a number of leaves per plant under integrated nutrient management treatments might be due to the steady release of nutrients throughout the crop growth period. The above findings conform with the findings of Farooqui *et al.* [14] and Anand *et al.* [15] The application of jeevamrit might have increased the microbial population which may have led to improved nutrient availability thereby, resulting in a maximum number of leaves per plant. Findings are also supported by Chatoo *et al.* [16] Increasing the dose of sulphur and jeevamrit application had a positive effect on several leaves as observed in treatment T₇. Similar results were also obtained by Verma *et al.* [11] and Patidar *et al.* [17].

The significant effect on bulb weight as a consequence of integrated nutrient management can be attributed to the increased nutritional status of the soil resulting in increased crop growth. This may be further attributed to the favorable effect of organic sources on microbial activity and root penetration in soil which might have caused a solubilizing effect on native nitrogen, phosphorus, potassium, sulphur and other nutrients. Maximum bulb weight in treatment T₇ (Table 2) might be due to the role of nitrogen on protein synthesis, chlorophyll and enzymatic activity; the role of P on root development, phosphor-lipid and phosphor-proteins formation as well as due to the role of K on the promotion of enzymes activity and enhancing the translocation of assimilates [18]. It might also be due to the role of sulphur in building up of sulphur-containing amino acids in plant cells, particularly in the early stage of plant growth and also it is the fourth major plant nutrient after nitrogen, phosphorus, and

potassium [5]. These results are also confirmed by Magray et al. [6] and Chattoo et al. [19].

Application of sulphur increases the uptake of N which is a chief constituent of chlorophyll. Due to this reason, a significant increase in chlorophyll content and other growth and yield attributes was observed. The similar results also found by Zaman et al. [20] Verma et al. [11] Damse et al. [21] and Shete et al. [22]. Sulphur is essential for building up sulphur-containing amino acids in plant cells, particularly in the early stage of plant growth and also it is the fourth major plant nutrient after nitrogen, phosphorus, and potassium [5]. Amino acids can

either directly or indirectly affect the physiological processes that contribute to plant growth and development, according to studies. Additionally, amino acids are widely known as bio-stimulants that promote plant growth, increase yield, and greatly lessen the harm brought on by abiotic stresses. Anand et al. [15] Patidar et al. [17] and Singh et al. [23] also reported a similar effect of sulphur. Kurubetta et al. (2017) reported that the application of jeevamrit has a significant effect on the yield parameters such as bulb weight, bulb diameter and number of cloves. Similar results of an increase in yield were reported by Manjutha et al. [24].

List 1. Treatment no. with treatment details

Treatment No.	Treatment Details
T ₁	No application of fertilizers and Jeevamrit
T ₂	100% recommended dose of NPK (125:75:60)kg/ha
T ₃	100% recommended dose of NPK + 40kg S/ha
T ₄	100% recommended dose of NPK + 50kg S/ha
T ₅	100% recommended dose of NPK + 60kg S/ha
T ₆	100% recommended dose of NPK + 40kg S/ha + 5% Jv* @ 1 L/m ²
T ₇	100% recommended dose of NPK + 50kg S/ha + 5% Jv @ 1 L/ m ²
T ₈	100% recommended dose of NPK + 60kg S/ha + 5% Jv @ 1 L/ m ²
T ₉	75% recommended dose of NPK + 40kg S/ha
T ₁₀	75% recommended dose of NPK + 50kg S/ha
T ₁₁	75% recommended dose of NPK + 60kg S/ha
T ₁₂	75% recommended dose of NPK + 40kg S/ha + 5% Jv @ 1 L/ m ²
T ₁₃	75% recommended dose of NPK + 50kg S/ha + 5% Jv @ 1 L/ m ²
T ₁₄	75% recommended dose of NPK + 60kg S/ha + 5% Jv @ 1 L/ m ²

*Jv = First jeevamrit drenching after 15 days of sowing and repeated at fortnightly intervals (total 14 applications)
FYM @ 250q/ha applied in all the plots (Except T₁)

Table 2. Effect of different treatments of integrated nutrient management on growth and yield

Tr. No.	Plant height (cm)	No. of leaves/pl ant	Days to harvest	Bulb weight (g)	Bulb diameter (cm)	No. of cloves/b ulb	Peeling index (%)	Bulb yield (q/ha)
T ₁	64.53	7.97	219.67	39.63	3.71	10.90	90.33	153.63
T ₂	72.63	8.73	230.33	51.67	4.53	12.50	93.82	194.33
T ₃	77.54	9.24	231.00	57.29	4.84	13.00	92.97	206.65
T ₄	80.48	9.61	231.67	60.22	5.05	13.22	90.92	213.62
T ₅	79.74	9.56	231.33	59.78	4.95	13.12	89.95	212.23
T ₆	82.65	9.87	234.00	61.57	5.06	13.37	93.65	220.88
T ₇	86.39	10.23	235.00	64.84	5.39	14.17	93.32	233.57
T ₈	86.15	10.15	235.33	64.25	5.28	13.90	93.54	231.66
T ₉	69.94	8.28	226.67	49.82	4.38	12.10	93.38	185.17
T ₁₀	72.55	8.71	228.00	51.37	4.49	12.47	92.26	191.40
T ₁₁	72.09	8.68	228.33	51.21	4.44	12.40	93.80	190.23
T ₁₂	73.83	8.90	229.00	53.53	4.57	12.63	92.94	196.09
T ₁₃	77.36	9.18	230.33	55.59	4.72	12.87	92.17	202.33
T ₁₄	77.29	9.07	230.33	55.22	4.65	12.87	90.43	202.03
CD	2.38	0.52	2.16	1.88	0.19	0.86	NA	9.52

0.05

3.2 Quality Parameters

Farooqui *et al.* [14] reported that the increasing dose of nitrogen increases the dry weight of bulbs and bulb yield up to 150 kg N/ha. The availability of nitrogen is of prime importance for growing plants as it is a major constituent of protein and amino acids. Similar observations have also been recorded by Yadav *et al.* [25] and Banafar and Gupta [26] Neelima *et al.* [27] reported that there was a significant improvement in the growth and yield in tomato plants with the combined application of liquid organic manures such as jeevamrit compared to RDF alone. Patidar *et al.* [17] reported that 50 kg sulphur per hectare with RDF significantly increases the dry matter content of the bulb. This might be due to the role of sulphur in improving amino acids and that the uptake of nutrients directly enhances the dry matter accumulation in the bulb. Similar results were also reported by Damse *et al.* [21] and Anand *et al.* [15].

Oleoresin is an extremely concentrated product containing all the flavouring ingredients soluble in a particular solvent. Oleoresin content decides the quality and market value of the particular variety. Oleoresin content was significantly enhanced with the increasing levels of sulphur (Table 3). These results are in accordance with

Jaggi [28] Banafar and Gupta [26] and Farooqui *et al.* [14] According to Sindhu and Sekhon (2000), the improvement in the quality attributes due to various fertilizer treatments is directly correlated with the physico-chemical and biological properties of soil which enables roots to proliferate more resulting in better uptake and utilization of nutrients required for enhancing the quality of crop. The present findings are consistent with those of Mridula and [29,30].

Singh *et al.* [23] and Chattoo *et al.* [31] also reported similar trends in total soluble solids with the application of sulphur. Sulfur is crucial for the growth and development of plants. It contributes to the production of amino acids like methionine, cysteine, and cystine. It is also responsible for the characteristic taste and smell of garlic like onion and mustard [32]. Jeevamrit application also enhanced the total soluble solids content due to the conversion of organically bound sulphur to the inorganic form [33,34,35].

The Allicin content of the bulb increased significantly with different treatments of integrated nutrient management might be due to sulphur application and an increase in inorganic sulphur in the soil. That ultimately enhances the sulphur-containing compounds.

Table 3. Effect of different treatments of integrated nutrient management on Quality parameters

Tr. No.	wt of unpeeled 100 cloves (g)	Dry matter content (%)	Oleoresin content (%)	TSS (°Brix)	Sulphur content (%)	Allicin (mg/g)
T ₁	309.33	36.53	0.74	32.13	1.02	4.73
T ₂	373.33	39.23	1.08	33.77	1.11	5.28
T ₃	400.00	41.00	1.30	34.80	1.31	5.76
T ₄	418.33	42.25	1.42	35.30	1.42	5.94
T ₅	414.00	42.23	1.39	35.27	1.42	5.91
T ₆	421.67	42.98	1.42	35.47	1.38	5.95
T ₇	445.67	44.15	1.60	36.30	1.50	6.19
T ₈	443.33	44.05	1.61	35.97	1.52	6.15
T ₉	354.67	38.70	0.99	33.20	1.17	5.00
T ₁₀	370.33	39.08	1.10	33.60	1.26	5.15
T ₁₁	366.00	39.01	1.09	33.53	1.28	5.12
T ₁₂	370.00	40.44	1.14	33.83	1.23	5.23
T ₁₃	384.33	40.95	1.28	34.13	1.34	5.36
T ₁₄	382.33	40.91	1.25	34.17	1.35	5.33
CD ₀	17.60	0.83	0.14	0.63	0.06	0.23

05

4. CONCLUSION

It can be concluded that among different treatments of integrated nutrient management the treatment comprising of 100% recommended dose of NPK + 50 kg S/ha + 5% Jv @ 1 L/ m² (T₇) performed best for most of the parameters under study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Diriba S, Nigussie D, Kebede W, Getachew T, Sharma JJ. Growth and nutrients content and uptake of garlic (*Allium sativum* L.) as influenced by different types of fertilizers and soils. *African Journal of Agricultural Research*. 2013;8:5387-5390.
2. Sterling SJ, Eagling R. Agronomic and allicin yield of Australian grown garlic (*Allium sativum*). *Acta Hort*. 2001;555:63 - 73.
3. Schulz V. Garlic. R. Hansel, V.E. Tayler (Eds.), *Rational Phytotherapy. A Physicians' Guide to Herbal Medicine* (3rd ed.), Springer. 1998;107-125.
4. Anonymous. 2018. National Horticulture Research and Development Foundation. <http://nhrdf.org> [February 18th, 2020].
5. Havlin JL, Beaton JD, Tisdale SL, Nelson WL. *Soil fertility and fertilizers: An introduction to nutrient management*. 7th edn. Person Education Inc. Singapore. 2004;221.
6. Magray MM, Chattoo MA, Narayan S, Mir SA. Influence of Sulphur and Potassium applications on yield, uptake & economics of production of garlic. *Int. J. Pure App. Biosci*. 2017;5:924-934.
7. Yoldas F, Ceylan S, Mordogan N, Esetlili BC. Effect of organic and inorganic fertilizers on yield and mineral content of onion. *Afr. J. Biotechnology*. 2011;10:11488-11492.
8. Marathe RA, Bharambe PR, Sharma Rajvir, Sharma UC. Leaf nutrient composition, its correlation with yield and economics of sunflower. *Karnataka Journal of Agriculture Sciences*. 2012;22: 198-199.
9. Devakumar N, Shubha SB, Gouder, Rao GGE. Microbial analytical studies of traditional organic preparations beejamrutha and jeevamrit. *Proceedings of the 4th ISOFAR Scientific Conference*. 2014;639-642.
10. Srinivasa MN, Naik N, Bhat SN, Nekar MM. Effect of organic liquid manures on growth, yield and quality of chilli. *Green Farming*. 2010;01:282-284.
11. Verma S, Choudhary MR, Yadav BL, Jakhar ML. Influence of vermicompost and sulphur on growth and yield of garlic (*Allium sativum* L.) under semi arid climate. *Journal of Spices and Aromatic Crops*. 2013;22:20–23.
12. Charities N, Carpentiers LJ. The role of sulphur in biology and its importance in Agriculture. *Soil Sci*. 1956;10:267–292.
13. Jawadagi RS, Basavaraj N, Hemla NB, Patil BN, Channappagoudar BB. Effect of planting geometry and organic sources of nutrients on growth, yield and quality of rabi onion cv. Bellary Red. *Karnataka Journal of Agricultural Sciences*. 2012;25:236-240.
14. Farooqui MA, Naruka IS, Rathore SS, Singh PP, Shaktawa RPS. Effect of nitrogen and sulphur levels on growth and yield of garlic. *Asian Journal of Food Agriculture and Industry*. 2009;5:18-23.
15. Anand M, Sankari A, Anita B. Influence of integrated nutrient management for garlic under nilgiris condition. *Journal of Current Microbiology and Applied Sciences*. 2017;6:3833-3838.
16. Chattoo MA, Ahmed N, Faheema S, Narayan S, Khan SH, Hussain K. Response of garlic to biofertilizer application. *The Asian Journal of Horticulture*. 2007;2:249-252.
17. Patidar M, Shaktawat RPS, Naruka IS. Effect of sulphur and vermicompost on growth, yield and quality of garlic. *Journal of Krishi Vigyan*. 2017;5:54-56.
18. El-Desuki M, Mahmoud AR, Hafiz MM. Response of onion plants to minerals and bio-fertilizers application. *Research Journal of Agriculture and Biological Science*. 2006;2:292-298.
19. Chattoo MA, Magray MM, Malik AA, Shah MD, Chisti JA. Effect of sources and levels of sulphur on growth, yield and quality of onion. *International Journal of Current Microbiology and Applied Sciences*. 2019;8:1462-1470.

20. Zaman MS, Hashem MA, Jahiruddin M, Rahim MA.. Effect of sulphur fertilization on the growth and yield of garlic. *Bangladesh Journal of Agriculture Research*. 2011;36:647-656.
21. Damse DN, Bhalekar MN, Pawar PK. Effect of integrated nutrient management on growth and yield of garlic. *The Bioscan*. 2014;9:1557-15560.
22. Shete MB, Chiktey HM, Jadhav SB, Bhalekar MN. Effect of sulphur on growth, yield and quality of garlic. *International Journal of Chemicals Studies*. 2018;6:552-555.
23. Singh CV, Gupta P, Kasana BS. Response of garlic to sulphur and boron application in terms of biochemical parameters. *International Journal of Current Microbiology and Applied Sciences*. 2018;7:2677-2687.
24. Manjutha GS, Upperi SN, Pujari BT, Yeledahalli, Kuligod VB. Effect of farm yard manure treated with jeevamrit on yield attributes, yield and economics of sunflower. *Karnataka Journal of Agriculture Sciences*. 2009;22: 198-199.
25. Yadav RL, Sen NL, Yadav BL. Response of onion to nitrogen and potassium fertilization under semi-arid condition. *Indian Journal of Horticulture*. 2003;60:176-178.
26. Banafar RNS, Gupta NK. Effect of fertilizer mixture with and without sulphur on quality of onion. In: National Seminar on Agro-technology, Quality, Processing and Export of Spices J. N. K. V. V. Jabalpur. 2005;60.
27. Nileema S, Gore, Sreenivasa MN. Influence of liquid organic manures on growth, nutrient content and yield of tomato in the sterilized soil. *Karnataka Journal of Agriculture Science*. 2011;24:153- 157.
28. Jaggi RC. Effect of sulphur levels and sources on composition and yield of onion. *International Journal of Agriculture Sciences*. 2004;74:219-220.
29. Mridula KR, Jayachandran BK. Quality of mango ginger as influenced by mineral nutrition. *Journal of Tropical Agriculture*. 2001;39:182-183.
30. Velmurugan M, Chezhiyan N, Jawaharlal M. Influence of organic manures and inorganic fertilizers on cured rhizome yield and quality of turmeric. *International Journal of Agriculture and Science*. 2008;1:142-145.
31. Chattoo MA, Magray MM, Parray FAH, Shah MD, Bhat TA. Effect of sulphur on growth, yield and quality of garlic. *Journal of Pharmacognosy and Phytochemistry*. 2018;7:2894-2896.
32. Tisdale SL, Nelson WL, Beaton JD. *Soil fertility and fertilizers, 4th edition*. Macmillan Publication Company, New York; 1985.
33. Anonymous. 2017. Food and agriculture organization of United Nations. FAOSTAT. Available:<http://www.fao.org/faostat/en/#data/QC>. [February 12th, 2019]
34. Anonymous. 2019. nhb.gov.in/area-pro/1st_Advance_Estimates_2018-19.xls.
35. Chesnin L, Yien CH. Turbidimetric determination of available sulphates. *Soil Sciences Society of America Proceeding*. 1950;15:149-151.

© 2023 Singh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/105491>