



# Review of Traditional Plants from Toliara Province in Southern Madagascar for the Treatment of Diabetes and Cancer

Jean Hugues Razafimahatratra <sup>a,b,c,\*</sup>,  
Carène Hanitriniaina <sup>a,b</sup>, Silvère Soavina <sup>a</sup>,  
Solofoniaina Razafimahefa <sup>d</sup>, Vincent Lequart <sup>e</sup>,  
Christian Manjvelo <sup>a,c</sup>, Patrick Martin <sup>e\*</sup>  
and Ruphin Fatiany <sup>a</sup>

<sup>a</sup> Département de Chimie, Université de Toliara, Madagascar.

<sup>b</sup> Institut d'Enseignement Supérieur du Menabe, Université de Toliara, Madagascar.

<sup>c</sup> Institut d'Enseignement Supérieur de Toliara, Université de Toliara, Madagascar.

<sup>d</sup> Institut d'Enseignement Supérieur Antsirabe Vakinankaratra, Université d'Antananarivo, Madagascar.

<sup>e</sup> Univ Artois, UniLaSalle, Unité Transformations & Agroressources – ULR7519, 62408 Béthune, France.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/EJMP/2023/v34i11-121171

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/109052>

**Short Communication**

**Received: 03/09/2023**

**Accepted: 08/11/2023**

**Published: 30/12/2023**

\*Corresponding author: E-mail: [jeanhuguesberoroha@yahoo.fr](mailto:jeanhuguesberoroha@yahoo.fr);

## ABSTRACT

In all countries, plants have always been and still are used for their health benefits as part of traditional medicine. The same is true of Madagascar, which also boasts considerable biodiversity, including plants that are totally endemic to the country. Our study aims to identify medicinal plants traditionally used to treat diabetes and cancer in the province of Toliara in Madagascar. At the end of the survey we carried out with the local population, using pre-established questionnaires, 34 plants were identified. Thus, among the species recorded, *Myrothamnus moschatus*, *Zanthoxylum decaryi* and *Calotropis procera* were targeted. These are multi-use species, they are used respectively to treat convulsion, hypertension and healing. Among the three species studied, two are endemic to Madagascar (*Myrothamnus moschatus* and *Zanthoxylum decaryi*), however the third is introduced. This study has enabled us to demonstrate the importance of ethnobotanical investigation in the field of research. The identification of these three species provides us with a basis for phytochemical development.

**Keywords:** medicinal plants; *Myrothamnus moschatus*; *Zanthoxylum decaryi*; *Calotropis procera*; ethnobotany.

## 1. INTRODUCTION

Many of the bioactive molecules currently in use are derived from phytochemical studies. Indeed, nature provides a number of remedies for various pathologies. Today, many plant- and animal-derived medicines are widely prescribed, such as madeglucyl anti-diabetic [1] and taxol anti-cancer [2] drugs, and the search for new active ingredients of natural origin remains a hot topic.

According to a report by the World Health Organization (WHO), despite the development of conventional medicine, the majority of people in developing countries like Madagascar still rely on traditional healers and medicinal plants for their care [3,4]. The inaccessibility and high cost of conventional medicine, as well as the effectiveness of medicinal plants, are contributing to an increase in the number of people practicing traditional medicine [5-7]. This increase also testifies to the preoccupation and importance of traditional medicine. The aim of this study is to inventory and identify medicinal plants used in the treatment of diabetes and cancer, with a view to creating a Traditionally Improved Medicine (TIM).

## 2. MATERIALS METHODS

### 2.1 Presentation of the Study Area

The province of Toliara is one of the largest provinces of Madagascar (Fig. 1) with an area of 161,405 km<sup>2</sup> with an estimated population of 4,199,643 inhabitants including 2,054,286 men and 2,145,357 women [8]. This province is distinguished from other provinces of Madagascar by its semi-arid climate,

predominated by the alternation of two seasons namely the dry season which lasts up to nine (09) months followed by a brief rainy season characterized by low annual rainfall less than 600mm. [9].

### 2.2 Ethno-botanical Survey

Ethno-botanical surveys, based on direct questioning of anti-diabetic and anti-cancer medicinal plants, were carried out from June to July 2021 in the province of Toliara among healers, herbalists and, above all, villagers. The survey was carried out using a pre-established questionnaire form. The questions were asked in the local dialect, and varied according to the region where the survey was carried out. They covered socio-demographic data (age, sex, marital status, level of education), plants and organs used, as well as preparation and administration methods.

### 2.3 Sampling

According to the 2004 administrative division, the province of Toliara is made up of four regions. The sample was drawn up using a random sampling method, divided into two groups: group 1 (South West region, Menabe region), group 2 (Androy, Anosy). In order to ensure the objectivity of the data, 3-year students from the Chemistry Department, the Institut d'Enseignement du Menabe and the Institut d'Enseignement Supérieur de Toliara were mobilized for practical and personal student work. They were divided into eight groups of five students. In each group, the presence of at least one student from one of the four regions is required.



Fig. 1. Study area

## 2.4 Species selection

Two basic criteria were adopted in selecting the anti-diabetic and anti-cancer medicinal plants studied.

- The first criterion is based on the medicinal plants most frequently used in the treatment of the two diseases (i.e. the frequency of use to treat diabetes and cancer).
- The second is based on bioavailability in the region surveyed.

## 2.5 Data Analysis

The citation frequency (Fc) of each plant surveyed is calculated using the formula in [10].

$$Fc = \frac{n_c}{N} \times 100$$

With;

$n_c$  = number of citations for the plant in question  
 $N$  = number of people surveyed

## 3. RESULTS AND DISCUSSION

### 3.1 Sociodemographic Characteristics of the Surveys (Table 1)


For this study, 108 people were surveyed, including 64.81% healers, 7.41% herbalists and 27.78% users. The majority of those surveyed were men (68.96%). Age ranged from under 30 to over 60, with the majority over 60 (50%). The majority of respondents came from the Androy region (49%) and the South-West region (38%). The highest percentages of respondents were healers (64.81%). With regard to educational level, the results show that 62.96% are illiterate, while university level is less represented (1.85%). In terms of marital status, the majority of respondents are married (72.22%).

### 3.2 Plant Parts Used

The most commonly used plant parts are leaves (20.59%), leafy stems (20.59%), stems (2.94%), stem bark (29.41%) and root bark (26.47%). The results show that all four organ parts are similarly used in the preparation of anti-cancer and anti-diabetic recipes in the province of Toliara (Fig. 2).

**Table 1. Socio-demographic characteristics of respondents (N=108).**

Features	Frequency (%)
<b>Gender</b>	
Men	68.96
Women	37.04
<b>Regional origin</b>	
Anosy	4
Androy	49
Menabe	17
South-West	38
<b>Age</b>	
< 30	14.81
30 ≤ x ≤ 60	35.19
x > 60	50.00
<b>Class</b>	
Healers	64.81
Herbalists	7.41
Users	27.78
<b>Level of education</b>	
No schooling	62.96
Primary	25.93
Secondary	9.26
University	1.85
<b>Marital status</b>	
Single	2.78
Divorced	10.19
Married	72.22
Widowed	14.81



**Fig. 2. Histograms of the different parts used**

### 3.3 Preparation Techniques

According to the ethno-botanical survey, two techniques are used: decoction 67.65%, maceration 32.35% (Fig. 3).

### 3.4 Route

Non-toxic plants are taken orally, while toxic plants are taken externally. A total of 34 anti-

diabetic and anti-cancer plant species have been identified in Toliara Province (Table 2).

### 3.5 Plant Selection Criteria

Based on the predefined criteria above, three medicinal plants were selected. These are: *Calotropis procera* (37.04%), *Myrothamnus moschatus* (31.48%), *Zanthoxylum decaryi* (25.00%).

**Table 2. Antidiabetic and anticancer medicinal plants in Toliara Province.**

N°	Family	Scientific name	Vernacular name	Parts used	Preparation technique	Disease	Frequency of quotation (%)
1	Amaranthaceae	<i>Amaranthus spinosus</i>	<i>Tsirapahy</i>	Leafy stem	Decoction	Antidiabetic	1.85
2	Anacardiaceae	<i>Anacardium occidentale</i>	<i>Mahabibo</i>	Stem bark	Decoction	Anticancer	0.93
3	Anacardiaceae	<i>Sclerocarya birrea</i>	<i>Sakoa</i>	Leaf	Decoction	Antidiabetic	2.78
4	Apocynaceae	<i>Calotropis procera</i>	<i>tapoky</i>	Leaf	Decoction	Anticancer/ Antidiabetic	37.04
5	Apocynaceae	<i>Roupellina boivini M.Pichon</i>	<i>Lalondo</i>	Leaf, stem bark, root bark	Maceration	Anticancer	1.85
6	Aristolochiaceae	<i>Aristolochia albida Duch.</i>	<i>Totonga</i>	Leaf root bark	Decoction	Anticancer	0.93
7	Asclépiadaceae	<i>Cryptostegia madagascariensis Boj.</i>	<i>Lombiry</i>	root bark	Maceration	Anticancer	0.93
8	Astéraceae	<i>Dicoma carbonaria Baker</i>	<i>Peha</i>	Leaf	Maceration	Anticancer	1.85
9	Bignoniaceae	<i>Rhigozum madagascariense Drake</i>	<i>Hazontà</i>	Leaf, Stem bark	Decoction	Anticancer	2.78
10	Bignoniaceae	<i>Phyllarthron bernierianum Seem.</i>	<i>Toirave</i>	Stem bark	Maceration	Anticancer	4.63
11	Borraginaceae	<i>Ehretia sp.</i>	<i>Lamoty</i>	root bark	Decoction	Anticancer	7.41
12	Cannabaceae	<i>Trema orientalis</i>	<i>Tsivakimbaratra</i>	Stem bark	Decoction	Antidiabetic	9.26
13	Capparidaceae	<i>Boscia madagascariensis DC.</i>	<i>Pake</i>	Stem bark	Maceration	Anticancer	5.56
14	Ebénaceae	<i>Diospyros sp</i>	<i>vandamena</i>	root bark	Maceration	Anticancer	8.33
15	Ebénaceae	<i>Diospyros sp.</i>	<i>Hazomby</i>	Stem bark	Decoction	Anticancer	11.11
16	Euphorbiaceae	<i>Gelonium boivinianum Mull</i>	<i>Hazombalala</i>	Leafy stem	Decoction	Anticancer	7.41
17	Fabaceae	<i>Chadsia grevei Drake</i>	<i>Remote</i>	root bark	Maceration	Anticancer	5.56
18	Fabaceae	<i>Indigofera tinctoria L</i>	<i>Netse</i>	Leaf	Decoction	Anticancer	6.48
19	Hypéricaceae	<i>Psorospermum sp</i>	<i>Hazompanihy</i>	Stem bark	Maceration	Anticancer	8.33
20	Labiatae	<i>Capitanopsis angustifolia</i>	<i>Reringitse</i>	Leafy stem	Decoction	Antidiabetic	9.26
21	Lamiaceae	<i>Portulaca oleracea</i>	<i>Kalabotretreka</i>	Leafy stem	Decoction	Antidiabetic	12.96
22	Lauraceae	<i>Ranvesera aromatica gmelin</i>	<i>Ravintsara</i>	Leafy stem	Decoction	Antidiabetic	1.85
23	Loganiaceae	<i>Strychnos sp</i>	<i>Mandakolahy</i>	root bark	Maceration	Anticancer	0.93
24	Loganiaceae	<i>Strychnos sp</i>	<i>Hazomby</i>	root bark	Decoction	Anticancer	0.93
25	Meliaceae	<i>Cedrelopsis grevei</i>	<i>Katrafay, katsafa, katrafa</i>	Stem bark	Maceration	Anticancer	1.85
26	Musaceae	<i>Musa paradisiaca</i>	<i>Kida</i>	Ripe banana peel	Decoction	Antidiabetic	4.63
27	Myrothamnaceae	<i>Myrothamnus moschatus</i>	<i>Maharoaky, Somorombato</i>	Leafy stem	Decoction	Anticancer/ Antidiabetic	31.48
28	Myrtaceae	<i>Psidium guajava</i>	<i>Goavy</i>	Leaf and fruit	Decoction	Antidiabetic	8.33
29	Nymphaeaceae	<i>Nymphaea stellata willd</i>	<i>Bekapaiké</i>	Leafy stem	Decoction	Antidiabetic	3.70
30	Olivaceae	<i>Olivax sp.</i>	<i>Ambihotse</i>	Stem	Decoction	Anticancer	7.41
31	Papilionaceae	<i>Cajanus indicus spreng</i>	<i>Ambatry</i>	Root bark	Decoction	Antidiabetic	2.78
32	Rhamnaceae	<i>Ziziphus mauritiana</i>	<i>Tsinefo</i>	Leaf	Decoction	Antidiabetic	0.93
33	Rubiaceae	<i>Paederia majungensis</i>	<i>Lengobe</i>	Dry root	Decoction	Antidiabetic	4.63
34	Rutaceae	<i>Zanthoxylum decaryi</i>	<i>Monongo</i>	Stem bark	Decoction	Antidiabetic/ Anticancer	<b>25.00</b>




Fig. 3. Histograms of preparation techniques

### 3.6 Monograph for Each Plant Selected According to pre-established Criteria

#### a) *Myrothamnus moschatus* (Myrothamnaceae) [11].

It's a dioecious shrub that only grows on bare, sunny rocks, hence the name "lithophyte plant". The flowering season is divided into two main parts: for the male, January to March and July to December; for the female, January to February and October to December. This plant is characterized by its ability to adapt to long periods of drought, and as soon as it rains, the leaves change color. This ability gives it the name "resurrection plant".

The plant is known by several vernacular names, which vary according to ethnic group:

- Maimbelona: mérina
- Fanalahy : betsileo
- Maharoaka et somoro: bara
- Tsangatsanga : tsimihety
- Maharoaky: antandroy

Geographical distribution: Southern part of Madagascar.

Traditional uses:

- Asthma: dried leaves are smoked like tobacco
- Cough: decoction of the whole plant
- External use: to heal wounds and facilitate wound healing

Pharmacological action tested from essential oil [12].

- Antimicrobial activity
- *In vitro*: cytotoxic

Use of essential oil in agriculture [13].

- Effective against crop arthropods and of medical interest

Use as phytomedicines [11]

- Essential oil: to prevent and treat convulsions and migraines

#### b) *Zanthoxylum decaryi*, Rutaceae [14]

A small tree, 5 to 8 cm high, very thorny when young, becoming almost ineradicable with age, with thick, scarred final branches. The hairless leaves measure from 15 to 25 cm and are grouped in a maximum of 12 terminal clusters. Male flowers are pubescent thyrses, tetramerous with stamens leading the carpels astray. Female inflorescences are clustered and very similar to the male flowers.

The plant is known by several Malagasy vernacular names, including monongo, monongovola and monongovola.

Geographical distribution: Southern part of Madagascar

Traditional uses: Antidiabetic, hypertension, malaria: bark decoction

#### c) *Calotropis procera*, Apocynaceae

A 5-metre-high latex shrub with large, glaucous leaves. Its purplish flowers are grouped together. Its fruits resemble small oval apples in the form of testicles.

The plant is known by several Malagasy vernacular names: tapoky, tsipopoky.

Geographical distribution: Southern Madagascar, Eastern India, Arabia, Africa.

Traditional uses:

- Decoction for external use to disinfect wounds (leaves)
- Latex accelerates healing
- Antidiabetic: decoction of leaves (take care to respect the dose)

Pharmacological action tested:

- Effects on biochemical markers of organic functions in OFA rats [15].
- Phytochemical properties and antioxidant activity of *Calotropis procera* (Ait.) R. Br. [16].

- Coagulant properties of *Calotropis procera* and its potential use in the food industry [17].
- Acute toxicological properties of *Calotropis procera* latex in rats [18].
- Coagulant properties of *Calotropis procera* and its potential use in the food industry [19].
- *In vitro* cytotoxicity against different human cancer cell lines of laticifer proteins of *Calotropis procera* (Ait.) R. Br [20].
- *Calotropis procera* (Aiton) Dryand (Apocynaceae) as an anti-cancer agent against canine mammary tumor and osteosarcoma cells [21].



Dry season plant



Wet season plant




Phytomedicine

Fig. 4. *Myrothamnus moschatus*



Fig. 5. *Zanthoxylum decaryi*



**Fig. 6. *Calotropis procera***

### 3.7 Discussion

The ethno-botanical surveys carried out in the province of Toliara questioned 108 people of different classes, the majority of whom were men (68.96%). When age is taken into account, the result is that they are over 60 (50%), which explains why the practice of traditional medicine is the prerogative of older men. These results are in line with the work carried out by Gbekley et al. [22], which also revealed the importance of ancestral knowledge. It emerges from this study that the majority of people surveyed are married (72.22%), illiterate (62.96%) and these results are corroborated with national data (Madagascar-INSTAT-RGPH3, 2018). The present study strictly concerns plants used against cancer and diabetes, among the species listed: *Calotropis procera*, *Myrothamnus moschatus* and *Zanthoxylum decaryi* particularly caught our attention due to the fact that they are the most solicited in the province of Toliara whose citation frequencies are 37.04%; 31.48% and 25% respectively. Moreover, the frequency of use testifies to the abundance of chemical groups that could have effects on people suffering from diabetes and/or cancer, but they

need to undergo scientific validation to justify their use. In traditional medicine, most recipes are prepared by decoction. This explains why the action of decocting recipes.

### 4. PROSPECTS AND CONCLUSION

The ethnobotanical survey carried out in the province of Toliara identified 34 medicinal plants used in the treatment of diabetes and cancer. From this list, *Calotropis procera*, *Myrothamnus moschatus* and *Zanthoxylum decaryi* are the most cited. To scientifically prove the therapeutic efficacy and safety of these three plants, a study is currently underway in partnership with the Institut Malgache de Recherches Appliquées (IMRA) and the Unité Transformations & Agroressources of the Université d'Artois in France. The use of these three medicinal plants to treat diabetes and cancer is another step in the valorization of these plants. It also enables us to contribute further knowledge on the progress of our research in terms of biological activity and the chemistry of natural products.

### CONSENT AND ETHICAL APPROVAL

It is not applicable.



## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Puri M, Masum H, Heys J, Singer PA. Harnessing biodiversity: The Malagasy Institute of Applied Research (IMRA) BMC International Health and Human Rights. 2010;10 (Supplement 1):p. S9. DOI:10.1186/1472-698x-10-s1-s9
- Weaver BA. How Taxol/paclitaxel kills cancer cells. Mol Biol Cell. 2014 Sep 15;25(18):2677-81. DOI:10.1091/mbc.E14-04-0916.
- Chominot A. Valorization of medicinal plants by the pharmaceutical industry; 2000.
- Xiaorui Z. Regulation of herbal medicines. The situation in the world. 1998;65.
- Fattahi FS, Zamani T. Poly (lactic acid) nanoparticles: A promising hope to overcome the cancers. Journal of Advanced Biomedical Sciences. 2021; 11(2):3791-3814. Available:https://doi.org/10.18502/jabs.v11i2.8766.
- Philippe Rasoanaivo, Ermenegilde Ralaibia, Filippo Maggi, Fabrizio Papa, Sauro Vittori, Marcello Nicoletti. Phytochemical investigation of the essential oil from the resurrection plant *Myrothamnus moschatus* endemic to Madagascar.pdf. 201224(3), 299-304.
- Salhi S, Fadli M, Zidane L, Douira A. Floristic and ethnobotanical studies of medicinal plants in the city of Kenitra (Morocco). Lazaroa. 2010;31:133-143.
- Madagascar-INSTAT-RGPH3. National Institute of Statistics; 2018.
- Monograph Atsimo Andrefana Region. Center for research, studies and support for economic analysis in Madagascar (Cream). 2013;208.
- Gbekley E, Agbodeka K, Karou SD, Anani K, Yao A. Bioactive compounds isolated from plants with anti-diabetic property: Review of literature/Isolated bioactive plant compounds with anti-diabetic property: Review. International Journal of Innovation and Applied Studies. 2017;19:839-849.
- Millogo-Rasolodimby. Myrothamnaceae, Family 94. In Flore de Madagascar et des Comoros. National Museum of Natural History. 1991;81-85.
- Rasoanaivo P, Ralaibia E, Maggi F, Papa F, Vittori S, Nicoletti M. Phytochemical investigation of the essential oil from the 'resurrection plant' *Myrothamnus moschatus* (Baillon) Niedenzu endemic to Madagascar. Journal of Essential Oil Research. 2012;24(3):299-304.
- Pavela R, Ferrati M, Spinozzi E, Maggi F, Petrelli R, Rakotosaona R, Ricciardi R, Benelli G. The Essential Oil from the Resurrection Plant *Myrothamnus moschatus* Is Effective against Arthropods of Agricultural and Medical Interest. Pharmaceuticals. 2022;15(12): 1511.
- Perrier de la Bathie H. Rutaceae, Family 104. In Flore de Madagascar et des Comoros. National Museum of Natural History. 1950;1-88
- Lohoues EE, Camara Cissé M, Tiahou Gnomlesson G, Monde A, Sess Essiagne D. Calotropis procera latex: Effects on biochemical markers of organic functions in the ofa rat. Phar11l. Med. Trans. Afr. 2006;14:81-95.
- Kumar A, Dandapat S, Kumar M, Sinha. Phytochemical properties and antioxidant activity of calotropis procera (Ait.) R. Br. Proceedings of International Conference on Harmony with Nature in Context of Ecotechnological Intervention and Climate Change. 2013;195-199.
- Baba-Moussa Farid, Baba-Moussa Lamine, Ahissou Hyacinthe, Bokossa Innocent, Capo-Chichi Bénédicte, Toukourou Fatiou, Sanni Ambaliou. Coagulant properties of *Calotropis procera* and its possibilities of use in the food industry. Sciences and Medicine. 2007; 05:7-12.
- Lohoues EE, Tiarou Gnomblesson G, Monde A, Djinh J, Sess Essiagne D. Acute Toxicological Properties of Calotropis Procera Latex in Rats. Pharm. Med. Trans. Afr. 2006;14:187-199.
- Farid BM, Lamine BM, Hyacinthe A, Innocent B, Bénédicte CC, Fatiou T, Ambaliou S. Coagulant properties of *Calotropis procera* and its possibilities of use in the food industry. 2007;5.
- Soares de Oliveira J, Pereira Bezerra D, Teixeira de Freitas CD, Delano Barreto Marinho Filho J, Odorico de Moraes M, Pessoa C, Costa-Lotufo LV, Ramos MV. *In vitro* cytotoxicity against different human cancer cell lines of

- laticifer proteins of *Calotropis procera* (Ait.). *Toxicology in Vitro*. 2007;21(8):1563-1573.
21. Rabelo AC, Borghesi J, Carreira ACO, Hayashi RG, Bessa F, Barreto R, da SN, da Costa RP, Cantanhede Filho AJ, Carneiro FJC, Miglino MA. *Calotropis procera* (Aiton) Dryand (*Apocynaceae*) as an anti-cancer agent against canine mammary tumor and osteosarcoma cells. *Research in Veterinary Science*. 2021; 138:79-89.
22. Gbekley E, Simplicite KD, Charlemagne G, Kodjovi A, Kokou A, Tchadjobo T, Amegnona A, Komlan B, Jacques S. Ethnobotanical study of plants used in the treatment of diabetes in traditional medicine in the Maritime region of Togo. *Pan African Medical Journal*. 2015;20:437-452.

---

© 2023 Razafimahatratra et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/109052>