



Pilot Study; Estimating Sample Size for the Prevalence of Pre-Clinical Left Ventricular Diastolic Dysfunction Using Two American College of Echocardiography Guidelines in a Semi-Urban Nigerian Population

**Udora Nneka Clara ^{a*}, Ajulufu Ifeyinwa Jennifer ^b,
Ossai Edmund Ndudi ^c and Ejim Emmanuel Chukwudi ^{a,d}**

^a *Department of Medicine, Cardiology Unit, University of Nigeria Teaching Hospital Ituku/Ozalla, Enugu, Nigeria.*

^b *General Medicine, William Harvey Hospital-East Kent Hospitals, Foundation Trust, Ashford, Kent, UK.*

^c *Department of Community Medicine, College of Health Sciences, Ebonyi State University Abakiliki, Nigeria.*

^d *Department of Medicine, University of Nigeria, Ituku/Ozalla Campus, Enugu, Nigeria.*

Authors' contributions

This work was carried out in collaboration among all authors. Author AIJ managed the literature searches and data collection process. Author OEN managed the statistical analyses of the study. Author EEC developed the concept and design of the study. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMMR/2024/v36i55422

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/115616>

Original Research Article

Received: 02/02/2024

Accepted: 05/04/2024

Published: 09/04/2024

*Corresponding author: E-mail: nnekaudora@gmail.com;

ABSTRACT

Aim: To estimate the sample size for the prevalence of pre-clinical left ventricular diastolic dysfunction using two American college of echocardiography guidelines in a semi-urban Nigerian population.

Study Design: Cross sectional, population based.

Place and Duration of Study: Neni, Anambra State, Nigeria on the 18th and 19th December 2023.

Methodology: Fifty subjects with age range of 18–70 years were consecutively recruited after a community awareness creation. Subjects gave informed oral and/or written consent before proceeding for an interviewer administered questioning of their biodata and clinical history. Anthropometric measurements were obtained followed by clinical examination and electrocardiography. Echocardiography was done using the American Society of Echocardiography (ASE) guideline with the left ventricular diastolic function assessed using the 2009 and 2016 guidelines.

Results: Mean age of participants was 54.3 ± 14.0 years consisting of 15 males and 35 females. The prevalence of pre-clinical left ventricular diastolic dysfunction was 16% and 14% using the ASE 2009 and 2016 guidelines respectively.

Conclusion: Prevalence enabling accurate sample size calculation for the main study was obtained. Modifications were made to the research proposal based on lessons learnt.

Keywords: Prevalence; pre-clinical; diastolic dysfunction; left ventricular; population; Nigeria, echocardiography.

1. INTRODUCTION

Sample size is the minimum number of subjects needed to answer a research question and it is crucial in the planning and designing of a study. Key components to appropriately estimate sample size are an acceptable level of significance (P value) dependent on alpha, power of the study- complement of beta ($1 - \beta$), effect size defined as the minimum difference that is clinically and biologically credible and population variability of an outcome which is usually unknown but can be estimated from a pilot or previous studies. [1,2] Others include 1-tailed or 2-tailed testing, outcome of interest, study design and allocation ratio [3].

Racial, gender and age differences exist in cardiovascular structure and function, risks of heart failure as well as cardiac response to these risks [4,5]. Black populations have been shown to have a higher arterial afterload, left ventricular (LV) dimension, mass and concentricity [6,7]. Significant differences also exist in the left atrial (LA) dimension, LV ejection fraction (EF), mitral annular velocity, tricuspid regurgitant jet pressure gradient and right ventricular dimension [4,6,8].

High-quality data is insufficient in Africa [9,10]. Evidence based medicine has become patient centered considering individual differences. Therefore, data from the White population may not be applied completely to native black

Africans. To the best of our knowledge, population prevalence on the subject matter is lacking in the African continent. Hence, aim of this pilot study to estimate the sample size for the prevalence of pre-clinical LV diastolic dysfunction (LVDD) in a semi-urban Nigerian population with normal ejection fraction.

2. MATERIALS AND METHODS

2.1 Settings

Neni, a semi-urban population with about 30,000 indigenes and non-indigenes according to a written data from the administrative office of Neni town union in April 2023. It is located in Anaocha Local Government Area (LGA) of Anambra state, Southeastern, Nigeria [11]. The town hosts the headquarters of the LGA, a diagnostic centre of a Federal Teaching Hospital and three primary health care centres.

2.2 Subjects

Fifty persons were recruited consecutively over two days on the 18th and 19th December 2023 at a government primary health care centre after awareness was created in churches and by the town crier.

2.2.1 Exclusion criteria

Included history of pulmonary vascular, parenchymal or obstructive airway disease,

history or evidence of any terminal illness e.g. malignancy, symptoms and signs of heart failure, atrial fibrillation, artificial pacemaker, poor echocardiographic window, ejection fraction less than 52% for males and 54% for females, unreliable determination of diastolic dysfunction, moderate or severe valvular heart disease and cardiomyopathy.

2.3 Data Collection

2.3.1 Clinical history

An interviewer-based questionnaire was administered to obtain the biodata, history of traditional cardiovascular disease risk factors, obstructive airway and heart failure symptoms.

2.3.2 Blood pressure

Blood pressure (BP) was checked using the oscillometric method (Omron M3 HEM-RML31 sphygmomanometer was used) on both arms and the arm with the higher reading was used for a recheck after 30mins.

2.3.3 Anthropometry

Weight and height measurements were taken with the subjects standing on a standard weighing scale and stadiometer with no shoes, heavy clothing, or head gear. Both readings were reported to the nearest kilogram and centimetre. Body mass index (BMI) and body surface area (BSA) were calculated using the Du bois formula.

2.3.4 Electrocardiography

A 12-lead electrocardiography (ECG) was performed by a trained technician using Cardiovit AT-1 Schillers (Switzerland) ECG machine.

2.3.5 Echocardiography

Echocardiography was performed using GE healthcare Vivid iq (China) echocardiography machine with a 3SC transducer. One cardiologist with experience in both Asian and Black populations performed the echocardiography using the American Society of Echocardiography (ASE) guideline,[12] in the standard parasternal long (PLAX), apical five(A5C), four(A4C) and two(A2C) chamber views [13]. A two-dimensional (2D)-guided direct measurement of the interventricular septal wall thickness at end diastole (IVSd), LV internal diameter at end diastole (LVIDd), and the posterior wall thickness

(PWTd) at end diastole measurements were obtained according to ASE guideline [12]. The machine automatically calculated the EF using the Teichholz's formula [13]. EF greater than or equal to 54% and 52% were considered normal for females and males respectively [14]. From the A4C view, using the pulse wave doppler (PWD), the sample volume was placed at the tip of the mitral leaflets. The mitral inflow E and A wave velocities were obtained. The machine automatically gave the E/A ratio and deceleration time (DT) on tracing the slope of the E wave. The A wave duration was taken in milliseconds. Subjects were asked to bear down to assess changes on the mitral E/A ratio during valsalva manoeuvre. Tissue doppler imaging (TDI) was used to obtain the mitral annular velocities with the sample volume placed at the septal and lateral annulus respectively to obtain the septal and lateral e' velocities. Average of the septal and lateral e' was calculated manually. Average E/ e' was obtained by dividing the E velocity with the average sum of the septal and lateral e' velocities. Continuous wave doppler (CWD) with sample volume placed at the vena contracta of the tricuspid regurgitant jet gave the maximum tricuspid regurgitation velocity (TRVmax) from the spectral envelope. Isovolumic relaxation time (IVRT) was obtained from the A5C view with the PWD, sample volume was placed in the LV cavity to obtain both the left ventricular outflow and mitral inflow spectral profiles. Due to technical challenges encountered in the field the pulmonary venous waves were not used for assessment. Left atrial maximum volume index (LAVI) was obtained in the A4C and A2C views at ventricular end systole using the formula: [15]

$$LAVI = \frac{8/3 \times (A1 \times A2)/L}{BSA} = \frac{8.5(A1 \times A2)/L}{BSA}$$

After the left atrial area was traced in both views and assigned A1 and A2 respectively. Length of left atrium (L) was assigned L1 and L2 respectively and was measured as a line drawn from the midpoint (of another line, drawn from the mitral septal and lateral annulus) to the roof of the left atrium (LA). The shorter of L1 and L2 was used in the formula.

2.3.5.1 ASE (2009) guideline for diagnosing diastolic dysfunction [16,17]

Diagnosis of diastolic dysfunction in the presence of normal EF was made when average e' < 9 cm/s and LAVI ≥34ml/m².

2.3.5.2 ASE (2016) guideline for diagnosing diastolic dysfunction [18]

Diagnosis of diastolic dysfunction in the presence of normal EF was made when three out of the four following parameters are present; septal $e^l < 7\text{cm/s}$ or lateral $e^l < 10\text{cm/s}$; Ave E/ $e^l > 14$; LAVI $> 34 \text{ ml/m}^2$; TRVmax $> 280\text{cm/s}$.

2.4 Statistical Analysis

IBM-SPSS version 25.0 (Chicago Illinois, USA) was used for analysis. Frequency tables, percentages, mean, and standard deviation were used to provide a descriptive summary of the study participants. Prevalence was calculated as the total number of subjects with diastolic dysfunction divided by the sample population.

3. RESULTS

Table 1 shows the socio-demographic profile of the respondents. Table 2 shows history of participants' cardiovascular disease risks and symptoms.

Prevalence of left ventricular diastolic dysfunction using the 2009 ASE guideline was 16% as shown in Table 3.

Prevalence of left ventricular diastolic dysfunction using the 2016 ASE guideline was 14% as shown in Table 4.

4. DISCUSSION

Prevalence of LVDD in a general population from previous studies [19,20,21,22] outside the African continent varied from 11.1 – 67.4% depending on the population characteristics and diagnostic criteria. The prevalence in a German study [19] dropped to 3.1% when left atrial enlargement was considered. This may explain the low prevalences in our study, as LAVI was a parameter used in both diagnostic guidelines [18,21,22]. All previous but the German study [19] did not consider the left atrial volume in their diagnostic criteria. A dilated LAVI is a marker of chronically elevated left ventricular filling pressure in the presence of other diastolic parameters and also has prognostic implications. A Tibetan study [21] had a prevalence of 34.39%

Table 1. Socio-demographic characteristics of respondents

Variable	Frequency (n=50)	Percent (%)
Age of respondents in years		
Mean±SD	54.3±14.0	
Age of respondents in groups		
<40 years	7	14.0
40-49 years	7	14.0
50-59 years	17	34.0
≥60 years	19	38.0
Gender		
Male	15	30.0
Female	35	70.0
Marital status		
Single	5	10.0
Married	30	60.0
Divorced	13	26.0
Widowed	2	4.0
Educational attainment of respondent		
No formal education	3	6.0
Primary education	12	24.0
Secondary education	16	32.0
Tertiary education	19	38.0
Employment status		
Unemployed	9	18.0
Self-employed	33	66.0
Paid employment	8	16.0

SD- standard deviation

Table 2. Cardiovascular risk factors and symptoms of participants

Variable	Frequency (n=50)	Percent (%)
History of hypertension		
Yes	24	48
No	26	52
History of diabetes mellitus		
Yes	7	14
No	43	86
History of alcohol use		
Yes	17	34
No	33	66
History of smoking		
Yes	0	0
No	50	100
History of cough		
Yes	1	2.0
No	49	98.0
History of breathlessness		
Yes	12	24
No	38	76
History of leg swelling		
Yes	1	2.0
No	49	98.0

Table 3. Left ventricular diastolic dysfunction using the 2009 ASE guideline

Variable	Frequency (n=50)	Percent (%)
Ave e^l < 9cm/s		
Yes	20	40.0
No	30	60.0
LAVI ≥34ml/m²		
Yes	29	58.0
No	21	42.0
Diastolic dysfunction (2009)		
Present	8	16.0
Absent	42	84.0

LAVI- left atrial maximum volume index; Ave e^l – average sum of the mitral annular septal and lateral velocities

using the 2009 ASE guideline in a population living at a higher altitude than our study participants. Higher altitude is an independent risk factor for LVDD [21,23] and may have contributed to the higher prevalence. Our study excluded subjects with renal dysfunction, coronary artery disease, significant valvular heart disease and cardiomyopathy which a Hispanic study [22] with a high prevalence of 67.4% included.

Limitations of this study are that the sample size calculation is dependent on the value chosen by the investigator(s) which is small therefore, results may differ from large scale studies. Also, causality cannot be established since it is an observational study.

Study is however, feasible with the following recommendations;

1. 1.Data collection should be carried out for nine days with three consecutive days dedicated to each of the three primary health care centres located in the town.
2. 2.Pulmonary venous waveform will not be used in grading diastolic dysfunction as there may be difficulty getting reliable waveforms in the field.
3. 3.Training rehearsals for the data collectors/ interviewers every morning of each study day.

Table 4. Left ventricular diastolic dysfunction using the 2016 ASE guideline

Variable	Frequency (n=50)	Percent (%)
Septal e' < 7cm/s or Lateral e' < 10cm/s		
Yes	28	72.0
No	14	28.0
Ave E/ e' >14		
Yes	1	2.0
No	49	98.0
LAVI >34 ml/m²		
Yes	36	56.0
No	22	44.0
TRVmax > 280cm/s		
Yes	14	28.0
No	36	72.0
Diastolic dysfunction (2016)		
Present	7	14.0
Absent	43	86.0

e' – mitral annular velocity; *LAVI*- left atrial maximum volume index; *Ave E/ e'* – left ventricular filling pressure; *TRVmax*- tricuspid regurgitation velocity maximum

5. CONCLUSION

This pilot study achieved its aim in calculating the sample size for the main study and also proves study's feasibility.

ACKNOWLEDGEMENTS

Special thanks to Chief (Mrs) I.S. Ajulufo and Chief B.A. Madu who obtained approval and acted as middlemen between investigators and the community.

CONSENT

Informed oral and/or written consent were obtained from all participants aged between 18 – 70years, after the Traditional Ruler and President General of the town union gave their permissions.

ETHICAL APPROVAL

Ethical approval was obtained from the Health Research Ethics Committee of the University of Nigeria Teaching Hospital Enugu, Nigeria with reference no NHREC/05/01/2008B-FWA00002458-1RB00002323.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Das S, Mitra K, Mandal M. Sample size calculation basic principles. Indian J Anaesth 2016; 60:652-656.
2. Noordzij M, Tripepi G, Dekker FW, Zoccali C, Tanck MW, Jager KJ. Sample size calculations basic principles and common pitfalls. Nephrol Dial Transplant. 2010;25: 1388-1393
DOI: 10.1093/ndt/gfp732
3. Farrokhyar F, Reddy D, Poolman RW, Bhandari M. Practical tips for surgical research why perform a priori sample size calculation. Can J Surg 2013;56:207-213.
4. LaBounty TM, Bach DS, Bossone E, Koliass TJ. Effect of race on echocardiographic measures of cardiac structure and function. Am J Cardiol. 2019;124:812-818.
DOI: 10.1016/j.amjcard.2019.05.049.
5. Chandra A, Skali H, Claggett B, Solomon SD, Rossi JS, Russell SD. Race and gender-based differences in cardiac structure and function and risk of heart failure. JACC. 2022;79:355-368.
6. Fernandes-Silva MM, Shah AM, Hegde S, Goncalves A, Claggett B, Cheng S. Race-related differences in left ventricular structural and functional remodeling in response to increased afterload the ARIC study. JACC Heart Fail 2017;3:157-165.

7. EchoNoRMAL collaboration. Ethnic-specific normative reference values for echocardiographic LA and LV size, LV mass and systolic function the EchoNoRMAL study. *JACC Cardiovasc Imaging*. 2015;8:656-665.
8. American Heart Association. Size, strength of hearts right side differs by age, gender, race/ethnicity. Available: <http://www.sciencedaily.com/releases/2011/06/110606161024.htm> Assessed 30 March 2024.
9. Marincola E, Kariuki T. Quality research in Africa and why it is important. *ACS Omega* 2020;5:24155–24157. Available: <https://dx.doi.org/10.1021/acsomega.0c04327>
10. Musa SM, Haruna UA, Manirambona E, Eshun G, Ahmad DM, Dada DA. Paucity of health data in Africa an obstacle to digital health implementation and evidence-based practice. *Public Health Rev*. 2023; 44:1-5. DOI: 10.3389/phrs.2023.1605821
11. Neni, Anambra. [Wiki] Available: https://en.wikipedia.org/wiki/Neni,_Anambra#Location Assessed 30 march 2024.
12. Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernade L, et al. Recommendation for cardiac chamber quantification by echocardiography in adults: An update from the American society of Echocardiography and the European Association of Cardiovascular imaging. *J Am Soc Echocardiogr*. 2015; 28:1-39.
13. Houghton AR. Making sense of Echocardiography. 2nd ed. Boca Raton: Taylor and Francis Group. 2014:1-306.
14. Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernade L et al. Recommendation for cardiac chamber quantification by echocardiography in adults: An update from the American society of Echocardiography and the European Association of Cardiovascular imaging. *J Am Soc Echocardiogr*. 2015; 28:1-39.
15. Rao MS. Left Atrial Volume Index (LAVI) in the evaluation of left ventricular diastolic dysfunction. *J Evol Med Dent Sci*. 2015;4:2532-2539. DOI: 10.14260/jemds/2015/365.
16. Kossaify A, Nasr M. Diastolic dysfunction and the new recommendations for echocardiographic assessment of left ventricular diastolic function summary of guidelines and novelties in diagnosis and grading. *J Diagn Med Sonogr*. 2019;35: 317–325. DOI: 10.1177/8756479319836781
17. Nagueh SF, Appleton CP, Gillebert TC, Marino PO, Oh JK, Smiseth OA et al. Recommendation for the evaluation of left ventricular diastolic function by echocardiography. *Eur J Echocardiograph*. 2009;10:165-193.
18. Nagueh SF, Smiseth OA, Appleton CP, Byrd B, Dokanish H, Edvardsen T et al. Recommendations for the evaluation of left ventricular diastolic dysfunction by echocardiography: An update from the American society of echocardiography and the European Association of cardiovascular imaging. *J Am Soc Echocardiograph*. 2016;29:277-314.
19. Fischera M, Baesslera A, Henseb HW, Hengstenberga C, Muscholla M, Holmera S et al. Prevalence of left ventricular diastolic dysfunction in the community results from a doppler echocardiographic-based survey of a population sample. *Eur Heart J*. 2003;24:3 20–328.
20. Kuznetsova T, Herbots H, Lopez B, Jin Y, Richart T, Thijs L et al. Prevalence of left ventricular diastolic dysfunction in a general population. *Circ Heart Fail*. 2009;2: 105-112. DOI: 10.1161/CIRCHEARTFAILURE.108.822627
21. Zheng C, Chen Z, Zhang L, Wang X, Dong Y, Wang J et al. Metabolic risk factors and left ventricular diastolic function in middle-aged Chinese living in the Tibetan plateau. *J Am Heart Assoc*. 2019;8: e010454. DOI: 10.1161/JAHA.118. 010454
22. Rodríguez Castro CE, Lyapin A, Pattathan M, Negrin J, Mukherjee D. Prevalence and predictors of left ventricular diastolic dysfunction in a Hispanic patient population. *Int J Angiol* 2013;22: 229–234.

23. Stembridge M, Ainslie PN, Donnelly J, MacLeod NT, Joshi S, Hughes MG et al. Cardiac structure and function in adolescent Sherpa; effect of habitual altitude and developmental stage. *Am J Physiol Heart Circ Physiol.* 2016;310: H740–H746.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/115616>