



# Isolation and Characterization of Yeasts from Jackfruit and Beet Root Juices

**G. Ashiwini<sup>a\*</sup> and B. Narayanaswamy<sup>a</sup>**

<sup>a</sup> Department of Agricultural Microbiology, University of Agricultural Sciences, Bengaluru-560 065, Karnataka, India.

## **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/CJAST/2022/v41i434003

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92983>

**Original Research Article**

**Received: 06/09/2022**

**Accepted: 08/11/2022**

**Published: 02/12/2022**

## **ABSTRACT**

**Aim:** Yeasts are present as natural microflora on surface of fruits and vegetables that have eminent importance from deputed as biocontrol agents to reduce pathogens of fruits and vegetables to the health benefits of humans.

**Materials and Methods:** Yeasts were isolated using standard plate count technique. Colony morphology of yeasts isolates were characterized under microscope for their cell shapes and sizes. Biochemical characteristics of yeasts were characterized for presence of effervescence in catalase test and production of acid and gas in Yeast Extract Peptone Dextrose (YEPD) broth. Utilization of carbon sources were evaluated by inoculating yeast isolates in YEPD broth with specific carbon source and phenol indicator. Ethanol tolerance were evaluated inoculating yeast isolates in YEPDA broth having different concentrations of ethanol viz., 0%, 5.0%, 10.0%, 12.5% and 15.0%.

**Results:** Population of yeast in jackfruit juices varied from  $20.5 \times 10^3$  to  $70.0 \times 10^3$  cfu per millilitre. Population of yeast varied in beet root juices from  $34.5 \times 10^3$  to  $43.0 \times 10^3$  cfu per millilitre. Juice of Chandra variety of jackfruit recorded the highest yeast population ( $7.0 \times 10^4$ ) when compared to

\*Corresponding author: E-mail: ashwini1592graju@gmail.com;

others. Juice of Detroit red dark variety of beetroot ( $4.3 \times 10^4$ ) recorded highest yeast population ( $4.3 \times 10^4$ ) when compared to others. Yeasts were also isolated from different fruits of which BRTY-I, JKFY-I, POMY-I, WATY-I, APLY-I, MUSKY-I and PINY-I were able to grow in jackfruit and beet root juices. The colonies of yeast varied from white to cream and few were pink. The cell shape varied from oval to ellipsoidal. All the yeast isolates were catalase positive. Most of the yeast isolates produced acid and gas. Yeast isolates were able to utilize glucose, galactose and sucrose. None of the yeast were able to utilize lactose. The highest ethanol tolerant yeasts were JCVY-I, JSVY-III, BDVY-I, PGY, *Saccharomyces boulardii* and *Saccharomyces cerevisiae* var. *ellipsoideus* NCIM 3207.

**Keywords:** Yeasts; utilization of sugars; ethanol tolerance; jackfruit juice; beet root juice.

## 1. INTRODUCTION

Fruits and vegetables provide a suitable niche to several bacteria, yeasts and moulds [1]. The term yeast was originally derived from the Dutch word "gist" which refers to the foam formed during the fermentation of beer wort. Yeast are single celled eukaryotic microorganisms belonging to ascomycetes or basidiomycetes genera of fungi that produce vegetatively by budding or fission and form sexual states which are not enclosed in a fruiting body [2].

Jackfruit (*Artocarpus heterophyllus*) is an ancient and most desirable fruit that is known for its therapeutic and nutritive values. Jackfruit contains vitamins A and C, thiamin, riboflavin, calcium, potassium, iron, sodium, zinc, niacin and has antioxidants [3]. The juicy pulp of the ripe fruit is eaten fresh and has wide potential for preparing different food items due to the presence of protein [4]. Jackfruit is rich in potassium which helps in lowering blood pressure by reversing the effects of sodium. It is good source of vitamin C essential for the production of collagen that gives firmness and strength to the skin and also maintains oral health. Dietary fibres present in jackfruit makes it a good bulk laxative [5].

Beet root (*Beta vulgaris*) on other hand is a vegetable loaded with significant levels of calcium, iron, magnesium, fibres and folate as well as vitamins A and C. [6]. In contrast to other fruits which are rich in fructose, beet roots have more of sucrose and small amounts of glucose and fructose [7]. The juice of beetroot is consumed as a natural remedy for sexual weakness and to reduce kidney and bladder stones. The therapeutic use of beetroot includes its antitumor, carminative, emmenagogue, hemostatic and renal protective properties and is a potential herb used in cardiovascular conditions [8].

Several factors could limit viability and survival in juices. As suggested by Tripathi and Giri [9], the major influencing parameters are intrinsic food parameters, such as pH, titrable acidity, molecular oxygen, water activity, presence of salt, sugar, artificial flavoring, coloring agents, chemical or microbial preservatives like hydrogen peroxide and bacteriocins, processing parameters such as extent of heat treatment, incubation temperature, cooling rate, volume, packaging materials, storage techniques and microbiological factors which mainly includes kind of strains, compatibility of different strains, inoculum proportion and rate. Among all these, pH is one of the chief significant factors affecting the probiotic viability.

Fermentation of fruits and vegetables occur spontaneously on their surfaces. Fermentation of milk, cereals and other substrates are indigenous to many regions of Asia, Africa, Europe, the Middle East and South America to produce beverages with health-promoting properties. Fermentation process increases its shelf life by enhancing nutritional value of substrates, bioavailability of nutrients, enzyme concentration and by neutralizing several toxic substances [10]. It is an effective way to preserve perishable food products. Yeast such as *Saccharomyces cerevisiae* are used as starter cultures in food fermentation. Apart from their use as starter cultures, some have probiotic properties, *Saccharomyces cerevisiae* var. *boulardii* and *Kluyveromyces marxianus* which are added to prepare beverages with therapeutic properties [11].

In developing countries, it is assessed that 50% of production is lost during storage due to infection caused by soft rot bacteria and fungi [12]. Therefore, the excess fruits and vegetables can be processed and nutrient bioavailability can be enhanced. In recent time several raw materials have been

extensively explored to determine if they are appropriate substrates to produce novel non-dairy functional foods (Vasudha and Mishra, 2013).

## 2. MATERIALS AND METHODS

### 2.1 Isolation of Yeasts

Yeasts were isolated from jackfruit and beet root samples collected from nearby villages. Yeasts were also isolated from apple, pomegranate, watermelon, papaya, pineapple and muskmelon obtained from local markets. Healthy and mature jackfruit, beet roots and other fruits was selected. Jackfruit was cut into half and bulbs were deseeded. The beet roots and other fruits were thoroughly washed and cut into thin pieces. All samples were then homogenized to obtain juice and strained using muslin cloth. Standard plate count method was used for isolation of yeasts.

One milliliter (ml) of samples were suspended in 100 ml sterile distilled water blanks. Diluted samples of 0.1 ml were spread on solidified Yeast Extract Peptone Dextrose Agar (YEPDA) media plates for isolation of yeasts. The dilutions from  $10^{-1}$  to  $10^{-3}$  were used in the isolation of yeasts. The petridishes containing yeasts were incubated for 24-48 hours at room temperature. Reference strains of *Saccharomyces cerevisiae*

var. *ellipsoideus* NCIM 3207 and Commercial strain of *Saccharomyces boulardii* (Econorm) were streaked on YEPDA.

The yeast cultures were streaked on Yeast Extract Peptone Dextrose Agar (YEPDA) petriplates to obtain pure cultures. The cultures were also maintained on agar slants for further analysis. The observations regarding colony characteristics were recorded and also compared with the reference and commercial strains. The total soluble solids of the samples (TSS) were also assessed with the help of 'ERMA' hand refractometer having a range of 0-32° Brix at room temperature.

### 2.2 Morphological Characterization of Yeasts

The colony morphology of isolated yeasts were studied by streaking on YEPDA media and incubating it for 48 hours at 28° C. After incubation, the colony morphological characters such as colour, texture and margins were observed as tools for preliminary identification. Cell shape of 1 day old cultures were observed under microscope (40X and 100X) using wet mount method. The observations were recorded as ellipsoidal or oval in shape [13].




Fig. 1. Isolation of yeast from different sources

## 2.3 Biochemical Characterization of Yeasts

### 2.3.1 Catalase test

A drop of yeast culture grown in YEPDA broth was added to a clean grease free slide, on it 0.3% hydrogen peroxide solution was added. Observation of immediate effervescence was recorded. Formation or absence of bubbles were indicated as catalase positive or catalase negative respectively [14].

### 2.4 Utilization of Different Carbon Sources

Yeast isolates were inoculated to test tubes containing yeast extract peptone broth with a specific carbohydrate (Glucose, lactose, mannitol, D-sorbitol and sucrose) and phenol red indicator (pH indicator). Red colour of phenol indicator at neutral pH changes to yellow at pH of 6.8 and below. Durham's tube were also added for detection of gas production. The test tubes were incubated at room temperature and observations were recorded after 48 hours [15].

### 2.5 Ethanol Tolerance

The experiment was conducted to evaluate the ethanol tolerance of yeast isolates at different concentrations of 0 (control), 5, 10 and 15 per cent (v/v). The yeast isolates were grown in YEPDA medium with varying quantities of ethanol with initial cell density of  $2 \times 10^6$  cfu/ml and were incubated at 30 °C. Ethanol tolerance was measured in terms of turbidity (OD values) after different intervals for a period of 24 and 48 hours using spectrophotometer at 600 nm [16].

## 3. RESULTS

### 3.1 Isolation of Yeasts

Jackfruit varieties, beet root varieties and other fruits and vegetables were collected from different locations of Karnataka and local super markets respectively. Yeasts were isolated from the juices of jackfruit, beet root and other fruits and vegetables viz., pomegranate, apple, watermelon, papaya, pineapple and muskmelon (Plate 1) by employing Standard Plate Count Method.

Population of yeast varied from  $4.1$  to  $7.0 \times 10^4$  in jackfruit juice. Population of yeast varied from  $4.1$  to  $4.3 \times 10^4$  in beet root juice. Juice of Chandra variety of jackfruit recorded the highest yeast population of  $7.0 \times 10^4$ . Juice of Detroit red dark

variety of beetroot recorded the highest yeast population of  $4.3 \times 10^4$  when compared to others. The results are presented in Table 1.

### 3.2 Morphological Characterization of Yeast

Colony colour of yeast isolates varied from whitish to cream and few were pink. Whereas, cell shape of yeast varied from oval to ellipsoidal when observed under microscope (100 X). The results are presented in Table 2.

### 3.3 Biochemical Characterization of Yeast

The catalase test was conducted to identify catalase positive and catalase negative yeast isolates. The presence of effervescence is noted as catalase positive (+) and absence of effervescence as catalase negative (-). All the yeast isolates were catalase positive (+). Most of the yeast isolates produced acid and gas except JCVY-IV, JSVY-III, JBVY-I, JBVY-II, BDVY-IV, AY and MMY. The results are presented in Table 3.

All the yeast isolates were able to utilize glucose. Most of the yeast isolates were able to utilize galactose and sucrose. None of the yeast were able to utilize lactose. The results are presented in Table 4. The ethanol tolerance of yeast isolates were tested at different concentrations viz., 0%, 5.0%, 10.0%, 12.5% and 15.0%. The highest ethanol tolerant yeasts were JCVY-I, JSVY-III, BDVY-I, PGY, *Saccharomyces boulardii* and *Saccharomyces cerevisiae* var. *ellipsoideus* NCIM 3207. The results are presented in Table 5.

## 4. DISCUSSION

In the present study, 25 yeast isolates were isolated from different varieties of jackfruit, beet root and other fruits. Yeasts were isolated from the juices of jackfruit, beet root and other fruits. Population of yeast and varied from  $4.1$  to  $7.0 \times 10^4$  in jackfruit juice. Population of yeast varied from  $4.1$  to  $4.3 \times 10^4$  in beet root juice. The microbial population on raw fruits and vegetables are estimated to be  $10^5$  to  $10^7$  CFU/g. Yeasts are most dominant group having population between  $10^2$  to  $10^6$  CFU/g. The most commonly found yeasts genera are *Saccharomyces*, *Candida*, *Pichia*, *Rhodotorula*, *Torulaspora*, *Cryptococcus*, *Hansenula* and *Debaromyces* [17]. Dushyantha et al. [18] isolated yeasts from jackfruit phyllosphere, perianth lobes and juice. Yeast JFY1 was found to be effective strain in jackfruit wine making.

**Table 1. Yeast and lactic acid bacterial (LAB) population on different varieties of jackfruit and beet root**

S. No.	Sources	Name of Varieties	TSS (°Brix)	Yeast population (x10 <sup>4</sup> CFU/ml)	Location
1.	Jackfruit	Chandra	26.5°	7.0	Tubagere, Doddaballapur
		Surya	25.7°	6.2	Hosahalli, Kunigal
		Bhyrachandra	26.2°	6.8	GKVK, Bengaluru
2.	Beet root	Detroit red dark	10.0°	4.3	Sulibele, Hoskote
		Crimson globe	9.2°	4.1	Nandi, Chikkaballapura

**Table 2. Colony morphology and cell shape of yeast isolates**

S.No.	Name of Yeast Isolates	Isolates code	Colony colour	Margin	Cell shape
1.	Jackfruit Chandra variety yeast- I	JCVY-I	Whitish cream	Rough	Oval
2.	Jackfruit Chandra variety yeast- II	JCVY-II	Light creamish	Smooth	Ellipsoidal
3.	Jackfruit Chandra variety yeast- III	JCVY-III	Creamish	Smooth	Ellipsoidal
4.	Jackfruit Chandra variety yeast- IV	JCVY-IV	Light Creamish	Smooth	Ellipsoidal
5.	Jackfruit Surya variety yeast- I	JSVY-I	Whitish	Smooth	Oval
6.	Jackfruit Surya variety yeast- II	JSVY-II	Light creamish	Smooth	Oval
7.	Jackfruit Surya variety yeast- III	JSVY-III	Creamish	Smooth	Ellipsoidal
8.	Jackfruit Bhyrachandra variety yeast- I	JBVY-I	Light creamish	Smooth	Oval
9.	Jackfruit Bhyrachandra variety yeast- II	JBVY-II	Whitish	Smooth	Oval
10.	Jackfruit Bhyrachandra variety yeast- III	JBVY-III	Light creamish	Smooth	Ellipsoidal
11.	Beet root Detroit red dark variety yeast- I	BDVY-I	Creamish	Smooth	Oval
12.	Beet root Detroit red dark variety yeast- II	BDVY-II	Creamish	Smooth	Oval
13.	Beet root Detroit red dark variety yeast- III	BDVY-III	Light creamish	Smooth	Oval
14.	Beet root Detroit red dark variety yeast- IV	BDVY-IV	Whitish	Smooth	Oval
15.	Beet root Crimson globe variety yeast- I	BCVY-I	Creamish	Smooth	Oval
16.	Beet root Crimson globe variety yeast- II	BCVY-II	Light creamish	Smooth	Oval
17.	Beet root Crimson globe variety yeast- III	BCVY-III	Whitish	Rough	Oval
18.	Pomegranate yeast	PMY	Whitish	Smooth	Oval
19.	Water melon yeast	WMY	Creamish	Smooth	Oval
20.	Apple yeast	AY	Creamish	Smooth	Oval
21.	Musk melon yeast	MMY	Whitish	Smooth	Oval
22.	Pineapple yeast	PY	Creamish	Smooth	Oval
23.	Beet root yeast –I	BRY- I	Whitish	Smooth	Oval
24.	Beet root yeast –V	BRY- V	Pinkish	Smooth	Oval
25.	Jackfruit yeast	JFY	Whitish	Smooth	Oval
26.	<i>Saccharomyces boulardii</i> (Probiotic strain)	S. <i>boulardii</i>	Whitish	Smooth	Oval
27.	<i>Saccharomyces cerevisiae</i> var. <i>ellipsoideus</i> NCIM 3207 (Reference strain)	NCIM 3207	Whitish	Rough	Ellipsoidal

**Table 3. Catalase of yeast isolates**

S.No.	Yeast isolates	Catalase Test
1.	JCVY-I	+
2.	JCVY-II	+
3.	JCVY-III	+
4.	JCVY-IV	+
5.	JSVY-I	+
6.	JSVY-II	+
7.	JSVY-III	+
8.	JBVY-I	+
9.	JBVY-II	+
10.	JBVY-III	+
11.	BDVY-I	+
12.	BDVY-II	+
13.	BDVY-III	+
14.	BDVY-IV	+
15.	BCVY-I	+
16.	BCVY-II	+
17.	BCVY-III	+
18.	PMY	+
19.	WMY	+
20.	AY	+
21.	MMY	+
22.	PY	+
23.	BRY- I	+
24.	BRY- V	+
25.	JFY	+
26.	<i>S. boulardii</i> (Probiotic strain)	+
27.	<i>Saccharomyces cerevisiae</i> var. <i>ellipsoideus</i> 3207 (Reference strain)	+

**Note:** '+'= Growth; '-'= No growth

**Table 4. Utilization of different carbon sources by yeast isolates**

S.No.	Yeast Isolates	Glucose		Galactose		Lactose		Mannose		D-Sorbitol		Sucrose	
		A	G	A	G	A	G	A	G	A	G	A	G
1.	JCVY-I	+	+	+	+	-	-	-	-	-	-	+	+
2.	JCVY-II	+	+	-	-	-	-	-	-	-	-	+	+
3.	JCVY-III	+	+	+	+	-	-	-	-	-	-	+	+
4.	JCVY-IV	+	-	+	-	-	-	-	-	-	-	+	-
5.	JSVY-I	+	-	-	-	-	-	-	-	-	-	-	-
6.	JSVY-II	+	+	+	+	-	-	-	-	+	+	-	-
7.	JSVY-III	+	-	+	-	-	-	-	-	-	-	-	-
8.	JBVY-I	+	-	+	-	-	-	+	-	-	-	+	-
9.	JBVY-II	+	-	+	-	-	-	-	-	-	-	+	-
10.	JBVY-III	+	-	+	-	-	-	-	-	-	-	-	-
11.	BDVY-I	+	-	+	-	-	-	-	-	+	+	+	+
12.	BDVY-II	+	+	-	-	-	-	-	-	-	-	+	+
13.	BDVY-III	+	+	+	+	-	-	-	-	-	-	-	-
14.	BDVY-IV	+	-	+	-	-	-	-	-	-	-	-	-
15.	BCVY-I	+	-	+	-	-	-	-	-	-	-	-	-
16.	BCVY-II	+	+	+	+	-	-	-	-	-	-	+	+
17.	BCVY-III	+	+	+	+	-	-	-	-	-	-	-	-
18.	PGY	+	+	-	-	-	-	-	-	-	-	+	+
19.	WMY	+	-	+	-	-	-	-	-	-	-	+	+
20.	MMY	+	-	+	-	-	-	-	-	-	-	+	-

S.No.	Yeast Isolates	Glucose		Galactose		Lactose		Mannose		D-Sorbitol		Sucrose	
		A	G	A	G	A	G	A	G	A	G	A	G
21.	AY	+	-	+	-	-	-	-	-	-	-	-	-
22.	PY	+	+	+	+	-	-	-	-	-	-	+	+
23.	BRY-I	+	+	+	+	-	-	+	+	+	+	+	+
24.	BRY-V	+	-	+	-	-	-	-	-	-	-	+	+
25.	JFY	+	+	+	+	-	-	-	-	+	+	+	+
26.	<i>Saccharomyces boulardii</i> (Probiotic strain)	+	+	+	+	-	-	-	-	-	-	+	+
27.	<i>Saccharomyces cerevisiae</i> var. <i>ellipsoideus</i> 3207 (Reference strain)	+	+	+		-	-	-	-	-	-	+	+

Note: '+'= Growth; '-'= No growth

Table 5. Ethanol tolerance of yeast isolates

S.No.	Yeast isolates	OD value @ 600 nm				
		Ethanol concentration%				
		0%	5.0%	10.0%	12.5%	15.0%
1.	JCVY-I	0.65	0.62	0.59	0.52	0.16
2.	JCVY-II	0.46	0.44	0.37	0.16	0.02
3.	JCVY-III	0.45	0.42	0.38	0.02	0.10
4.	JCVY-IV	0.45	0.43	0.34	0.18	0.06
5.	JSVY-I	0.44	0.42	0.37	0.19	0.14
6.	JSVY-II	0.45	0.43	0.36	0.11	0.09
7.	JSVY-III	0.54	0.46	0.39	0.38	0.13
8.	JBVY-I	0.46	0.43	0.41	0.10	0.06
9.	JBVY-II	0.44	0.42	0.38	0.09	0.05
10.	JBVY-III	0.45	0.43	0.33	0.20	0.07
11.	BDVY-I	0.57	0.55	0.51	0.46	0.14
12.	BDVY-II	0.45	0.44	0.32	0.16	0.03
13.	BDVY-III	0.44	0.43	0.37	0.14	0.09
14.	BDVY-IV	0.45	0.44	0.37	0.13	0.07
15.	BCVY-I	0.44	0.43	0.42	0.12	0.06
16.	BCVY-II	0.45	0.44	0.34	0.23	0.05
17.	BCVY-III	0.47	0.42	0.40	0.12	0.05
18.	PGY	0.55	0.54	0.48	0.43	0.11
19.	WMY	0.54	0.53	0.42	0.12	0.03
20.	MMY	0.55	0.53	0.56	0.44	0.04
21.	AY	0.54	0.52	0.37	0.05	0.11
22.	PY	0.45	0.5	0.42	0.36	0.13
23.	BRY-I	0.44	0.43	0.36	0.35	0.15
24.	BRY-V	0.35	0.34	0.38	0.31	0.05
25.	JFY	0.54	0.52	0.55	0.46	0.13
26.	<i>Saccharomyces boulardii</i> (Probiotic strain)	0.72	0.73	0.63	0.50	0.17
27.	<i>Saccharomyces cerevisiae</i> var. <i>ellipsoideus</i> 3207 (Reference strain)	0.77	0.73	0.75	0.58	0.21

Note: '+'= Growth; '-'= No growth

Colony colour of yeast isolates were mainly white or cream. Whereas, the cell shape of yeast varied from oval to ellipsoidal when observed under microscope (100 X). Ali and Khan [19] reported on seven yeast species from soil

samples, fruits and fermented products and studied their morphological characteristics. Colonies formed by yeast isolates were round, smooth and cream, white to whitish cream colored. Individual cells were oval, elongate and

ovoid to spherical when young and hexagonal when mature.

All the yeast isolates were catalase positive (+). Most of the yeast isolates produced acid and gas except JCVY-IV, JSVY-III, JBVY-I, JBVY-II, BDVY-IV, AY and MMY. Andreolli et al., [20] investigated on catalase activity of 13 basidiomyceteous yeast strains isolated from grapes, apples and relative juices and few identified in previous investigations. The results revealed that catalase activity was maximum in *Papiliotrema flavescens* Yz3 and *Papiliotrema terrestris* Yvp3.

Most of the yeast isolates utilized glucose, galactose and sucrose. None of the yeast were able to utilize lactose. The ethanol tolerant yeasts were JCVY-I, JSVY-III, BDVY-I, PGY, *Saccharomyces boulardii* and *Saccharomyces cerevisiae* var. *ellipsoideus* NCIM 3207 in varying levels of ethanol concentrations viz., 0%, 5.0%, 10.0%, 12.5% and 15.0%. Senapati et al., [21] reported on 12 yeasts species isolated from fruits of local markets of Assam that were catalase positive. The isolates had carbohydrate utilization ability to ferment different sugars namely glucose, lactose and mannitol. All strains were able to ferment glucose and mannitol. Isolates J7 and L1 were observed to be grown in 5%, 10%, 12%, 15% and 18% ethanol concentrations which is a very important character of the isolates to utilize them in fermentation process.

## 5. CONCLUSION

Juices of Chandra variety of jackfruit with 26.5 °Brix and Detroit red dark variety of beetroot with 10.0 °Brix recorded highest yeast population of  $7.0 \times 10^4$  and  $4.3 \times 10^4$  respectively when compared to others varieties. The colonies of yeast isolates varied from white to cream and few were pink. Whereas, the cell shape varied from oval to ellipsoidal. All the yeast isolates were catalase positive. All the yeast isolates were able to utilize glucose. Most of the yeast isolates were able to utilize glucose, galactose and sucrose. None utilized lactose. The ethanol tolerance of yeast isolates were tested and highest ethanol tolerant yeasts were JCVY-I, JSVY-III, BDVY-I, PGY, *Saccharomyces boulardii* and *Saccharomyces cerevisiae* var. *ellipsoideus* NCIM 3207.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Trias R, Bañeras L, Montesinos E, Badosa E. Lactic acid bacteria from fresh fruit and vegetables as biocontrol agents of phytopathogenic bacteria and fungi. *Int Microbiol.* 2008;11(4):231-6. DOI: 10.2436/20.1501.01.66, PMID 19204894.
2. Boekhout T, Kurtzman CP. Principles and methods used in yeast classification, and an overview of currently accepted yeast genera. *Nonconventional yeasts in biotechnology* KLAUS W, editor; 1996; 1-81.
3. Hettiaratchi UPK, Ekanayake S, Welihinda J. Nutritional assessment of jackfruit (*Artocarpus heterophyllus*) meal. *Ceylon Med J.* 2011;56(2):54-8. DOI: 10.4038/cmj.v56i2.3109, PMID 21789865.
4. Elevitch CR, Manner HI. *Artocarpus heterophyllus* (jackfruit). In: *Species profiles for pacific island agroforestry*. Permanent Agriculture Resources series. Honolulu, HI: Western Region Sustainable Agriculture Research and Education; 2006.
5. Ranasinghe RASN, Maduwanthi SDT, Marapana RAUJ. Nutritional and health benefits of jackfruit (*Artocarpus heterophyllus* Lam.): A review. *Int J Food Sci.* 2019;2019:4327183. DOI: 10.1155/2019/4327183, PMID 30723733.
6. Panghal A, Virkar K, Kumar V, B. Dhull SB, Gat Y, Chhikara N. Development of probiotic beetroot drink. *Curr Res Nutr Food Sci.* 2017;5(3):257-62. DOI: 10.12944/CRNFSJ.5.3.10
7. Bavec M, Turinek M, Grobelnik-Mlakar S, Slatnar A, Bavec F. Influence of industrial and alternative farming systems on contents of sugars, organic acids, total phenolic content, and the antioxidant activity of red beet (*Beta vulgaris* L. ssp. *vulgaris* Rote Kugel). *J Agric Food Chem.* 2010;58(22):11825-31. DOI: 10.1021/jf103085p, PMID 20964342.
8. Dambalkar VS, Rudrawar BD, Poojari VR. Study of physico-chemical properties and sensory attributes of beetroot-orange RTS drink. *Int J Sci Res.* 2015;4(10):589-94.
9. Tripathi MK, Giri SK. Probiotic functional foods: survival of probiotics during processing and storage. *J Funct Foods.* 2014;9:225-41.



- DOI: 10.1016/j.jff.2014.04.030
10. Septembre-Malaterre A, Remize F, Poucheret PP. Fruits and vegetables, as a source of nutritional compounds and phytochemicals: changes in bioactive compounds during lactic fermentation. *Food Res Int.* 2018;104:86-99.  
DOI: 10.1016/j.foodres.2017.09.031, PMID 29433787.
  11. Moradi R, Nosrati R, Zare H, Tahmasebi T, Saderi H, Owlia P. Screening and characterization of in vitro probiotic criteria of *Saccharomyces* and *Kluyveromyces* strains. *Iran J Microbiol.* 2018;10(2):123-31. PMID 29997753.
  12. Gunny AAN, Fang LP, Misnan NM, Gopinath SC, Salleh NHM, Hashim RHR, Mat MHC. Microwave-assisted solvent-free extraction of essential oil from *Coleus aromaticus*: anti-phytopathogenic potential for fruit post-harvesting. *Biotech Publishing.* 2021;11:1-11.
  13. Aneja KR. Experiments in microbiology, plant pathology and biotechnology. New age international P. Ltd., New Delhi; 2018.
  14. Balazevic DJ, Ederes GM. Principles of biochemical tests in diagnostic microbiology. New York: Wiley and Co. 1975;405-15.
  15. Nwachukwu IK, Ewelike CN, Ibekwe VI. Isolation, identification and screening of local yeast strains for attributes important for industrial ethanol production. *J Raw Mater Res.* 2015;3(1):38-45.
  16. Khaing TW, Weine N, Mya MO. Isolation, characterization and screening of thermotolerant, ethanol tolerant indigenous yeasts and study on the effectiveness of immobilized cell for ethanol production. *J Sci Technol.* 2008;1:12-4.
  17. Pimentel TC, Gomes de Oliveira LI, de Lourdes Chaves Macedo E, Costa GN, Dias DR, Schwan RF et al. Understanding the potential of fruits, flowers, and ethnic beverages as valuable sources of techno-functional and probiotics strains: current scenario and main challenges. *Trends Food Sci Technol.* 2021;114:25-59.  
DOI: 10.1016/j.tifs.2021.05.024
  18. Dushyantha DK, Raghavendrakumar M, Suvarna VC. Study on fermentation of jackfruit (*Artocarpus heterophyllus* L.) juice by beneficial lactic acid bacteria. *Res J Pharm Biol Chem Sci.* 2011;2:732-9.
  19. Ali MN, Khan MM. Screening, identification and characterization of alcohol tolerant potential bioethanol producing yeasts. *Curr. Res. J Microbiol Biotechnol.* 2014; 2(1):316-24.
  20. Andreolli M, Lampis S, Lorenzini M, Zapparoli G. Features of basidiomycetous yeasts from grapes and apples associated with crop environment and fermenting juice. *J Appl Microbiol.* 2021;131(4): 1932-41.  
DOI: 10.1111/jam.15083, PMID 33759285.
  21. Senapati SS, Saikia D, Bhagawati P. Isolation and characterization of amyolytic yeast strains isolated from indigenous fruits of Assam. India; 2021.

© 2022 Ashiwini and Narayanaswamy; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/92983>